

Europäisches Patentamt

European Patent Office

Office européen des brevets



EP 0 919 542 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication: 02.06.1999 Bulletin 1999/22

(21) Application number: 98122107.0

(22) Date of filing: 23.11.1998

(51) Int. Cl.⁶: **C07C 233/64**, C07C 235/42, C07C 317/32, C07C 323/42, C07D 295/192, A01N 37/18, A01N 43/00, A01N 43/84, A01N 57/00

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU

MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 25.11.1997 JP 339393/97

(71) Applicant: NIHON NOHYAKU CO., LTD. Chuo-ku Tokyo100-0027 (JP)

(72) Inventors:

Tohnishi, Masanori
 Sakai-shi (JP)

Nakao, Hayami
 Kawachinagano-shi (JP)

Kohno, Eiji
 Habikino-shi (JP)

 Nishida, Tateki Tondabayashi-shi (JP) Furuya, Takashi Izumisano-shi (JP)

(11)

Shimizu, Toshiaki
 Kawachinagano-shi (JP)

 Seo, Akira Hashimoto-shi (JP)

Sakata, Kazuyuki
 Kawachinagano-shi (JP)

Fujioka, Shinsuke
 Kawachinagano-shi (JP)

 Kanno, Hideo Ibaraki-shi (JP)

(74) Representative:
Grünecker, Kinkeldey,
Stockmair & Schwanhäusser
Anwaltssozietät
Maximilianstrasse 58
80538 München (DE)

(54) Phthalic acid diamide derivatives, agricultural and horticultural insecticides, and a method for application of the insecticides

(57) The present invention provides a phthalic acid diamide derivative represented by the general formula (I),

$$Z^{1}$$

$$C-N(R^{1}) R^{2}$$

$$Ym$$

$$Z^{2}$$

$$Z^{2}$$

$$(I)$$

wherein R^1 , R^2 and R^3 may be same or different, and are each a hydrogen atom, a cyano group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group, or a group of the formula - A^1 - Q_ℓ ; X may be the same or different and are each a hydrogen atom, a halogen atom, a cyano group, a nitro group, a substituted phenyl group, a naphthyl group, a substituted naphthyl group, a heterocyclic group, a substituted heterocyclic group; or a group of the formula - A^2 - R^7 ; \underline{n} is an integer of 1 to 4; Y may be same or different and are each a hydrogen atom, a halogen atom, a cyano group, a nitro group, a phenyl group, a substituted phenyl group, a naphthyl group, a substituted naphthyl group, a heterocyclic group, a substituted heterocyclic group or a group of the formula - A^2 - R^7 ; \underline{n} is an integer of 1 to 5; Z^1 and Z^2 are each an oxygen atom or a sulfur atom), and an agricultural and horticultural insecticide containing said phthaldiamide derivative, as well as to provide a method for use of said insecticide.

The agricultural and horticultural insecticides of the present invention show excellent activities for controlling injurious insects.

Description

BACKGROUND OF THE INVENTION

5 FIELD OF THE INVENTION

[0001] The present invention relates to phthalic acid diamide derivatives, agricultural and horticultural insecticides containing said derivative as an active ingredient, and a method for application of the insecticides.

O RELATED ART

15

25

30

35

50

55

[0002] Japanese Patent Application Nos. 59-163353 and 61-180753 and J.C.S. Perkin I, 1338-1350, (1978), etc. disclose some of the phthalic acid diamide derivatives of the present invention but neither describe nor suggest their usefulness as agricultural and horticultural insecticides.

SUMMARY OF THE INVENTION

a C₃-C₆ alkynylene group;

[0003] The present inventors earnestly studied in order to develop a novel agricultural and horticultural insecticide, and consequently found that the phthalic acid diamide derivatives represented by the general formula (I) of the present invention are novel compounds not known as agricultural and horticultural insecticides in any literature and that said derivatives including the compounds disclosed in the above references can be used for a new purpose as agricultural and horticultural insecticides. Thus, the present invention has been accomplished.

DETAILED DESCRIPTION OF THE INVENTION

[0004] The present invention relates to phthalic acid diamide derivatives of the general formula (I),

$$\begin{array}{c}
X_{1} \\
C-N(R^{1}) R^{2}
\end{array}$$

$$\begin{array}{c}
Y_{m} \\
Y_{m}
\end{array}$$
(1)

wherein R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a cyano group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group or a group of the formula - A^1 - Q_ℓ (wherein A^1 is -O-, -S-, - SO_2 -, -C(=O)-, a group of the formula - $N(R^4)$ - (wherein R^4 is a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenylcarbonyl group, or a substituted phenylcarbonyl group having at least one substituent which may be the same or different, and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkoxy group, a C_1 - C_6 alkoxy group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 -C

a di- C_1 - C_6 alkylamino group which may be the same or different), a C_1 - C_6 alkylene group, a C_3 - C_6 alkenylene group or

(1) when A^1 is -O- or a group of the formula -N(R^4)-(wherein R^4 is the same as defined above), then Q is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a benefit group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_1 - C_6 alkynyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a mono- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a mono- C_1 - C_6 alkyl group or a substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group having at least one substituted phenyl- C_1 - C_4 alkyl group

uent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a C_1 - C_6 alkynyl group, a C_1 - C_6 alkynyl group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different;

10

15

20

25

30

35

40

45

50

55

(2) when A^1 is -S-, -SO₂- or -C(=O)-, then Q is a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a mono- C_1 - C_6 alkylamino group, a di- C_1 - C_6 alkylamino group which may be the same or different, a C_1 - C_6 alkoxycarbonylamino group, a C_1 - C_6 alkoxycarbonyl-C1-C6 alkylamino group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_2 alkylsulfinyl group, a C_1 - C_2 fonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylamino group, a substituted phenylamino group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a $di-C_1-C_6$ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or a pyrazolyl group), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 -C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different,

(3) when A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C_3 - C_6 alkynylene group, then Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₁-C₆ alkyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a di-C₁-C₆ alkoxyphosphoryl group which may be the same or different, a di-C₁-C₆ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkylyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_2 - C_1 - C_2 - C_3 - C_4 - C_6 fonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-

C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsul fonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, or a group of the formula -Z3-R5 (wherein Z3 is -O-, -S-, -SO-, -SO2- or a group of the formula -N(R6)-(wherein R6 is a hydrogen atom, a C1-C6 alkylcarbonyl group, a halo-C1-C6 alkylcarbonyl group, a C1-C6 alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a phenyl C_1 - C_4 alkoxycarbonyl group, or a substituted phenyl C₁-C₄ alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C₁-C₅ alkylamino group and a di-C₁-C₅ alkylamino group which may be the same or differ-

R⁵ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkylcarbonyl group, a halo C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl C₁-C₄ alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁- C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a C_2 a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different);

ℓ is an integer of 1 to 4); further,

5

10

15

20

25

30

35

40

45

50

55

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X may be the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkenyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6

5

10

15

20

25

30

35

45

50

55

C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁- C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 -Cgroup and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A2-R7 (wherein A2 is -O-, -S-, -SO-, -SO2-, C(=O)-, -C(=NOR8)- (wherein R8 is a hydrogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C3-C6 alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a phenyl-C₁-C₄ alkyl group, or a substituted phenyl-C₁-C₄ alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 -C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different), a C1-C6 alkylene group, a halo-C1-C6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene aroup:

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo C_1 - C_6 alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₃-C₆ cycloalkenyl group, a halo-C₃-C₆ cycloalkenyl group, a di- C_1 - C_6 alkoxyphosphoryl group which may be the same or different, a di- C_1 - C_6 alkoxythiophosphoryl group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 sulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 -C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁- C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, or a group of the formula -A3-R9 (wherein A3 is -C(=O)-, -SO2-, a C1-C6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_3 - C_6 alkynylene group or a halo-C3-C6 alkynylene group,

5

10

15

20

25

30

35

40

45

50

55

(i) when A3 is -C(=O)- or -SO2-, then R9 is a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1- C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 -C₆ alkylamino group which may be the same or different,

(ii) when A^3 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo-C2-C6 alkenylene group, a C3-C6 alkynylene group or a halo-C3-C6 alkynylene group, then R9 is a hydrogen atom, a halogen atom, a cyano group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, or a group of the formula -A4-R10 (wherein A4 is -O-, -S-, -SO-, -SO₂-, -C(=O)-, or a group of the formula -N(R¹¹)- (wherein R¹¹ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1- C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alky sulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the

same or different); and

5

10

15

20

25

30

35

40

45

50

55

 R^{10} is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₃-C₆ cycloalkenyl group, a halo-C₃-C₆ cycloalkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkylnyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different));

(2) when A² is -C(=O)- or a group of the formula -C(=NOR⁸)-(wherein R⁸ is the same as defined above), then R7 is a C1-C6 alkyl group, a halo C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfinyl group, a halo-C1-C6 alkyl sulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having one or more substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 -C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C₁-C₅ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1-C_6 alkyl group, a halo- C_1-C_6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁- C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the

same or different.

5

10

15

20

25

30

35

40

45

50

55

(3) when A² is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, C₂-C₆ alkenylene group, a halo-C₂-C₆ alkenylene group, a C₂-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group, then R⁷ is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 -C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 -C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R¹³)-(wherein R¹³ is a hydrogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a C2-C6 alkynyl group, a C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 -C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkytthio group, a halo-C1-C6 alkytthio group, a C1-C6 alkytsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different); and R¹² is a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkynyl group, a C2-C6 alkynyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C2-C6 alkynyl group, a C2-C6 alkynyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C2-C6 alkynyl group group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl

group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylsulfinyl group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkenyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, or a group of the formula - C_1 - C_6 alkenylene group, a halo- C_2 - C_6 alkylene group, a C_2 - C_6 alkylene group, a C_2 - C_6 alkylene group, a C_2 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group.

15

10

20

25

30

35

40

45

50

55

(i) when A⁶ is -C(=O)- or -SO₂-, then R¹⁴ is a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 nyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different;

(ii) when A^6 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo-C2-C6 alkenylene group, a C2-C6 alkynylene group or a halo-C3-C6 alkynylene group, then R14 is a hydrogen atom, a halogen atom, a cyano group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a C₁-C₆ alkylcarbonyl group, a halo-C₁-C₆ alkylcarbonyl group, a C1-C6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylthio

group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆- alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulf nyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different)));

n is an integer of 1 to 4;

5

10

15

20

25

30

35

40

45

50

55

further, X may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole), by combining together with the adjacent carbon atoms in the phenyl ring, and said condensed ring may have at least one substituents, which may be the same or different, and selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different;

Y is the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a di-C1-C6 alkoxyphosphoryl group which may be the same or different, a di-C1-C6 alkoxythiophosphoryl group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a cyano group, a C_2 - C_6 alkenyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a cyano group, a C_2 - C_6 alkenyl group, a cyano group, a C_2 - C_6 alkenyl group, a cyano group, a C_2 - C_6 alkenyl group, a cyano group, a C_2 - C_6 alkenyl group, a cyano group, a C_2 - C_6 alkenyl group, a C_2 - C_6 a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may he the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A2-R7 (wherein A² and R⁷ are the same as defined above);

m is an integer of 1 to 5;

5

10

15

20

25

30

35

40

45

50

55

further, Y may form a condensed ring (the condensed ring is the same as defined above), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituents, which may be the same or different, and selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group, a sulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein said heterocyclic group is the same as defined above) having at lease one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆

5

10

15

20

proportions.

50

55

alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different; Z^1 and Z^2 are each represents an oxygen atom or a sulfur atom; provided that,

- (1) when X, R^1 and R^3 are hydrogen atoms at the same time; \underline{m} is an integer of 2; Y at 2-position is a fluorine atom and Y at 3-position is a chlorine atom; then R^2 is not ethyl group, isopropyl group, cyclohexyl group, 2-propenyl group, methylthiopropyl group and α -methylbenzyl group,
- (2) when X and R^3 are hydrogen atoms at the same time; \underline{m} is an integer of 2; Y at 2-position is a fluorine atom and Y at 3-position is a chlorine atom; then the 4 to 7 membered ring by combining R^1 and R^2 to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom is not morpholino group,
- (3) when X, R^1 and R^3 are hydrogen atoms at the same time; and R^2 is 1,2,2-trimethylpropyl group; then Y is not a hydrogen atom,
- (4) when X, R¹ and R³ are hydrogen atoms at the same time; R² is 2,2-dimethylpropyl group; and m is an integer of 1; then Y is not 2-ethoxy group, and
- (5) when X, R^1 and R^3 are hydrogen atoms at the same time; and R^2 is <u>tert</u>-butyl group group; and \underline{m} is an integer of 1; then Y is not 4-chlorine atom, 2-nitro group, 4-nitro group, 3-methoxy group, 4-methoxy group and 2,6-dimethyl groups;

agricultural and horticultural insecticides containing as an active ingredient any of the phthalic acid diamide derivatives of the general formula (I) including known compounds; and a method for application of the insecticides.

- [0005] In the definition of the general formula (I) representing the phthalic acid diamide derivative of the present invention, the halogen atom includes chlorine atom, bromine atom, iodine atom and fluorine atom. The term " C_1 - C_6 alkyl" means a linear or branched alkyl group of 1 to 8 carbon atoms, such as methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, n-pentyl, n-hexyl, or the like. The term " C_1 - C_8 alkylene" means a linear or branched alkylene group of 1 to 8 carbon atoms, such as methylene, ethylene, propylene, trimethylene, dimethylmethylene, tetramethylene, i-butylene, dimethylethylene, pentamethylene, hexamethylene, heptamethylene, octamethylene or the like. The term "halo- C_1 - C_6 alkyl" means a substituted and linear or branched alkyl group of 1 to 6 carbon atoms having as the substituent(s) one or more halogen atoms which may be the same or different.
- [0006] As the ring which R¹ and R² form by combining to each other, i.e., the 4- to 7-membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom, there can be exemplified azetidine ring, pyrrolidine ring, pyrrolidine ring, pyrrolidine ring, pyrrolidine ring, piperidine ring, imidazolidine ring, oxazolidine ring, thiazolidine ring, isoxazolidine ring, isothiazolidine ring, tetrahydropyridine ring, piperazine ring, morpholine ring, thiomorpholine ring, dioxazine ring, dithiazine ring, etc. [0007] The phthalic acid diamide derivative of the general formula (I) of the present invention contains an asymmetric carbon atom or some asymmetric center in the structural formula in some cases or has two optical isomers in some cases. The present invention includes these optical isomers and all mixtures containing the optical isomers in arbitrary
- 40 [0008] Preferable examples of each substituent of the phthalic acid diamide derivative of the general formula (I) of the present invention are as follows. Preferable examples of each of R¹ and R² which may be the same or different are hydrogen atom, C₁-C₆ alkyl groups such as methyl, ethyl, i-propyl, etc. Preferable examples of R³ are hydrogen atom, and C₁-C₆ alkyl groups such as methyl, ethyl, n-propyl, i-propyl, n-butyl, etc. Preferable examples of x are halogen atoms, nitro group, halo-C₁-C₆ alkyl groups, halo-C₁-C₆ alkoxy groups, halo-C₁-C₆ alkylthio groups, etc.
 45 examples of Y are halo-C₁-C₆ alkyl groups, halo-C₁-C₆ alkoxy groups, halo-C₁-C₆ alkylthio groups, etc.
 - [0009] The phthalic acid diamide derivative of the general formula (I) of the present invention can be produced, for example, by any of the processes illustrated below.

Production process 1.

[0010]

wherein R1, R2, X, n, Y and m are as defined above.

[0011] A phthalic anhydride derivative of the general formula (V) is reacted with an aniline of the general formula (IV) in the presence of an inert solvent to obtain a phthalimide derivative of the general formula (III). The phthalimide derivative (III) is reacted with an amine of the general formula (II) after or without being isolated, whereby a phthalic acid diamide derivative of the general formula (I-1) can be produced.

(1) General formula (V) \rightarrow general formula (III)

[0012] As the inert solvent used in this reaction, any solvent may be used so long as it does not markedly inhibit the progress of the reaction. There can be exemplified aromatic hydrocarbons such as benzene, toluene, xylene, etc.; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, etc., chlorinated aromatic hydrocarbons such as chlorobenzene, dichlorobenzene, etc.; a cyclic or cyclic ethers such as diethyl ether, dioxane, tetrahydrofuran, etc., esters such as ethyl acetate, etc.; amides such as dimethylformamide, dimethylacetamide, etc.; acids such as acetic acid, etc.; dimethyl sulfoxide; and 1,3-dimethyl-2-imidazolidinone. These inert solvents may be used alone or as a mixture thereof.

[0013] Since the reaction is an equimolar reaction, it is sufficient that the reactants are used in equimolar amounts, though either of them may be used in excess. If necessary, the reaction may be carried out under dehydrating conditions

[0014] As to the reaction temperature, the reaction can be carried out in a temperature range of room temperature to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.

[0015] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced. The desired compound can be subjected to the subsequent reaction without isolation from the reaction solution.

[0016] The phthalic anhydride derivative of the general formula (V) can be produced by the process described in J. Org. Chem., <u>52</u>, 129 (1987), J. Am. Chem. Soc., <u>51</u>, 1865 (1929), J. Am. Chem. Soc., <u>63</u>, 1542 (1941), etc. The aniline of the general formula (IV) can be produced by the process described in J. Org. Chem., <u>29</u>, 1 (1964), Angew. Chem. Int. Ed. Engl., <u>24</u>, 871 (1985), Synthesis, <u>1984</u>, 667, Bulletin of the Chemical Society of Japan, <u>1973</u>, 2351, DE-2606982, JP-A-1-90163, etc.

(2) General formula (III) → general formula (I-1)

[0017] In this reaction, there can be used the inert solvents exemplified above as the inert solvent used in the reaction (1).

[0018] Since the reaction is an equimolar reaction, it is sufficient that the reactants are used in equimolar amounts, though the amine of the general formula (II) may be used in excess.

[0019] As to the reaction temperature, the reaction can be carried out in a temperature range of room temperature to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.

[0020] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced.

Production process 2.

[0021]

15

20

25

$$R^{1}(R^{2})NH$$
 $C-N(R^{1})R^{2}$
 $C-NH$
 $C-NH$

40 wherein R¹, R², n, X, Y and m are as defined above, and X' is a halogen atom or a nitro group, provided that X is other than a hydrogen atom or a nitro group.

[0022] A phthalimide derivative of the general formula (III-1) is reacted with a reactant corresponding to X in the presence of an inert solvent to obtain a phthalimide derivative of the general formula (III). The phthalimide derivative (III) is reacted with an amine of the general formula (II) after or without being isolated, whereby a phthalic acid diamide derivative of the general formula (I-1) can be produced.

(1) General formula (III-1) → general formula (III)

[0023] This reaction can be carried out according to the methods described in J. Org. Chem., <u>42</u>, 3415 (1977), Tetrahedron, <u>25</u>, 5921 (1969), Synthesis, <u>1984</u>, 667, Chem. Lett., <u>1973</u>, 471, J. Org. Chem., <u>39</u>, 3318 (1974), J. Org. Chem., <u>39</u>, 3327 (1974), etc.

(2) General formula (III) → general formula (I-1)

55 [0024] This reaction can be carried out according to production process 1-(2).

Production process 3

[0025]

wherein R1, R2, X, Y, m and n are as defined above.

45

[0026] A phthalic anhydride of the general formula (V-1) is reacted with an aniline of the general formula (IV) in the presence of an inert solvent to obtain a phthalimide derivative of the general formula (III-2). The phthalimide derivative (III-2) is subjected to catalytic reduction with hydrogen after or without isolation to obtain a phthalimide derivative of the general formula (III-3). The phthalimide derivative (III-3) is converted to a phthalimide derivative of the general formula (III) by adding a diazotizing agent and then a metal salt after or without isolation of the phthalimide derivative (III-3). The phthalimide derivative (III) is reacted with an amine of the general formula (II) after or without being isolated, whereby a phthalic acid diamide derivative of the general formula (I-1) can be produced.

(I-1)

- (1) General formula (V-1) → general formula (III-2)
- [0027] The desired compound can be produced by this reaction in the same manner as in production process 1-(1).
- 5 (2) General formula (III-2) → general formula (III-3)
 - [0028] Any solvent may be used in this reaction so long as it does not markedly inhibit the progress of the reaction. There can be exemplified alcohols such as methanol, ethanol, propanol, etc.; acyclic or cyclic ethers such as diethyl ether, dioxane, tetrahydrofuran, etc., and acids such as acetic acid, etc. These inert solvents may be used alone or as a mixture thereof.
 - [0029] As the catalyst for catalytic reduction used in this reaction, there can be exemplified palladium carbon, Raney nickel, palladium black, platinum black, etc. The amount of the catalyst used may be properly chosen in a range of 0.1 to 10% by weight based on the weight of the phthalimide derivative of the general formula (III-2). This reaction is carried out under a hydrogen atmosphere and the hydrogen pressure may be properly chosen in a range of 1 to 10 atmospheric pressure.
 - [0030] As to the reaction temperature, the reaction can be carried out in a temperature range of room temperature to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.
 - [0031] After completion of the reaction, the desired compound is isolated from the reaction mixture containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced. The desired compound can be subjected to the subsequent reaction without isolation from the reaction mixture.
 - (3) General formula (III-3) → general formula (III)
 - [0032] In this reaction, an acidic solvent can be used as an inert solvent. The acidic solvent includes, for example, an aqueous hydrochloric acid solution, an aqueous hydrochloric acid solution, an aqueous hydrochloric acid solution, an aqueous sulfuric acid solution, acetic acid and trifluoroacetic acid. These acidic solvents may be used alone or as a mixture thereof. In addition, these acidic solvents may be used in admixture with ethers such as tetrahydrofuran, dioxane, etc.
 - [0033] The diazotizing agent includes, for example, sodium nitrite, nitrosyl hydrogensulfate and alkyl nitrites. The amount of the diazotizing agent used may be properly chosen in a range of equal amount to excess amount relative to the amount of the phthalimide derivative of the general formula (III-3).
- [0034] As to the reaction temperature, the reaction can be carried out in a temperature range of -50°C to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.
 - [0035] As the metal salt added after the production of a diazonium salt, there can be used, for example, cuprous chloride, cuprous bromide, potassium iodide, copper cyanide, potassium xanthate and sodium thiorate. The amount of the metal salt used may be properly chosen in a range of 1 equivalent to excess equivalents per equivalent of the phthalimide derivative of the general formula (III-3).
 - [0036] After completion of the reaction, the desired compound is isolated from the reaction mixture containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced. The desired compound can be subjected to the subsequent reaction without isolation from the reaction mixture.
- 45 [0037] The reaction can be carried out according to the method described in Org. Synth., IV, 160 (1963), Org. Synth., III, 809 (1959), J. Am. Chem. Soc., 92, 3520 (1970), etc.
 - (4) General formula (III) → general formula (I-1)
- 50 [0038] The desired compound can be produced by this reaction in the same manner as in production process 1-(2).

25

Production process 4.

[0039]

35

50

wherein R1, R2, X, Y, m and n are as defined above.

[0040] A phthalimide derivative of the general formula (III-2) is reacted with an amine of the general formula (II) in the presence of an inert solvent to obtain a phthalic acid diamide derivative of the general formula (I-3). The phthalic acid diamide derivative (I-3) is subjected to catalytic reduction with hydrogen after or without isolation to obtain a phthalic acid diamide derivative of the general formula (I-2). A phthalic acid diamide derivative of the general formula (I-1) can be produced from the phthalic acid diamide derivative (I-2) by adding a diazotizing agent and then a metal salt after or without isolating the phthalic acid diamide derivative (I-2).

(1) General formula (III-2) → general formula (I-3)

[0041] The desired compound can be produced by this reaction in the same manner as in production process 1-(2).

(2) General formula (I-3) → general formula (I-2)

[0042] The desired compound can be produced by this reaction in the same manner as in production process 3-(2).


(3) General formula (I-2) → general formula (I-1)

[0043] The desired compound can be produced by this reaction in the same manner as in production process 3-(3).

Production process 5.

[0044]

55



wherein R¹, R², R³, X, n, Y and m are as defined above.

[0045] A phthalic anhydride derivative of the general formula (V) is reacted with an amine of the general formula (II)

in the presence of an inert solvent to obtain a phthalamide of the general formula (III-4). The phthalamide (III-4) is treated as follows after or without isolation. When R² of the phthalamide (III-4) is a hydrogen atom, the phthalamide (III-4) is condensed into a compound of the general formula (VI) in the presence of a condensing agent, and the compound (VI) is reacted with an aniline of the general formula (IV) in the presence of an inert solvent after or without being isolated. When R² of the phthalamide (III-4) is other than a hydrogen atom, the phthalamide (III-4) is condensed with an aniline of the general formula (IV) in the presence of a condensing agent. Thus, a phthalic acid diamide derivative of the general formula (I) can be produced.

[0046] Alternatively, a phthalic anhydride derivative of the general formula (V) is reacted with an aniline of the general formula (IV) in the presence of an inert solvent to obtain a phthalanilide of the general formula (III-5). The phthalanilide (III-5) is treated as follows after or without isolation. When R³ of the phthalanilide (III-5) is a hydrogen atom, the phthalanilide (III-5) is condensed into a compound of the general formula (VI-1) in the presence of a condensing agent, and the compound (VI-1) is reacted with an amine of the general formula (II) in the presence of an inert solvent after or without being isolated. When R³ of the phthalanilide (III-5) is other than a hydrogen atom, the phthalanilide (III-5) is condensed with an amine of the general formula (II) in the presence of a condensing agent. Thus, a phthalic acid diamide derivative of the general formula (I) can be produced.

- (1) General formula (V) or general formula (VI-1) → general formula (III-4) or general formula (I), respectively
- [0047] The desired compound can be produced by this reaction in the same manner as in production process 1-(2).
- (2) General formula (III-4) or general formula (III-5) → general formula (VI) or general formula (VI-1), respectively
- [0048] The desired compound can be produced by this reaction according to the method described in J. Med. Chem., 10, 982 (1967).
- (3) General formula (VI) or general formula (V) → general formula (I) or general formula (III-5), respectively
- [0049] The desired compound can be produced by this reaction in the same manner as in production process 1-(2).
- 30 (4) General formula (III-4) or general formula (III-5) \rightarrow general formula (I)
 - [0050] The desired compound can be produced by reacting the phthalamide derivative of the general formula (III-4) or the general formula (III-5) with the aniline of the general formula (IV) or the amine of the general formula (II), respectively, in the presence of a condensing agent and an inert solvent. If necessary, the reaction can be carried out in the presence of a base.
 - [0051] The inert solvent used in the reaction includes, for example, tetrahydrofuran, diethyl ether, dioxane, chloroform and dichloromethane. As the condensing agent used in the reaction, any condensing agent may be used so long as it is used in usual amide synthesis. The condensing agent includes, for example, Mukaiyama reagent (e.g. 2-chloro-N-methylpyridinium iodide), 1,3-dicyclohexylcarbodiimide (DCC), carbonyldiimidazole (CDI) and diethyl phosphorocyanidate (DEPC). The amount of the condensing agent used may be properly chosen in a range of 1 mole to excess moles per mole of the phthalamide derivative of the general formula (III-4) or the general formula (III-5).
 - [0052] As the base usable in the reaction, there can be exemplified organic bases such as triethylamine, pyridine, etc. and inorganic bases such as potassium carbonate, etc. The amount of the base used may be properly chosen in a range of 1 mole to excess moles per mole of the phthalamide derivative of the general formula (III-4) or the general formula (III-5).
 - [0053] As to the reaction temperature, the reaction can be carried out in a temperature range of 0°C to the boiling point of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it ranges from several minutes to 48 hours.
- [0054] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced.

55

20

Production process 6

[0055]

5 O || |-C-OR¹⁵ C-OR¹⁵ Halogenation 10 Ç-Hal С-ОН || О (VII) (VII-1) 15 20 (IV) (VII-1)_ 25 (III-6) О ∥ -С-ОН 30 ОΗ (四-6) -35 (111-5) 40 Condensation (III-5) 45

55

50

(VI-1)

$$\begin{array}{c|c}
R^{1}(R^{2}) \text{ NH} & C-N(R^{1}) R^{2} \\
\hline
(VI-1) & C-N H
\end{array}$$
(I-1)

wherein R1, R2, X, n, Y and m are as defined above, Hal is a halogen atom, and R15 is a (C1-C3)alkyl group.

[0056] A phthalic acid ester derivative of the general formula (VII) is halogenated into a phthaloyl halide of the general formula (VII-1) in the presence or absence of an inert solvent. The phthaloyl halide (VII-1) is reacted with an aniline of the general formula (IV) in the presence of an inert solvent and a base after or without being isolated, to obtain a phthalanilide of the general formula (III-6). The phthalanilide (III-6) is hydrolyzed into a phthalanilide of the general formula (III-5) in the presence or absence of an inert solvent after or without being isolated. The phthalanilide (III-5) is condensed into a phthalic anhydride derivative of the general formula (VI-1) after or without being isolated. The phthalic anhydride derivative (VI-1) is reacted with an amine of the general formula (II), whereby a phthalic acid diamide derivative of the general formula (I-1) can be produced.

(1) General formula (VII) → general formula (VII-1)

30

45

[0057] As the inert solvent usable in this reaction, any solvent may be used so long as it does not markedly inhibit the progress of the reaction. There can be exemplified aromatic hydrocarbons such as benzene, toluene, xylene, etc.; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, etc., chlorinated aromatic hydrocarbons such as chlorobenzene, dichlorobenzene, etc.; acyclic or cyclic ethers such as diethyl ether, dioxane, tetrahydrofuran, etc., and esters such as ethyl acetate, etc. These inert solvents may be used alone or as a mixture thereof

[0058] As the halogenating agents, there can be used, for example, thionyl chloride, phosphoryl chloride, and phosphorus trichloride. The amount of the halogenating agent used may be properly chosen in a range of 1 to 10 equivalents per equivalent of the phthalic acid ester of the general formula (VII).

[0059] As to the reaction temperature, the reaction can be carried out in a temperature range of 0°C to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.

[0060] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced. The desired compound can be subjected to the subsequent reaction without isolation from the reaction solution.

[0061] The phthalic acid ester of the general formula (VII) can be produced, for example, by the process described in J. Med. Chem., 31, 1466 (1988).

(2) General formula (VII-1) → general formula (III-6)

[0062] As the inert solvent used in this reaction, there may be used, for example, the inert solvents exemplified in production process 1-(1).

[0063] As the base, an inorganic base or an organic base may be used. As the inorganic base, there may be used, for example, hydroxides of alkali metals, such as sodium hydroxide, potassium hydroxide, etc. As the organic base, there may be used triethylamine, pyridine, etc. The amount of the base used may be properly chosen in a range of 0.5 to 3 equivalents per equivalent of the phthaloyl halide of the general formula (VII-1).

[0064] Since the reaction is an equimolar reaction, it is sufficient that the reactants are used in equimolar amounts, though the amount of the aniline of the general formula (IV) used may be properly chosen in a range of 0.5 to 2 equivalents per equivalent of the phthaloyl halide of the general formula (VII-1).

[0065] As to the reaction temperature, the reaction can be carried out in a temperature range of 0°C to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.

[0066] After completion of the reaction, the desired compound is isolated from the reaction solution containing the

desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced. The desired compound can be subjected to the subsequent reaction without isolation from the reaction solution.

5 (3) General formula (III-6) → general formula (III-5)

[0067] As the inert solvent usable in this reaction, there may be used water, alcohols (e.g. methanol, ethanol and propanol) as water-soluble solvents, and mixed solvents of water and a water-soluble solvent.

[0068] As the base used for the hydrolysis, there may be used, for example, hydroxides of alkali metals, such as sodium hydroxide, potassium hydroxide, etc. The amount of the base used may be properly chosen in a range of 1 to 10 equivalents per equivalent of the phthalanilide of the general formula (III-6).

[0069] As to the reaction temperature, the reaction can be carried out in a temperature range of 0°C to the reflux temperature of the inert solvent used. Although the reaction time is varied depending on the scale of reaction, the reaction temperature, etc., it may be properly chosen in a range of several minutes to 48 hours.

[0070] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be produced. The desired compound can be subjected to the subsequent reaction without isolation from the reaction solution.

20 (4) General formula (III-5) → general formula (VI-1)

[0071] The desired compound can be produced by this reaction according to production process 5-(2).

(5) General formula (VI-1) → general formula (I-1)

25

30

35

40

45

50

55


[0072] The desired compound can be produced by this reaction according to production process 1-(2).

Production process 7.

[0073]

50

55



wherein R1, R2, R3, X, Y, m, n, Z1 and Z2 are as defined above.

30

40

[0074] A benzamide derivative of the general formula (VIII-1) or the general formula (VIII-2) or a thiobenzamide derivative of the general formula (VIII-3) or the general formula (VIII-4) obtained by thiocarbonylation of the benzamide derivative of the general formula (VIII-1) or the general formula (VIII-2), respectively, is subjected to ortho-metallation by using a metal reagent such as butyllithium. The compound thus obtained is directly reacted with an isocyanate or isothiocyanate derivative of the general formula (IX-1) or (IX-2), or the compound is reacted with carbon dioxide to obtain a phthalamide derivative of the general formula (III-4') or the general formula (III-5'), which is treated in the same manner as in production processes 5-(1) to 5-(4). Thus, a phthalic acid diamide derivative of the general formula (I) can be produced.

(1) General formula (VIII-1) or general formula (VIII-2) → general formula (VIII-3) or general formula (VIII-4), respectively

[0075] The desired compound can be produced by this reaction according to the method described in J. Org. Chem., 46, 3558 (1981).

(2) General formula (VIII-1), general formula (VIII-2), general formula (VIII-3) or general formula (VIII-4) → general formula (I)

[0076] In this step, the benzamide derivative of the general formula (VIII-1) or the general formula (VIII-2) or the thiobenzamide derivative of the general formula (VIII-3) or the general formula (VIII-4) obtained by thiocarbonylation of the benzamide derivative of the general formula (VIII-1) or the general formula (VIII-2), respectively, is subjected to ortho-lithiation according to the method described in J. Org. Chem., 29, 853 (1964). The compound thus obtained is reacted with the isocyanate or isothiocyanate derivative of the general formula (IX-1) or (IX-2) at -80°C to room temperature, whereby the desired compound can be produced.

[0077] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be obtained.

(3) General formula (VIII-1), general formula (VIII-2), general formula (VIII-3) or general formula (VIII-4) \rightarrow general formula (III-4) or the general formula (III-5)

[0078] In this step, the desired compound can be produced by carrying out the same ortho-lithiation as in the above step (2) and introducing carbon dioxide into the ortho-lithiation product at -80°C to room temperature.

[0079] After completion of the reaction, the desired compound is isolated from the reaction solution containing the desired compound by a conventional method, and if necessary, purified by recrystallization, column chromatography, etc., whereby the desired compound can be obtained.

(4) General formula (III-4') or general formula (III-5') → general formula (I)

[0080] In this step, the desired compound can be produced in the same manner as in production process 1-(2) or 5-(4).

[0081] Tables 1 and 2 show typical examples of the phthalic acid diamide derivative of the general formula (I) used as the active ingredient of the agricultural and horticultural insecticide of the present invention, but the examples are not intended in any way to limit the scope of the present invention.

General formula (I):

20 [0082]

35

40

45

50

[Table 1] Table 1 $(Z^1, Z^2 = 0)$

10	No	R 1	R ²	Rэ	Хn	Ym	Physical Properties (melting point: °C
	1	Н	H	H	3-NO ₂	2-CH ₃ -5-Cl	173-175
15	2	СНз	Н	H	H	4-CF 3	129-131
	3	СНз	Н	H	3-NO ₂	2-CH ₃ -5-Cl	169-171
•	4	СНз	Н	Н	3-NO2	2-CH ₃ -4-OCHF ₂	167-169
20	5	СНэ	CH 3	H	6-NO ₂	2-CH ₃ -5-Cl	171-173
	6	СНэ	СНз	Н	6-NO2	2-CH ₃ -4-OCHF ₂	167-169
25	7	C2H5	Н	Н	Н	4-CF 3	134-136
-	8	C 2H 5	Н	Н	3-C1	2-CH ₃ -4-OCHF ₂	179-180
	9	C2H5	Н	Н	6-C1	2-CH 3-4-OCHF 2	189-190
30	10	C2H5	Н	Н	3-NO ₂	2-CH3-5-Cl	175-177
	11	C ₂ H ₅	Н	Н	3-NO ₂	2-CH ₃ -4-OCHF ₂	207-208
35							

Table 1 (Cont'd)

5	No	R1	R ²	Ra	V-	V	Physical
		n.	K²	K3	Xn	Ym	Properties (melting point: °C
10	12	C ₂ H ₅	C ₂ H ₅	Н	H	4-CF 3	148-150
	13	C ₂ H ₅	C ₂ H ₅	Н	3-NO ₂	2-CH ₃ -5-Cl	175-177
15	14	n-C3H7	H	Н	H	4-CF ₃	138-140
15	15	n-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-0CHF ₂	171-173
	16	n-C ₃ H ₇	Н	H	6-C1	2-CH ₃ -4-OCHF ₂	189-191
20	17	n-C3H7	Н	Н	3-N0 ₂	2-CH ₃ -4-0CHF ₂	184-186
	18	n-C3H7	Н	H	3-NO ₂	2-CH₃-5-Cl	187-189
	19	n-C3H7	Н	Н	5-CF 3	2,6-(C ₂ H ₅) ₂	230-232
25	20	i-C3H7	H	Н	H	Н	192-194
	21	i-C₃H₁	Н	H	Н	2-NO ₂	198-200
	22	i-C₃H₁	H	H	H	4-NO ₂	139-141
30	23	i-C₃H7	Н	Н	Н	4-F	199-201
	24	i-C₃H₁	H	Н	H	2-CH ₃	191-193
	25	i-C ₃ H ₇	H	Н	H	4-CF 3	198-200
35	26	i-C ₃ H ₇	Н	Н	Н	3-CF ₃	174-176
	27	i-C3H7	H	Н	Н	4-CF 2CF 2CF 3	237-238
40	28	i-C ₃ H ₇	H	H	H	4-(CF ₂) ₃ CF ₃	137-139
	29	i-C ₃ H ₇	H	Н	Н	4-0CF ₃	155-157
	30	i-C3H7	H	Н	Н	4-OCF 2CHFOC 3F 7-n	220-222
45	31	i-C ₃ H ₇	H	H	Н	3-SCF ₃	176-178
	32	i-C3H7	H	Н	H	4-SCHF 2	169-170
	33	i-C ₃ H ₇	H	H	H	4-SCH2CF3	166-167
50	34	i-C ₃ H ₇	H	H	H	4-SCF 2CHF 2	169-170

Table 1 (Cont'd)

5		<u> </u>		T	1		D1
	No	R1	R²	Rз	Xn	Ym	Physical Properties (melting point: °C
10	35	i-C ₃ H ₇	Н	Н	Н	4-S(CF ₂) ₃ CF ₃	159-161
	36	i-C ₃ H ₇	Н	Н	Н	4-SCF(CF ₃) ₂	145-147
15	37	i-C3H7	Н	Н	H	4-SCF ₂ CBrF ₂	158-160
	38	i-C₃H7	Н	H	Н	4-SOCF 2CBrF 2	180-182
	39	i-C3H7	н	Н	Н	4-SO(CF ₂) ₃ CF ₃	192-193
20	40	i-C ₃ H ₇	Н	Н	Н	4-S02CH2CF3	169-170
	41	i-C ₃ H ₇	Н	Н	Н	2,3-Cl ₂	151-153
	42	i-C ₃ H ₇	Н	Н	Н	2,4-Cl ₂	162-164
25	43	i-C₃H₁	Н	Н	Н	3,4-F ₂	172-174
	44	i-C₃H₁	Н	Н	H	2,4-(CH ₃) ₂	162-163
	45	i-C₃H₁	Н	Н	Н	2-C1-4-CF ₃	197-199
30	46	i-C₃H₁	Н	Н	Н	2-C1-4-CF(CF ₃) ₂	201-202
	47	i-C₃H₁	H	Н	Н	2-C1-4-0CF ₃	151-153
35	48	i-C ₃ H ₇	H	н	H	2-Br-4-0CF ₃	146-147
35	49	i-C3H7	H	Н	H	2-CH ₃ -3-Cl	196-198
	50	i-C3H7	H	Н	H	2-CH3-4-Cl	180-182
40	51	i-C ₃ H ₇	H	Н	H	2-CH3-5-Cl	161-162
	52	i-C ₃ H ₇	H	Н	H	2-CH3-4-Br	159-261
	53	i-C ₃ H ₇	H	Н	H	2-CH3-5-F	168-170
45	54	i-C ₃ H ₇	H	H	H	2-CH3-5-C4H8-t	203-204
	55	i-C ₃ H ₇	Н	Н	H	2-CH3-4-CF2CF3	157-159
	56	i-C ₃ H ₇	H	H	H	2-CH3-4-CF2CF2CF3	177-178
50	57	i-C ₃ H ₇	Н	H	H	2-CH ₃ -4-CF(CF ₃) ₂	230-231

Table 1 (Cont'd)

No	5			Γ	Γ	Ι		Physical
To State		No	R1	R 2	Rз	Xn	Ym	Properties
10								
59 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₃ 172-173 60 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHF ₂ 145-146 61 i-C ₃ H ₇ H H H 2-CH ₃ -3-OCF ₂ CHC1F 172-174 62 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHC1F 142-144 63 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHC1F 142-144 64 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CBrF ₂ 164-166 64 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CBrFCF ₃ 151-152 65 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CBrFCF ₃ 163-164 67 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFCCF ₃ 146-148 68 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ CCH ₃ 165-166 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ CCH ₃ 160-162 71 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 163-165 72 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 185-187 40 74 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-C1-PhCH ₂ O) 188-189 45 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-C1-PhCH ₂ O) 188-189 46 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-C1-PhCH ₂ O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H	10	58	i-CaHa	н	н	н	2-CH4-OCHE-	
60 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHF ₂ 145-146 61 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHC1F 172-174 62 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHC1F 142-144 20 63 i-C ₃ H ₇ H H H H 2-CH ₃ -4-CF ₂ CBrF ₂ 164-166 64 i-C ₃ H ₇ H H H B 2-CH ₃ -4-CF ₂ CCl ₂ F 172-173 65 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCF ₂ CHFCF ₃ 151-152 25 66 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCF ₂ CHFCF ₃ 163-164 67 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCH ₂ CH ₃ CH ₃ 165-166 70 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCH ₂ CCH ₃ 165-166 70 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCH ₂ CCH ₃ 165-166 70 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCH ₂ COCH ₃ 163-165 72 i-C ₃ H ₇ H H H B 2-CH ₃ -4-OCH ₂ COOCH ₃ 163-165 72 i-C ₃ H ₇ H H H B 2-CH ₃ -4-CH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 185-187 40 74 i-C ₃ H ₇ H H H B 2-CH ₃ -4-(2-C1-4-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhC) 188-189 45 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhC) 188-189 46 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhC) 188-189 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhC) 188-189 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167		i						
61 i-C ₃ H ₇ H H H H 2-CH ₃ -3-OCF ₂ CHC1F 172-174 62 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHC1F 142-144 20 63 i-C ₃ H ₇ H H H H 2-CH ₃ -4-CF ₂ CBrF ₂ 164-166 64 i-C ₃ H ₇ H H H H 2-CH ₃ -4-CF ₂ CC1 ₂ F 172-173 65 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 151-152 25 66 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 163-164 67 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 178-180 30 69 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ CH ₃ 69 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ CH ₃ 70 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ CH ₃ 71 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ CH ₃ 72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ CH ₃ 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GH ₂ COCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GH ₂ COCH ₃ 121-122 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(G-CH ₂ COCH ₃ 185-187 75 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 185-187 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 188-189 45 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 188-189 46 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167								
62 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHClF 142-144 63 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CBrF ₂ 164-166 64 i-C ₃ H ₇ H H H H 2-CH ₃ -4-CF ₂ CCl ₂ F 1772-173 65 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 151-152 25 66 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 163-164 67 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 178-180 30 69 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ OCH ₃ 165-166 70 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ OCH ₃ 165-166 70 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ SCH ₃ 160-162 71 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 163-165 72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 40 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 188-189 45 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 188-189 46 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167	15							
20 63 i-C ₃ H ₇ H H H H 2-CH ₃ -4-CF ₂ CBrF ₂ 164-166 64 i-C ₃ H ₇ H H H H 2-CH ₃ -4-CF ₂ CBrF ₂ 172-173 65 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 151-152 25 66 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 163-164 67 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 163-164 68 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 178-180 30 69 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ OCH ₃ 165-166 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ SCH ₃ 160-162 71 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ SCH ₃ 163-165 72 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 40 74 i-C ₃ H ₇ H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 40 75 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 45 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 45 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhS) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-C1- 184-185)								
64 i-C ₃ H ₇ H H H 2-CH ₃ -4-CF ₂ CCl ₂ F 172-173 65 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 151-152 25 66 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CBrFCF ₃ 163-164 67 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CBrFCF ₃ 163-164 68 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ CCH ₃ 165-166 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ CCH ₃ 165-166 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ CCH ₃ 160-162 71 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 163-165 72 i-C ₃ H ₇ H H H 2-CH ₃ -4-OCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-(GH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-(GH ₂ COOCH ₃ 150-152 75 i-C ₃ H ₇ H H H 2-CH ₃ -4-(GH ₂ CH ₃ -PhO) 185-187 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(GH ₂ CH ₂ -PhO) 183-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(GH ₂ -CH ₂ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GH ₂ -CH ₂ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GH ₂ -CH ₂ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GH ₂ -CH ₂ -PhO) 185-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GH ₂ -CH ₂ -PhS) 181-182		62	i-C3H7	H	H	H	2-CH ₃ -4-OCF ₂ CHC1F	142-144
65 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 151-152 66 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFCF ₃ 163-164 67 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCF ₂ CHFOCF ₃ 146-148 69 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ OCH ₃ 165-166 70 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ OCH ₃ 160-162 71 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COCH ₃ 163-165 72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 150-152 75 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(2-C1-4-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 45 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-C1- 184-185)	20	63	i-C₃H7	H	H	H	2-CH3-4-CF2CBrF2	164-166
25		64	i-C₃H₁	Н	H	H	2-CH3-4-CF2CCl2F	172-173
67 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CF ₂ CHFOCF ₃ 146-148 68 i-C ₃ H ₇ H H H H 2-CH ₃ -4-SC ₃ H ₇ -i 178-180 69 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ OCH ₃ 165-166 70 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ SCH ₃ 160-162 71 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ SCH ₃ 163-165 72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 150-152 75 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(2-C1-4-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhS) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-C1- 184-185)		65	i-C ₃ H ₇	Н	H	H	2-CH3-4-OCF2CHFCF3	151-152
68	25	66	i-C ₃ H ₇	H	Н	H	2-CH3-4-OCF2CBrFCF3	163-164
30 69 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ 0CH ₃ 70 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ SCH ₃ 160-162 71 i-C ₃ H ₇ H H H H 2-CH ₃ -4-COOCH ₃ 163-165 72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCF ₃ -PhO) 185-187 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCF ₃ -PhO) 150-152 75 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCH ₂ COOCH ₃ 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCH ₂ COOCH ₃ 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCH ₂ COOCH ₃ 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCH ₂ COOCH ₃ 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(GCH ₂ COOCH ₃ 184-185		67	i-C3H7	Н	H	Н	2-CH ₃ -4-OCF ₂ CHFOCF ₃	146-148
70 i-C ₃ H ₇ H H H 2-CH ₃ -4-0CH ₂ OCH ₃ 165-166 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-0CH ₂ SCH ₃ 160-162 71 i-C ₃ H ₇ H H H 2-CH ₃ -4-COOCH ₃ 163-165 72 i-C ₃ H ₇ H H H 2-CH ₃ -4-0CH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 74 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 150-152 75 i-C ₃ H ₇ H H H 2-CH ₃ -4-(2-C1-4-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-C1-PhS) 181-182 78 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-0) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 78 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 70 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 71 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 72 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 73 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 74 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 75 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-C1- 184-185 75 i-C ₃ H ₇ H H H 184-185 76 i-C ₃ H ₇ H H H 184-185 77 i-C ₃ H ₇ H H H H 184-185 78 i-C ₃ H ₇ H H H 184-185 79 i-C ₃ H ₇ H H H 184-185 70 i-C ₃ H ₇ H H H 184-185 70 i-C ₃ H ₇ H H H 184-185 71 i-C ₃ H ₇ H H 184-185 72 i-C ₃ H ₇ H H H 184-185 73 i-C ₃ H ₇ H H H 184-185 74 i-C ₃ H ₇ H H H 184-185 75 i-C ₃ H ₇ H H H 184-185 76 i-C ₃ H ₇ H		68	i-C ₃ H ₇	Н	Н	H	2-CH3-4-SC3H7-i	178-180
71 i-C ₃ H ₇ H H H H 2-CH ₃ -4-COOCH ₃ 72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-OCH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 150-152 75 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(2-C1-4-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhS) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-C1- 184-185	30	69	i-C ₃ H ₇	н	H	H	2-CH3-4-0CH2OCH3	165-166
72 i-C ₃ H ₇ H H H H 2-CH ₃ -4-0CH ₂ COOCH ₃ 121-122 73 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(F ₅ -PhO) 185-187 74 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-CF ₃ -PhO) 150-152 75 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(2-C1-4-CF ₃ -PhO) 183-185 76 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-Ph-CH ₂ O) 188-189 77 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(4-C1-PhS) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-C1- 184-185		70	i-C3H7	Н	H	Н	2-CH3-4-0CH2SCH3	160-162
72	35	71	i-C3H7	H	H	H	2-CH ₃ -4-COOCH ₃	163-165
74 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-CF ₃ -Ph0) 150-152 75 i-C ₃ H ₇ H H H 2-CH ₃ -4-(2-Cl-4-CF ₃ -Ph0) 183-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ 0) 188-189 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-PhS) 181-182 78 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-0) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-Cl- 184-185		72	i-C3H7	Н	H	H	2-CH3-4-0CH2COOCH3	121-122
75 i-C ₃ H ₇ H H H 2-CH ₃ -4-(2-Cl-4-CF ₃ -Ph0) 183-185 76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ 0) 188-189 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-PhS) 181-182 78 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-0) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-Cl- 184-185		73	i-C ₃ H ₇	H	H	Н	2-CH ₃ -4-(F ₅ -PhO)	185-187
76 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-Ph-CH ₂ O) 188-189 77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-PhS) 181-182 78 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-O) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-Cl- 184-185	40	74	i-C3H7	H	Н	H	2-CH ₃ -4-(3-CF ₃ -PhO)	150-152
77 i-C ₃ H ₇ H H H 2-CH ₃ -4-(4-Cl-PhS) 181-182 78 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-0) 165-167 79 i-C ₃ H ₇ H H H H 2-CH ₃ -4-(3-Cl- 184-185		75	i-C ₃ H ₇	H	Н	H	2-CH ₃ -4-(2-Cl-4-CF ₃ -PhO)	183-185
78 i-C ₃ H ₇ H H H 2-CH ₃ -4-(5-CF ₃ -2-Pyi-0) 165-167 79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-Cl- 184-185		76	i-C ₃ H ₇	Н	Н	Н	2-CH ₃ -4-(4-Cl-Ph-CH ₂ 0)	188-189
79 i-C ₃ H ₇ H H H 2-CH ₃ -4-(3-Cl- 184-185	45	77	i-C3H7	H	H	Н	2-CH ₃ -4-(4-Cl-PhS)	181-182
		78	i-C3H7	H .	Н	Н	$2-CH_3-4-(5-CF_3-2-Pyi-0)$	165-167
5-CF ₃ -2-Pyi-0)		79	i-C ₃ H ₇	H	Н	н	2-CH ₃ -4-(3-C1-	184-185
	50						5-CF ₃ -2-Pyi-0)	

Table 1 (Cont'd)

5	No	Rı.	R2	Rз	Χn	Ym	Physical Properties
	110	.		-	****	••	(melting
							point: ℃
10	80	i-C ₃ H ₇	H	H	H	$4-(3-C1-5-CF_3-2-Pyi-S)$	173-175
	81	i-C₃H7	H	H	H	$2-CH_3-4-P=0(0C_2H_5)_2$	134-136
	82	i-C ₃ H ₇	H	H	H	$2-CH_3-4-OP=S(OCH_3)_2$	132-134
15	83	i-C ₃ H ₇	H	Н	H	2-CF ₃ -4-0CHF ₂	147-149
	84	i-C ₃ H ₇	Н	Н	H	3,5-Cl ₂ -4-OCHF ₂	`183-185
20	85	i-C3H7	Н	Н	H	3-N=C(CF ₃)-NH-4	217-218
	86	i-C ₃ H ₇	Н	H	H	3-N=C(CF ₃)-N(CH ₃)-4	171-173
	87	i-C3H7	Н	H	3-C1	4-C ₄ H ₉ -n	169-171
25	88	i-C ₃ H ₇	Н	Н	3-C1	4-C4H9-t	224-226
	89	i-C₃H₁	H	Н	3-C1	4-CF(CF ₃) ₂	198-200
	90	i-C ₃ H ₇	Н	Н	3-C1	4-CF2CF2CF3	203-204
30	91	i-C ₃ H ₇	Н	Н	3-C1	4-(CF ₂) ₃ CF ₃	176-178
	92	i-C3H7	Н	Н	3-C1	4-0CHF 2	205-207
	93	i-C ₃ H ₇	Н	Н	3-C1	4-OCF 2CHFOC 3F 7-n	169-171
35	94	i-C ₃ H ₇	Н	Н	3-C1	4-SCH ₃	231-232
	95	i-C ₃ H ₇	Н	Н	6-C1	4-SCH ₃	193-195
	96	i-C ₃ H ₇	H	Н	3-C1	4-SOCH ₃	178-182
40	97	i-C ₃ H ₇	Н	Н	3-C1	4-S0 ₂ CH ₃	208-210
	98	i-C ₃ H ₇	Н	H	3-C1	4-SCHF 2	220-222
45	99	i-C ₃ H ₇	Н	Н	3-C1	3-SCF 3	189-191
	100	i-C ₃ H ₇	Н	Н	3-C1	3-SOCF 3	183-187
	101	i-C ₃ H ₇	Н	Н	3-C1	4-SCH ₂ CF ₃	191-193
50	102	i-C ₃ H ₇	Н	Н	3-C1	4-SCF 2CHF 2	198-200

Table 1 (Cont'd)

5	No	R 1	R²	R³	Xn	Ym.	Physical Properties (melting point: °C
10	103	i-C ₃ H ₇	Н	Н	3-C1	4-SCF 2CBrF 2	201-203
	104	i-C₃H₁	Н	Н	3-C1	4-SCF(CF ₃) ₂	221-223
	105	i-C₃H₁	Н	Н	3-C1	4-S(CF ₂) ₃ CF ₃	199-200
15	106	i-C ₃ H ₇	Н	Н	3-C1	4-SOCF(CF ₃) ₂	204-206
	107	i-C ₃ H ₇	Н	H	3-C1	4-S02CH2CF3	202-204
20	108	i-C3H7	Н	Н	3-C1	4-S02CF2CHF2	227-230
	109	i-C₃H₁	Н	Н	3-C1	4-COCH ₃	217-219
;	110	i-C3H7	Н	Н	3-C1	4-Ph	215-217
25	111	i-C3H7	Н	Н	3-C1	2,3-Cl ₂	168-169
	112	i-C3H7	Н	H	3-C1	2,4-Cl ₂	190-192
	113	i-C3H7	Н	H	3-C1	2,4-F ₂	188-190
30	114	i-C3H7	Н	H	3-C1	2-C1-4-F	172-173
	115	i-C₃H₁	Н	Н	3-C1	2-F-4-Cl	181-182
	116	i-C₃H₁	H	H	3-C1	2,3,4-F ₃	174-176
35	117	i-C₃H₁	Н	Н	3-C1	2,3-(CH ₃) ₂	187-189
	. 118	i-C₃H₁	Н	Н	3-C1	2-CH₃-3-C1	200-202
40	119	i-C₃H₁	Н	Н	3-C1	2-CH3-4-C1	213-215
₩	120	i-C3H7	Н	H	3-C1	2-CH₃-5-C1	183-185
	121	i-C3H7	Н	H	3-C1	2-CH3-4-Br	210-212
45	122	i-C₃H₁	Н	Н	3-C1	2-CH3-4-I	206-208
	123	i-C₃H₁	Н	Н	3-C1	2-CH ₃ -4-0CH ₃	191-192
	124	i-C₃H₁	H	Н	3-C1	2,3-(CH ₃) ₂ -4-0CH ₃	208-210
50	125	i-C ₃ H ₇	H	Н	3-C1	2-C1-4-CF ₃	156-157

Table 1 (Cont'd)

5	No	R 1	R2	R ³	Xn	Ym	Physical Properties (melting point: °C
10	126	i-C ₃ H ₇	H	Н	3-C1	2-C1-4-CF(CF ₃) ₂	204-206
:	127	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-CF ₃	219-220
	128	i-C3H7	H	H	3-C1	2-CH ₃ -4-CF ₂ CF ₃	199-200
15	129	i-C₃H₁	Н	H	3-C1	2-CH ₃ -4-0CF ₂ CCl ₃	169-171
	130	i-C ₃ H ₇	Н	H	3-C1	2-CH3-4-CF2CF2CF3	214-215
20	131	i-C₃H₁	Н	Н	3-C1	2-CH ₃ -4-CF(CF ₃) ₂	220-222
	132	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-(CF ₂) ₃ CF ₃	188-189
	133	i-C₃H7	Н	Н	3-C1	2-CH ₃ -4-(CF ₂) ₅ CF ₃	161-163
25	134	i-C₃H₁	H	H	3-C1	3-C1-4-0CHF ₂	197-199
	135	i-C3H7	H	Н	3-C1	2-C1-4-0CF ₃	158-159
	136	i-C ₃ H ₇	Н	Н	3-C1	2-Br-4-0CF ₃	169-170
30	137	i-C3H7	Н	Н	3-C1	3-F-4-0CHF ₂	211-212
	138	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-OCHF ₂	193-195
	139	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-OCF ₃	199-201
35	140	i-C ₃ H ₇	Н	H	3-C1	2-CH ₃ -4-OCBrF ₂	181-182
	141	i-C ₃ H ₇	Н	H	3-C1	2-CH ₃ -4-OCF ₂ CHF ₂	202-204
40	142	i-C3H7	Н	н	3-C1	2-CH ₃ -3-OCF ₂ CHC1F	169-171
40	143	i-C ₃ H ₇	Н	н	3-C1	2-CH3-4-OCF2CHC1F	194-196
	144	i-C3H7	H	Н	3-C1	2-CH ₃ -4-OCF ₂ CBrF ₂	193-194
45	145	i-C3H7	н	Н	3-C1	2-CH ₃ -4-OCF ₂ CCl ₂ F	202-203
	146	i-C3H7	н	Н	3-C1	2-CH ₃ -4-OCF ₂ CHFCF ₃	186-187
	147	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-OCH ₂ CF ₂ CHF ₂	207-208
50	148	i-C ₃ H ₇	H	H	3-C1	2-CH ₃ -4-OCF ₂ CBrFCF ₃	205-206
Ĺ							

Table 1 (Cont'd)

5	No	R 1	R ²	R³	Xn	Ym	Physical Properties (melting point: °C
10	149	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-OCF ₂ CHFOCF ₃	179-181
	150	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-0CHF ₂ -5-C1	191-192
	151	i-C₃H₁	H	H	3-C1	3,5-C1 ₂ -4-OCHF ₂	205-207
15	152	i-C₃H₁	Н	Н	3-C1	2-CH ₃ -4-OCF ₂ CHF ₂ -5-Cl	211-212
	153	i-C ₃ H ₇	Н	Н	3-C1	2-CH3-4-SC3H7-i	189-191
20	154	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-SCHF ₂	189-191
20	155	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-SOCHF ₂	173-176
	156	i-C₃H7	Н	Н	3-C1	2-CH3-4-S02CHF2	168-170
25	157	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-(F ₅ -PhO)	224-226
	158	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-(5-CF ₃ -2-Pyi-0)	189-191
	159	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-(3-Cl-	204-205
30						5-CF ₃ -2-Pyi-0)	
	160	i-C3H7	Н	Н	3-C1	4-(3-C1-5-CF ₃ -2-Pyi-S)	213-215
	161	i-C ₃ H ₇	Н	H	3-C1	2-CH ₃ -4-P=0(OC ₂ H ₅) ₂	71-73
35	162	i-C₃H7	H	H	3-C1	2-CH ₃ -4-OP=S(OCH ₃) ₂	168-170
	163	i-C ₃ H ₇	H	H	3-C1	2-CF ₃ -4-0CHF ₂	194-196
40	164	i-C ₃ H ₇	H	H	3-C1	3-CF ₃ -4-0CHF ₂	208-209
	165	i-C ₃ H ₇	H	Н	3-C1	3-N=C(CF ₃)-0-4	248-250
	166	i-C ₃ H ₇	Н	Н	3-C1	3-N=C(CF ₃)-NH-4	194-196
45	167	i-C ₃ H ₇	Н	н	3-C1	3-N=C(CF ₃)-N(CH ₃)-4	225-227
	168	i-C ₃ H ₇	H	н	4-C1	Н	190-192
	169	i-C3H7	н	Н	4-C1	4-F	213-215
50	170	i-C ₃ H ₇	Н	H	4-C1	2-CH ₃	208-210

Table 1 (Cont'd)

5	No	R1	R2	Ra	Xn	Υm	Physical Properties (melting point: °C
10	171	i-C ₃ H ₇	Н	Н	4-C1	3-CF 3	196-198
	172	i-C₃H₁	Н	Н	4-C1	4-0CF ₃	192-194
	173	i-C3H7	Н	Н	4-C1	2,4-Cl ₂	174-176
15	174	i-C₃H₁	Н	н	4-C1	3,4-F ₂	231-233
	175	i-C ₃ H ₇	Н	Н	4-C1	2,3-Cl ₂	186-188
	176	i-C3H7	н	Н	4-C1	2-CH₃-3-C1	203-205
20	177	i-C₃H₁	Н	Н	4-C1	2-CH3-4-C1	206-208
	178	i-C3H7	Н	H	4-C1	2-CH3-5-C1	207-208
25	179	i-C3H7	Н	Н	4-C1	2-CH3-5-F	229-231
	180	i-C ₃ H ₇	Н	Н	4-C1	2-CH3-4-0CHF2	223-224
	181	i-C ₃ H ₇	Н	H	5-C1	Н	186-188
30	182	i-C ₃ H ₇	Н	Н	5-C1	4-F	209-211
	183	i-C3H7	Н	Н	5-C1	2-CH ₃	187-189
	184	i-C ₃ H ₇	Н	Н	5-C1	3-CF 3	198-200
35	185	i-C3H7	Н	Н	5-C1	4-0CF ₃	180-182
	186	i-C3H7	Н	Н	5-C1	2,3-Cl ₂	167-169
	187	i-C3H7	н	Н	5-C1	2,4-Cl ₂	165-167
40	188	i-C3H7	H	Н	5-C1	3,4-F ₂	207-209
	189	i-C ₃ H ₇	Н	Н	5-C1	2-CH ₃ -3-C1	204-206
	190	i-C ₃ H ₇	Н	Н	5-C1	2-CH3-4-C1	202-204
45	191	i-C₃H₁	Н	H	5-C1	2-CH ₃ -5-Cl	209-210
	192	i-C ₃ H ₇	Н	Н	5-C1	2-CH₃-5-F	192-194
50	193	i-C₃H7	Н	Н	5-C1	2-CH ₃ -4-0CHF ₂	188-189

Table 1 (Cont'd)

5	No	R1	R2	R³	Xn	Ym	Physical Properties (melting point: °C
10	194	i-C ₃ H ₇	Н	Н	5-C1	2,3,4-F ₃	224-226
	195	i-C ₃ H ₇	Н	Н	6-C1	4-C₄H₀-n	194-196
	196	i-C ₃ H ₇	Н	Н	6-C1	4-C4H9-t	235-237
15	197	i-C3H7	Н	Н	6-C1	4-CF ₂ CF ₂ CF ₃	216-217
	198	i-C ₃ H ₇	Н	Н	6-C1	4-CF(CF ₃) ₂	209-211
20	199	i-C ₃ H ₇	Н	Н	6-C1	4-(CF ₂) ₃ CF ₃	196-198
	200	i-C3H7	Н	Н	6-C1	4-0CHF 2	223-225
	201	i-C₃H₁	Н	H	6-C1	4-OCF 2 CHFOC 3F 7-n	205-207
25	202	i-C ₃ H ₇	Н	H	6-C1	4-SCH ₂ CF ₃	189-190
	203	i-C₃H₁	H	H	6-C1	4-SCF 2CHF 2	211-213
	204	i-C₃H₁	Н	H	6-C1	4-SCF(CF ₃) ₂	250-252
30	205	i-C₃H₁	H	H	6-C1	4-S(CF ₂) ₃ CF ₃	210-212
	206	i-C ₃ H ₇	H	H	6-C1	3-SOCF 3	212-215
	207	i-C ₃ H ₇	H ·	H	6-C1	4-COCH ₃	230-232
35	208	i-C3H7	H	H	6-C1	2,3-Cl ₂	179-180
	209	i-C ₃ H ₇	H	H	6-C1	2,4-Cl ₂	199-200
40	210	i-C ₃ H ₇	H	H	6-C1	2,4-F ₂	196-198
	211	i-C ₃ H ₇	H	H	6-C1	2-C1-4-F	196-197
	212	i-C ₃ H ₇	Н	н	6-C1	2-F-4-C1	184-186
45	213	i-C ₃ H ₇	Н	H	6-C1	2,3-(CH ₃) ₂	214-216
	214	i-C ₃ H ₇	Н	н	6-C1	2-CH ₃ -4-Cl	233-235
	215	i-C ₃ H ₇	H	Н	6-C1	2-CH ₃ -5-Cl	204-206
50	216	i-C ₃ H ₇	н	н	6-C1	2-CH3-4-Br	242-244

Table 1 (Cont'd)

5	No	R 1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	217	i-C3H7	Н	Н	6-C1	2-CH ₃ -4-I	236-238
	218	i-C ₃ H ₇	Н	Н	6-C1	2-CH ₃ -4-0CH ₃	195-197
45	219	i-C3H7	Н	Н	6-C1	2,3-(CH ₃) ₂ -4-0CH ₃	242-244
15	220	i-C ₃ H ₇	Н	Н	6-C1	2-C1-4-CF ₃	171-172
	221	i-C ₃ H ₇	Н	Н	6-C1	2-CH ₃ -4-CF ₃	234-236
20	222	i-C₃H7	Н	H	6-C1	2-CH ₃ -4-OCF ₂ CCl ₃	169-171
	223	i-C3H7	Н	H	6-C1	2-CH ₃ -4-CF ₂ CF ₃	215-217
	224	i-C ₃ H ₇	Н	H	6-C1	2-CH ₃ -4-CF(CF ₃) ₂	238-240
25	225	i-C ₃ H ₇	Н	H	6-C1	2-CH ₃ -4-(CF ₂) ₃ CF ₃	177-178
	226	i-C ₃ H ₇	Н	H	6-C1	2-CH ₃ -4-(CF ₂) ₅ CF ₃	167-169
	227	i-C ₃ H ₇	Н	H	6-C1	3,5-Cl ₂ -4-OCHF ₂	196-198
30	228	i-C ₃ H ₇	H	H	6-C1	2-CH ₃ -4-OCF ₂ CCl ₂ F	218-220
	229	i-C ₃ H ₇	H	H	6-C1	2-CH3-4-OCF2CBrF2	214-215
	230	i-C ₃ H ₇	H	Н	6-C1	2-CH3-4-OCH2CF2CHF2	212-213
35	231	i-C ₃ H ₇	H	Н	6-C1	2-Cl-4-CF(CF ₂) ₂	212-214
	232	i-C ₃ H ₇	H	Н	6-C1	3-C1-4-0CHF ₂	204-206
40	233	i-C ₃ H ₇	Н	Н	6-C1	3-F-4-0CHF 2	225-227
	234	i-C ₃ H ₇	Н	Н	6-C1	2-C1-4-0CF ₃	161-162
:	235	i-C ₃ H ₇	Н	H	6-C1	2-Br-4-0CF ₃	188-189
45	236	i-C ₃ H ₇	Н	Н	6-C1	2-CH ₃ -4-OCHF ₂	213-215
	237	i-C ₃ H ₇	Н	н	6-C1	2-CH ₃ -4-OCF ₉	212-214
	238	i-C ₃ H ₇	н	н	6-C1	2-CH ₃ -4-OCBrF ₂	195-196
50	239	i-C ₃ H ₇	Н	Н	6-C1	2-CH ₃ -4-OCF ₂ CHF ₂	199-201
ĺ							

Table 1 (Cont'd)

5	No	R1	<u>R</u> 2	R3	Xn		Ym	Physical Properties (melting point: °C
10	240	i-C ₃ H ₇	Н	H	6-C1	2-CH ₃ -3	-OCF 2 CHC 1 F	195-197
	241	i-C ₃ H ₇	Н	H	6-C1	2-CH3-4	-OCF 2 CHC 1 F	204-213
	242	i-C ₃ H ₇	H	H	6-C1	2-CH3-4	-OCF 2 CHF CF 3	199-200
15	243	i-C3H7	Н	H	6-C1	2-CH3-4	-OCF 2 CBrFCF 3	226-227
	244	i-C ₃ H ₇	Н	H	6-C1	2-CH3-4	-OCF 2 CHFOCF 3	210-212
20	245	i-C ₃ H ₇	H	H	6-C1	2-CH₃-4	-OCHF 2-5-C1	234-235
	246	i-C ₃ H ₇	H	H	6-C1	2-CH₃-4	-OCF 2CHF 2-5-C1	230-232
	247	i-C ₃ H ₇	H	H	6-C1	2-CH3-4	-SCHF 2	199-201
25	248	i-C ₃ H ₇	Н	Н	6-C1	2-CH3-4	-(F ₅ -Ph0)	243-245
	249	i-C ₃ H ₇	H	Н	6-Cl	2-CH3-4	-(5-CF ₃ -2-Pyi-0)	116-120
	250	i-C ₃ H ₇	H	H	6-C1	2-CH3-4	-(3-C1-	219-221
30						5-	CF ₃ -2-Pyi-0)	
	251	i-C ₃ H ₇	H	H	6-C1	2-CH₃-4	-P=0(0C ₂ H ₅) ₂	146-147
	252	i-C ₃ H ₇	H	Н	6-C1	2-CH3-4	-OP=S(OCH ₃) ₂	183-185
35	253	i-C ₃ H ₇	H	H	6-C1	2-CF ₃ -4	-OCHF 2	234-236
	254	i-C ₃ H ₇	H	Н	6-C1	3-CF ₃ -4	-OCHF 2	204-205
40	255	i-C ₃ H ₇	Н	Н	6-C1	3-N=C(C	F ₃)-0-4	270-272
	256	i-C ₃ H ₇	Н	H	6-C1	3-N=C(C	F ₃)-NH-4	213-215
	257	i-C ₃ H ₇	H	H	6-C1	3-N=C(C	F ₃)-N(CH ₃)-4	239-241
45								·
	258	i-C ₃ H ₇	H	H	3,6-C	12	2-CH3-4-0CHF2	221-222
	259	i-C ₃ H ₇	H	H	3,6-C	12	2-CH3-4-Cl	234-235
50	260	i-C ₃ H ₇	H	H	3,4,5	,6-Cl4	2-CH3-4-Cl	265-266

Table 1 (Cont'd)

5						T	Physical
	No	R1	R2	Rз	Xn	Ym	Properties (melting point: °C
10	261	i-C3H7	Н	H	3-Br	4-CF ₃	221-223
	262	i-C ₃ H ₇	Н	H	3-Br	4-0CF ₃	208-210
15	263	i-C ₃ H ₇	H	H	3-Br	2,3-(CH ₃) ₂	248-250
	264	i-C3H7	Н	Н	3-Br	2,4-(CH ₃) ₂	223-224
	265	i-C ₃ H ₇	H	Н	3-Br	2,4,6-(CH ₃) ₃	254-255
20	266	i-C ₃ H ₇	Н	Н	3-Br	2-CH3-3-C1	215-217
	267	i-C ₃ H ₇	Н	Н	3-Br	2-CH3-4-Cl	176-178
	268	i-C3H7	Н	Н	3-Br	2-CH₃-5-Cl	196-198
25	269	i-C ₃ H ₇	Н	H	3-Br	2,3-(CH ₃) ₂ -4-Cl	222-224
	270	i-C ₃ H ₇	Н	Н	3-Br	2,4-(CH ₃) ₂ -3-Cl	236-238
30	271	i-C ₃ H ₇	Н	Н	3-Br	2-C ₂ H ₅ -4-Cl	205-207
	272	i-C3H7	Н	Н	3-Br	2-CH3-4-Br	220-222
	273	i-C3H7	Н	Н	3-Br	2,3-(CH ₃) ₂ -4-Br	200-202
35	274	i-C ₃ H ₇	H	H	3-Br	2-CH3-4-I	203-205
	275	i-C ₃ H ₇	Н	Н	3-Br	2-CH3-4-F	223-224
	276	i-CáH7	H	H	3-Br	2-C1-4-CF ₃	156-157
40	277	i-C ₃ H ₇	Н	Н	3-Br	2-CH ₃ -4-CF ₃	227-228
	278	i-C ₃ H ₇	Н	Н	3-Br	2-CH ₃ -4-CF ₂ CF ₃	201-202
45	279	i-C ₃ H ₇	Н	H	3-Br	2-CH3-4-CF2CF2CF3	199-200
	280	i-C ₃ H ₇	Н	Н	3-Br	2-CH ₃ -4-CF(CF ₃) ₂	222-224
	281	i-C ₃ H ₇	H	Н	3-Br	2-CH ₃ -4-(CF ₂) ₃ CF ₃	190-191
50	282	i-C ₃ H ₇	Н	H	3-Br	2-CH ₃ -4-OCH ₃	199-200
	1						

Table 1 (Cont'd)

5	No	R1	R2	R3	Xn	Ym	Physical Properties (melting
10							point: ℃
10	283	i-C3H7	H	Н	3-Br	2-CH ₃ -4-0CH ₂ CF ₂ CHF ₂	206-207
	284	i-C₃H7	H	Н	3-Br	2,4-(CH ₃) ₂ -3-0CHF ₂	187-189
15	285	i-C₃H₁	Н	Н	3-Br	2,3-(CH ₃) ₂ -4-OCH ₃	206-208
,,	286	i-C3H7	Н	H	3-Br	2-C1-4-OCF ₃	165-167
	287	i-C ₃ H ₇	Н	H	3-Br	2-Br-4-0CF ₃	179-180
20	288	i-C ₃ H ₇	Н	H	3-Br	2-CH ₃ -4-0CHF ₂	205-207
	289	i-C ₃ H ₇	Н	Н	3-Br	2-CH ₃ -4-0CF ₃	211-213
	290	i-C ₃ H ₇	н	Н	3-Br	2-CH3-4-0CBrF2	178-180
25	291	i-C ₃ H ₇	Н	Н	3-Br	2-CH3-4-OCF2CHFCF3	196-197
	292	i-C ₃ H ₇	H	H	3-Br	2-CH3-4-OCF2CHC1F	194-195
	293	i-C ₃ H ₇	H	H	3-Br	2-CH ₃ -4-OCF ₂ CHF ₂	205-207
30	294	i-C ₃ H ₇	H	H	3-Br	2-CH ₃ -3-Cl-4-OCHF ₂	229-230
	295	i-C ₃ H ₇	H	Н	3-Br	2,3-(CH ₃) ₂ -4-OCHF ₂	219-220
	296	i-C ₃ H ₇	H	H	3-Br	2-CH ₃ -4-SCH ₃	215-217
35	297	i-C ₃ H ₇	H	H	3-Br	2-CH ₃ -4-(3-CF ₃ -Ph0)	156-158
	298	i-C ₃ H ₇	H	H	3-Br	2-CH ₃ -4-(3-Cl-	206-208
40						5-CF ₃ -2-Pyi-0)	
***	299	i-CaH7	H	H	3-Br	2-CH ₃ -4-(5-CF ₃ -	182-184
						2-Pyi-0)	
45	300	i-C ₃ H ₇	Н	Н	3-Br	-3-0CH ₂ 0-4-	195-198
	301	i-C ₃ H ₇	Н	Н	6-Br	4-CF ₃	190-192
	302	i-CaH7	Н	Н	6-Br	4-0CF ₃	210-212
50	303	i-C3H7	н	н	6-Br	2,3-(CH ₃) ₂	250-252

Table 1 (Cont'd)

5	No	Ri	R2	R3		·	Physical
	NO	π.	K²	K.	Xn	Ym	Properties (melting point: °C
10	304	i-C ₃ H ₇	Н	Н	6-Br	2,4,6-(CH ₃) ₃	272-274
	305	i-C ₃ H ₇	Н	Н	6-Br	2-CH ₃ -3-C1	214-216
	306	i-C ₃ H ₇	Н	H	6-Br	2-CH ₃ -4-Cl	198-200
15	307	i-C ₃ H ₇	Н	H	6-Br	2-CH3-5-Cl	194-196
	308	i-C₃H₁	Н	Н	6-Br	2,3-(CH ₃) ₂ -4-Cl	227-229
20	309	i-C ₃ H ₇	Н	Н	6-Br	2,4-(CH ₃) ₂ -3-Cl	249-251
	310	i-C₃H₁	Н	Н	6-Br	2-C ₂ H ₅ -4-Cl	243-245
	311	i-C₃H7	Н	Н	6-Br,	2-CH ₃ -4-Br	227-228
25	312	i-C ₃ H ₇	Н	Н	6-Br	2,3-(CH ₃) ₂ -4-Br	209-211
	313	i-C3H7	Н	H	6-Br	2-CH ₃ -4-I	227-229
	314	i-C ₃ H ₇	Н	H	6-Br	2-CH ₃ -4-F	231-232
30	315	i-C3H7	Н	Н	6-Br	2-C1-4-CF ₃	169-170
	316	i-C ₃ H ₇	Н	H	6-Br	2-CH ₃ -4-CF ₃	232-234
25	317	i-C ₃ H ₇	H	H	6-Br	2-CH ₃ -4-CF(CF ₃) ₂	236-238
35	318	i-C ₃ H ₇	Н	Н	6-Br	2-CH ₃ -4-(CF ₂) ₃ CF ₃	208-210
	319	i-C ₃ H ₇	H	Н	6-Br	2-CH3-4-OCH2CF2CHF2	209-211
40	320	i-C ₃ H ₇	H	Н	6-Br	2,4-(CH ₃) ₂ -3-OCHF ₂	247-249
	321	i-C ₃ H ₇	H	Н	6-Br	2,3-(CH ₃) ₂ -4-OCH ₃	250-252
	322	i-C ₃ H ₇	H	Н	6-Br	2-CH ₃ -4-OCH ₃	220-222
4 5	323	i-C ₃ H ₇	H	н	6-Br	2-C1-4-0CF ₃	182-183
	324	i-C ₃ H ₇	H	н	6-Br	2-Br-4-0CF ₃	195-196
	325	i-C3H7	Н	Н	6-Br	2-CH ₃ -4-OCHF ₂	225-226
50	326	i-C3H7	H	H	6-Br	2-CH ₃ -4-OCF ₃	223-225

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ī	Ym	Physical
	NO	n.	W.	L.	All	İ	IM	Properties (melting
	ļ							point: °C
10	327	i-C₃H₁	H	Н	6-Br	2-0	CH3-4-OCBrF2	194-196
	328	i-C₃H₁	Н	H	6-Br	2-0	CH 3-4-OCF 2 CHFCF 3	212-213
15	329	i-C ₃ H ₇	Н	Н	6-Br	2-0	CH3-4-OCF2CHC1F	211-213
	330	i-C ₃ H ₇	H	Н	6-Br	2-0	CH 3-4-OCF 2 CHF 2	214-215
	331	i-C3H7	H	H	6-Br	2,3	3-(CH ₃) ₂ -4-0CHF ₂	228-229
20	332	i-C₃H7	Н	Н	6-Br	2-0	CH 3 - 3 - C1 - 4 - OCHF 2	224-225
	333	i-C₃H7	Н	Н	6-Br	2-0	CH3-4-SCH3	215-217
	334	i-C3H7	Н	H	6-Br	2-0	CH3-4-(3-CF3-Ph0)	194-195
25	335	i-C ₃ H ₇	Н	H	6-Br	2-0	CH3-4-(5-CF3-	201-203
							-2-Pyi-0)	
	336	i-C ₃ H ₇	H	H	6-Br	2-0	H₃-4-(3-C1-5-	234-236
30							CF ₃ -2-Pyi-0)	
	337	i-C3H7	H	H	6-Br	-3-	-OCH 2O-4-	205-207
25	338	i-C ₃ H ₇	Н	H	3,4-Br ₂	2-0	CH 3 - 4 - OCHF 2	196-197
35	339	i-C ₃ H ₇	Н	Н	3,4-Br ₂	2-0	CH3-4-C1	199-201
	340	i-C ₃ H ₇	Н	Н	3,6-Br	2-0	CH 3 - 4 - OCHF 2	233-234
40	341	i-C ₃ H ₇	H	Н	3,6-Br ₂	2-0	CH3-4-C1	245-247
	342	i-C ₃ H ₇	Н	Н	5,6-Br ₂		2-CH3-4-OCHF2	208-210
	343	i-C ₃ H ₇	Н	н	5,6-Br ₂		2-CH3-4-Cl	259-261
45	344	i-C ₃ H ₇	Н	Н	3,4,5,6-B	r4	2-CH3-4-C1	270-272
	345	i-C ₃ H ₇	H	H	3-I		4-C1	230-232
	346	i-C ₃ H ₇	H	Н	3-I		4-Br	251-253
50	347	i-C ₃ H ₇	H	H	3-I		4-I	231-233
į								

Table 1 (Cont'd)

5	No	B1	R²	R³	Xn	Ym	Physical Properties (melting point: °C
10	348	i-C₃H₁	H	H	3-I	3-CF 3	194-197
	349	i-C ₃ H ₇	Н	H	1-E	4-CF 3	223-224
	350	i-C ₃ H ₇	Н	Н	3-I	4-CF 2 CF 2 CF 3	217-219
15	351	i-C₃H7	Н	Н	3-I	4-CF(CF ₃) ₂	209-211
	352	i-C ₃ H ₇	Н	Н	3-I	4-0CF ₃	222-223
20	353	i-C ₃ H ₇	Н	Н	3-I	4-OCF 2 CHFOCF 3	192-194
20	354	i-C ₃ H ₇	Н	Н	3-I	4-SCHF ₂	204-206
•	355	i-C ₃ H ₇	н	Н	3-I	4-SCH ₂ CF ₃	195-197
25	356	i-C₃H₁	- Н	Н	3-I	4-SCF 2CHF 2	196-198
	357	i-C3H7	Н	H	3-I	4-SCF ₂ CBrF ₂	203-205
	358	i-C₃H₁	Н	Н	3-I	4-SCF(CF ₃) ₂	170-172
30	359	i-C3H7	Н	Н	3-1	4-S(CF ₂) ₃ CF ₃	185-187
	360	i-C3H7	Н	Н	3-I	3,4-F ₂	227-229
	361	i-C₃H₁	Н	Н	3-I	2-CH ₃ -3-Cl	222-224
35	362	i-C3H7	Н	H	3-I	2-CH ₃ -4-Cl	215-217
	363	i-C ₃ H ₇	Н	H	3-I	2-CH ₃ -5-Cl	210-212
	364	i-C ₃ H ₇	Н	Н	3-I	2,4-(CH ₃) ₂ -3-Cl	226-228
40	365	i-C ₃ H ₇	Н	Н	3-I	2,3-(CH ₃) ₂ -4-Cl	235-237
	366	i-C ₃ H ₇	Н	Н	3-I	2-CH ₃ -4-Br	227-229
45	367	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-I	201-203
	368	i-C ₃ H ₇	Н	Н	3-I	2-CH ₃ -4-F	227-228
	369	i-C ₃ H ₇	Н	Н	3-I	2-C1-4-CF ₃	170-171
50	370	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -3-CF ₃	179-181
•							

Table 1 (Cont'd)

5	No	R1	R2	R ³	Xn	Ym	Physical Properties (melting point: °C
10	371	i-C ₃ H ₇	Н	H	3-I	2-CH ₃ -4-CF ₃	202-203
	372	i-C3H7	H	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	195-196
	373	i-C3H7	H	Н	3-I	2-CH ₃ -4-CF ₂ CF ₂ CF ₃	193-195
15	374	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-CF(CF ₃) ₂	211-213
	375	i-C ₃ H ₇	H.	H	3-I	2-CH ₃ -4-(CF ₂) ₃ CF ₃	203-204
20	376	i-C ₃ H ₇	H	H	3-I	2-CH3-4-0CH3	204-206
	377	i-C ₃ H ₇	H	H	3-I	2-CH3-4-0-C3H7-i	209-211
	378	i-C ₃ H ₇	H	H	3-I	2,3-(CH ₃) ₂ -4-0CH ₃	220-222
25	379	i-C ₃ H ₇	H	H	3-I	2-CH3-4-0CH2CF3	223-224
	380	i-C ₃ H ₇	H	H	3-I	2-CH3-4-OCF2CBrF2	228-230
	381	i-C ₃ H ₇	H	H	3-I	2-CH3-4-0CF2CC12F	230-231
30	382	i-C ₃ H ₇	H	H	3-I	3-F-4-0CHF 2	208-210
	383	i-C ₃ H ₇	H	Н	3-I	3,5-Cl ₂ -4-OCHF ₂	234-236
	384	i-C ₃ H ₇	H	H	3-I	3-0CH3-4-0CHF2	196-198
35	385	i-C ₃ H ₇	H	H	3-I	3,4-(OCHF ₂) ₂	171-172
	386	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-0CF ₃	214-216
40	387	i-C ₃ H ₇	H	H	3-I	2-CH 3-4-0CHF 2	207-209
7 0	388	i-C ₃ H ₇	H	H	1-8	2-CH ₃ -4-0CH ₂ CF ₂ CHF ₂	229-231
	389	i-C ₃ H ₇	H	H	1-E	2-CH3-4-0CBrF2	181-182
45	390	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-0CF ₂ CHF ₂	197-199
	391	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-OCF ₂ CHF ₂ -5-C1	198-200
	392	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-OCF ₂ CHC1F	200-201
50	393	i-C ₃ H ₇	Н	н	3-I	2-CH ₃ -4-OCF ₂ CHFCF ₃	213-214

Table 1 (Cont'd)

5	No	R 1	R2	Rз	Xn	Ym	Physical Properties (melting
							point: °C
10	394	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-OCF ₂ CBrFCF ₃	233-234
	395	i-C₃H7	H	H	3-I	2-CH ₃ -4-OCF ₂ CHFOCF ₃	213-215
15	396	i-C₃H7	H	H	3-I	2-CH ₃ -4-OCHF ₂ -5-Cl	230-232
,,	397	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-(F ₅ -Ph0)	245-247
	398	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-(3-CF ₃ -Ph0)	168-170
20	399	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-(5-CF ₃ -2-	186-188
						Pyi-0)	
:	400	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-(3-Cl-5-CF ₃ -	212-214
25						2-Pyi-0)	
	401	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-SO ₂ CH ₃	172-175
	402	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-SC ₃ H ₇ -i	190-192
30	403	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-SCF ₂ CF ₂ CF ₃	227-228
	404	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-(4-Cl-PhS)	191-192
	405	i-C ₃ H ₇	H	Н	3-I	4-(3-C1-5-CF ₃ -2-Pyi-S)	198-200
35	406	i-C ₃ H ₇	H	Н	3-I	2-Br-4-0CF ₃	196-198
	407	i-C ₃ H ₇	H	Н	3-I	2-C1-4-CF2CF2CF3	162-164
40	408	i-C3H7	H	Н	3-I	2-C1-4-0CF ₃	173-175
	409	i-C3H7	H	Н	3-I	2-CF 3-4-0CHF 2	219-220
	410	i-C3H7	H	H	3-I	3-CF ₃ -4-0CHF ₂	128-130
45	411	i-C ₃ H ₇	Н	Н	6-I	4-C1	251-253
	412	i-C3H7	Н	Н	6-I	4-Br	270-272
	413	i-C ₃ H ₇	H	Н	6-I	4-I	242-244
50	414	i-C3H7	Н	Н	6-I	3-CF ₃	210-212

Table 1 (Cont'd)

5					Τ	T	D
	No	R1	R2	Ra	Xn	Ym	Physical Properties (melting point: °C
10	415	i-C ₃ H ₇	Н	H	6-I	4-CF ₃	201-202
	416	i-C₃H₁	Н	H	6-I	4-CF(CF ₃) ₂	238-240
15	417	i-C3H7	H	Н	6-I	4-CF 2CF 2CF 3	238-240
	418	i-C3H7	H	Н	6-I	4-0CF ₃	193-194
	419	i-C ₃ H ₇	Н	Н	6-I	4-OCF 2 CHFOC 3 F 7-n	213-214
20	420	i-C₃H7	Н	Н	I-9	4-SCH ₂ CF ₃	217-219
	421	i-C₃H₁	H	Н	6-I	4-SCHF ₂	224-226
	422	i-C ₃ H ₇	H	Н	6-I	4-SCF 2 CHF 2	213-215
25	423	i-C ₃ H ₇	H	H	6-I	4-SCF ₂ CBrF ₂	220-222
	424	i-C ₃ H ₇	H	H	6-I	4-SCF 2CF 2CF 3	196-197
	425	i-C ₃ H ₇	Н	H	6-I	4-SCF(CF ₃) ₂	216-218
30	426	i-C3H7	Н	H	6-I	4-S(CF ₂) ₃ CF ₃	201-203
	427	i-C3H7	Н	H	6-I	2-CH ₃ -3-Cl	252-254
	428	i-C3H7	H	H	6-I	2-CH ₃ -4-Cl	244-246
35	429	i-C3H7	H	H	6-I	2,4-(CH ₃) ₂ -3-Cl	260-262
	430	i-C3H7	Н	H	6-I	2-CH3-4-Br	241-243
40	431	i-C ₃ H ₇	H	H	6-I	2-CH3-4-I	213-215
•	432	i-C ₃ H ₇	H	H	6-I	2-CH ₃ -4-F	251-252
	433	i-C3H7	Н	H	6-I	2-C1-4-CF ₃	195-196
45	434	i-C3H7	Н	H	6-I	2,3-(CH ₃) ₂ -4-Cl	253-255
*	435	i-C3H7	Н	H	6-I	2-CH ₃ -3-CF ₃	245-251
	436	i~C3H7	н	Н	6-I	2-CH3-4-CF3	220-221
50	437	i-C ₃ H ₇	H	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	203-205

Table 1 (Cont'd)

5	No	R 1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	438	i-C ₃ H ₇	H	H	6-1	2-CH ₃ -4-CF ₂ CF ₂ CF ₃	154-156
	439	i-C3H7	Н	Н	6-I	2-CH ₃ -4-CF(CF ₃) ₂	237-239
	440	i-CaH7	H	K	1-6	2-CH ₃ -4-(CF ₂) ₃ CF ₃	168-170
15	441	i-C ₃ H ₇	H	H	6-I	2-CH ₃ -4-OCH ₃	215-217
	442	i-C₃H7	H	Н	6-I	2-CH ₃ -4-0-C ₃ H ₇ -i	212-214
20	443	i-C3H7	H	Н	6-I	2-CH ₃ -4-0CH ₂ CF ₃	233-234
	444	i-C3H7	H	н	6-I	2-CH3-4-OCF2CBrF2	242-244
:	445	i-C3H7	H	H	6-I	2-CH ₃ -4-OCF ₂ CCl ₂ F	251-253
25	446	i-C ₃ H ₇	Н	Н	6-I	2-CH ₃ -4-OCF ₂ CBrFCF ₃	251-253
	447	i-C ₃ H ₇	H	Н	6-I	2-CH ₃ -4-OCH ₂ CF ₂ CHF ₂	235-237
	448	i-C ₃ H ₇	Н	Н	6-I	3-F-4-0CHF 2	214-216
30	449	i-C3H7	Н	H	6-I	3,5-C12-4-OCHF2	211-213
	450	i-C3H7	Н	Н	6-I	3-0CH3-4-0CHF2	215-217
	451	i-C ₃ H ₇	н	Н	6-I	2,3-(CH ₃) ₂ -4-OCH ₃	253-254
35	452	i-C3H7	H	Н	6-I	2-CH ₃ -4-OCBrF ₂	192-194
	453	i-C ₃ H ₇	Н	H	6-I	2-CH ₃ -4-OCF ₂ CHF ₂	216-218
40	454	i-C₃H₁	Н	Н	6-I	2-CH ₃ -4-OCF ₂ CHF ₂ -5-Cl	230-232
10	455	i-C ₃ H ₇	H	Н	6-I	2-CH3-4-OCF2CHC1F	205-207
	456	i-C ₃ H ₇	Н	Н	6-I	2-CH ₃ -4-OCF ₂ CHFCF ₃	222-223
45	457	i-C ₃ H ₇	Н	H	6-I	2-CH ₃ -4-OCF ₂ CHFOCF ₃	258-260
	458	i-C₃H₁	Н	Н	6-I	2-CH ₃ -4-(3-CF ₃ -Ph0)	198-199
	459	i-C3H7	Н	H	6-I	2-CH ₃ -4-(F ₅ -PhO)	262-264
50	460	i-C₃H₁	Н	Н	6-I	2-CH ₃ -4-(5-CF ₃ -2-Pyi-0)	245-246

Table 1 (Cont'd)

5	No	R1	R ²	Rз	Xn	Ym	Physical Properties (melting point: °C
10	461	i-C3H7	Н	H	6-I	2-CH ₃ -4-(3-Cl-	231-232
						5-CF ₃ -2-Pyi-0)	
	462	i-C3H7	H	H	6-I	2-CH3-4-SC3H7-i	197-199
15	463	i-C3H7	Н	H	6-I	2-CH ₃ -4-(4-Cl-PhS)	211-213
	464	i-C ₃ H ₇	H	H	6-I	2-CH ₃ -4-0CF ₃	230-232
20	465	i-C ₃ H ₇	H	H	I-9	2-CF 3-4-0CHF 2	238-239
	466	i-C3H7	н	H	6-I	2-Br-4-0CF ₃	215-217
	467	i-C ₃ H ₇	н	H	6-I	2-C1-4-0CF ₃	186-188
25	468	i-C3H7	H	H	6-I	2-C1-4-CF ₂ CF ₂ CF ₃	199-200
	469	i-C₃H7	Н	H	6-I	2-CH3-4-OCHF2	226-228
	470	i-C₃H7	H	H	6-I	2-CH ₃ -4-0CHF ₂ -5-Cl	239-240
30	471	i-C3H7	Н	H	6-I	3-CF 3-4-0CHF 2	238-239
	472	i-C ₃ H ₇	H	H	3-F	4-(CF ₂) ₃ CF ₃	187-188
	473	i-C ₃ H ₇	H	Н	3-F	4-CF ₂ CF ₂ CF ₃	182-183
<i>35</i>	474	i-C ₃ H ₇	H	H	3-F	4-CF(CF ₃) ₂	206-208
	475	i-C ₃ H ₇	H	Н	3- F	4-0CF ₃	197-199
40	476	i-C ₃ H ₇	H	H	3-F	4-OCF2CHFOC3F7-n	142-144
40	477	i-CaH7	H	H	3-F	4-SCHF ₂	190-192
	478	i-CaH7	Н	Н	3- F	4-SCH ₂ CF ₃	157-158
45	479	i-C3H7	H	Н	3-F	4-SCF 2CHF 2	177-178
	480	i-CsH7	Н	Н	3-F	4-SCF ₂ CBrF ₂	197-199
	481	i-C ₃ H ₇	Н	Н	3-F	4-SCF(CF ₃) ₂	206-208
50	482	i-C ₃ H ₇	Н	Н	3-F	4-S(CF ₂) ₃ CF ₃	173-174

Table 1 (Cont'd)

			,				
5	.,	D .		,,	,,	,,,	Physical
	No	R:	R2	Rз	Xn	Ym	Properties
							(melting point: °C
10	483	i-C3H7	H	Н	3-F	4-SOCH ₂ CF ₃	115-119
	484	i-C ₃ H ₇	H	Н	3-F	4-SOCF 2CBrF 2	181-182
	485	i-C3H7	Н	Н	3-F	4-SOCF(CF ₃) ₂	195-197
15	486	i-C3H7	Н	Н	3-F	4-SO(CF ₂) ₃ CF ₃	175-176
	487	i-C₃H₁	Н	Н	3-F	4-S02CH2CF3	199-202
20	488	i-C₃H₁	Н	Н	3-F	2,3-Cl ₂	175-177
20	489	i-C₃H₁	Н	Н	3-F	2-CH ₃ -3-C1	193-194
	490	i-C₃H₁	н	Н	3-F	2-CH ₃ -4-Cl	192-194
25	491	i-C3H7	Н	Н	3-F	2-CH ₃ -5-Cl	191-193
	492	i-C ₃ H ₇	Н	H	3-F	2-CH ₃ -4-I	192-194
	493	i-C3H7	Н	Н	3-F	2-CH ₃ -5-F	175-177
30	494	i-C₃H₁	Н	Н	3-F	2-CH ₃ -3-F	187-189
	495	i-C ₃ H ₇	Н	Н	3-F	2-CH ₃ -4-CF ₂ CF ₃	213-214
	496	i-C ₃ H ₇	Н	H	3-F	2-CH ₃ -4-CF ₂ CF ₂ CF ₃	191-192
35 .	497	i-C3H7	Н	H	3-F	2-CH ₃ -4-CF(CF ₃) ₂	241-243
	498	i-C ₃ H ₇	H	Н	3-F	2-CH ₃ -4-(CF ₂) ₃ CF ₃	138-139
40	499	i-C ₃ H ₇	H	Н	3-F	2-CH ₃ -3-0CHF ₂	172-174
40	500	i-C ₃ H ₇	Н	Н	3-F	2-CH ₃ -4-OCHF ₂	160-162
	501	i-C ₃ H ₇	H	Н	3-F	2-CH ₃ -4-OCF ₂ CCl ₃	162-163
45	502	i-C ₃ H ₇	H	H	3-F	2-CH3-4-OCF2CC12F	207-208
	503	i-C ₃ H ₇	H	Н	3-F	2-CH3-4-OCF2CBrF2	196-197
	504	i-C ₃ H ₇	H	H	3-F	2-C1-4-CF ₃	169-170
50	505	i-C3H7	H	Н	3-F	2-C1-4-CF 2CF 2CF 3	169-170

Table 1 (Cont'd)

5	No	Rı	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	506	i-C ₃ H ₇	Н	H	3-F	3,5-C1 ₂ -4-0CHF ₂	201-202
	507	i-C ₃ H ₇	Н	н	3-F	2-C1-4-CF(CF ₃) ₂	223-225
	508	i-C ₃ H ₇	Н	H	3-F	2-Cl-4-0CF ₃	169-170
15	509	i-C3H7	H	H	3-F	2-Br-4-0CF ₃	164-165
	510	i-C ₃ H ₇	Н	H	3-F	2-CH ₃ -4-OCF ₃	183-184
20	511	i-C ₃ H ₇	Н	H	3-F	2-CH3-4-OCBrF2	177-178
	512	i-C ₃ H ₇	H	H	3-F	2-CH ₃ -4-OCF ₂ CHF ₂	172-173
	513	i-C ₃ H ₇	Н	H	3-F	2-CH3-4-OCF2CHC1F	168-169
25	514	i-C3H7	Н	H	3-F	2-CH ₃ -4-OCF ₂ CHFCF ₃	160-162
	515	i-C₃H₁	H	H	. 3-F	2-CH ₃ -4-OCF ₂ CHFOCF ₃	148-150
	516	i-C3H7	H	H	3-F	2-CH3-4-OCF2CBrFCF3	148-150
30	517	i-C3H7	H	Н	3-F	2-CH ₃ -4-OCHF ₂ -5-Cl	187-188
	518	i-C₃H₁	H	H	3-F	2-CH3-4-SC3H7-i	165-167
	519	i-C3H7	H	H	3-F	2-CH ₃ -4-(3-CF ₃ -PhO)	135-136
35	520	i-C3H7	H	H	3-F	2-CH ₃ -4-(F ₅ -PhO)	206-207
	521	i-C ₃ H ₇	H	H	3-F	2-CH ₃ -4-(2-Cl-4-	215-217
40				(-		CF ₃ -PhO)	
**	522	i-C₃H₁	H	H	3-F	2-CH ₃ -4-(4-Cl-PhS)	176-178
	523	i-C ₃ H ₇	H	H	3-F	2-CH ₃ -4-(5-CF ₃ -2-Pyi-0)	175-176
45	524	i-C ₃ H ₇	Н	H	3-F	2-CH ₃ -4-(3-Cl-	188-190
						5-CF ₃ -2-Pyi-0)	
	525	i-C3H7	H	H	3-F	4-(3-C1-5-CF ₃ -2-Pyi-S)	213-215
50	526	i-C ₃ H ₇	Н	Н	3-F	2-CH ₃ -4-OP=S(OCH ₃) ₂	175-177

Table 1 (Cont'd)

5	No	R1	R2	R3	Хn	Ym	Physical Properties (melting point: °C
10	527	i-C ₃ H ₇	Н	Н	3-F	2-CF ₃ -4-0CHF ₂	180-182
	528	i-C ₃ H ₇	Н	Н	3-F	-3-0CH ₂ 0-4-	197-199
	529	i-C₃H₁	Н	Н	4-F	2-CH ₃ -4-C1	217-218
15	530	i-C3H7	Н	Н	4-F	2-CH ₃ -5-Cl	202-203
	531	i-C3H7	Н	Н	4-F	2-CH ₃ -4-0CHF ₂	191-193
20	532	i-C ₃ H ₇	H	Н	5-F	2-CH3-4-Cl	197-198
	533	i-C ₃ H ₇	Н	H	5-F	2-CH ₃ -4-CF ₂ CF ₂ CF ₃	213-215
	534	i-C3H7	Н	H	5- F	2-CH ₃ -4-OCHF ₂	181-182
25	535	i-C ₃ H ₇	Н	H	6-F	4-CF ₂ CF ₂ CF ₃	201-202
	536	i-C ₃ H ₇	Н	H	6-F	4-(CF ₂) ₃ CF ₃	156-158
	537	i-C ₃ H ₇	Н	H	6-F	4-0CF ₃	212-214
30	538	i-C ₃ H ₇	Н	H	6-F	4-OCF ₂ CHFOC ₃ F ₇ -n	178-180
	539	i-C ₃ H ₇	H	Н	6-F	4-SCH ₂ CF ₃	176-178
	540	i-C ₃ H ₇	H	Н	6-F	4-SCF ₂ CHF ₂	230-232
35	541	i-C3H7	Н	H	6-F	4-SCF(CF ₃) ₂	218-220
	542	i-C3H7	H	H	6-F	4-S(CF ₂) ₃ CF ₃	178-181
40	543	i-C ₃ H ₇	Н	н	6-F	2,3-Cl ₂	158-160
	544	i-C3H7	Н	Н	6-F	2-CH ₃ -3-Cl	182-184
	545	i-C ₃ H ₇	Н	Н	6-F	2-CH ₃ -4-Cl	204-206
45	546	i-C ₃ H ₇	Н	Н	6-F	2-CH ₃ -5-Cl	196-199
	547	i-C ₃ H ₇	н	Н	6-F	2-CH ₃ -4-I	213-215
	548	i-C ₃ H ₇	н	H	6-F	2-CH ₃ -3-F	165-167
50	549	i-C ₃ H ₇	H	H	6-F	2-CH ₃ -5-F	181-183
Į			l				

Table 1 (Cont'd)

5	No	R1	R ²	R 3	Xn	Ym	Physical Properties (melting point: °C
10	550	i-C3H7	Н	Н	6-F	2-C1-4-CF ₃	190-191
	551	i-C₃H₁	Н	Н	6-F	2-CH ₃ -4-CF ₂ CF ₃	222-223
	552	i-C ₃ H ₇	H	Н	6-F	2-CH ₃ -4-OCF ₂ CCl ₃	184-185
15	553	i-C ₃ H ₇	н	H	6-F	2-CH ₃ -4-OCF ₂ CCl ₂ F	214-215
	554	i-C3H7	Н	H	6-F	2-CH3-4-OCF2CBrF2	208-210
20	555	i-C ₃ H ₇	Н	Н	6-F	2-CH ₃ -4-CF ₂ CF ₂ CF ₃	168-170
	556	i-C3H7	H	H	6-F	2-CH ₃ -4-CF(CF ₃) ₂	255-257
	557	i-C₃H7	H	H	6-F	2-CH ₃ -4-(CF ₂) ₃ CF ₃	157-159
25	558	i-C3H7	H	H	6-F	2-CH ₃ -3-0CHF ₂	177-179
	559	i-C ₃ H ₇	Н	H	6-F	2-CH ₃ -4-0CHF ₂	176-178
	560	i-C ₃ H ₇	Н	H	6-F	3,5-Cl ₂ -4-OCHF ₂	198-200
30	561	i-C ₃ H ₇	H	H	6-F	2-C1-4-CF(CF ₃) ₂	241-243
	562	i-C3H7	H	H	6-F	2-C1-4-0CF ₃	171-172
	563	i-C ₃ H ₇	H	H	6-F	2-Br-4-OCF ₃	181-182
35	564	i-C ₃ H ₇	H	Н	6-F	2-CH3-4-0CF3	193-195
	565	i-C ₃ H ₇	H	Н	6-F	2-CH3-4-OCBrF2	181-183
40	566	i-C3H7	Н	Н	6-F	2-CH ₃ -4-OCF ₂ CHF ₂	185-187
9	567	i-C3H7	Н	Н	6-F	2-CH3-4-OCF2CHC1F	175-176
	568	i-C ₃ H ₇	Н	Н	6-F	2-CH ₃ -4-OCF ₂ CHFCF ₃	176-178
45	569	i-C ₃ H ₇	Н	н	6-F	2-CH3-4-OCF2CBrFCF3	217-219
	570	i-C ₃ H ₇	Н	Н	6-F	2-CH ₃ -4-OCF ₂ CHFOCF ₃	183-185
	571	i-C ₃ H ₇	н	Н	6-F	2-CH ₃ -4-0CHF ₂ -5-Cl	209-211
50	572	i-C ₃ H ₇	Н	Н	6-F	2-CH ₃ -4-(3-CF ₃ -PhO)	184-185

Table 1 (Cont'd)

5	No	R:	R ²	R 3	Xn		Ym	Physical Properties (melting point: °C
10	573	i-C ₃ H ₇	Н	Н	6-F	2-CH₃	-4-(F ₅ -Ph0)	227-228
	574	i-C3H7	Н	Н	6-F	2-CH₃	-4-(2-C1-4-CF ₃ -PhO)	220-222
15	575	i-C3H7	H	Н	6-F	2-CH₃	-4-(4-Cl-PhS)	190-193
	576	i-C₃H7	Н	н	6-F	2-CH3	-4-(5-CF ₃ -2-Pyi-0)	206-207
20	577	i-C₃H7	Н	н	6-F	2-CH₃	-4-(3-Cl-	177-179
20							$5-CF_3-2-Pyi-0)$	
	578	i-C3H7	Н	Н	6-F	2-CH₃	-4-0P=S(OCH ₃) ₂	188-190
25	579	i-C3H7	Н	Н	6-F	2-CF 3	-4-0CHF ₂	223-225
	580	i-C3H7	Н	Н	6-F	-3-0C	H ₂ 0-4-	201-203
30								
30	581	i-C ₃ H ₇	Н	Н	3,6-F	2	2-CH 3-4-OCHF 2	203-204
	582	i-C₃H₁	Н	Н	3,6-F	2	2-CH ₃ -4-Cl	221-222
35	583	i-C3H7	Н	Н	3,4,5	,6-F ₄	2-CH ₃ -5-Cl	189-191
	584	i-C₃H₁	H	Н	3-NO2		2,3-Cl ₂	201-203
	585	i-C3H7	H	Н	3-NO ₂		н	236-238
40	586	i-C3H7	Н	Н	3-NO2		2-C1	190-192

Table 1 (Cont'd)

55

r				T	T		
5	No	R1	R2	Rз	Xn	Ym	Physical Properties
				•			(melting point: ℃
10	587	i-C3H7	Н	Н	3-NO ₂	3-C1	227-229
	588	i-C ₃ H ₇	Н	Н	3-NO ₂	4-C1	238-240
	589	i-C3H7	Н	н	3-NO ₂	2-Br	170-172
15	590	i-C ₃ H ₇	Н	Н	3-NO ₂	3-Br	196-198
	591	i-C ₃ H ₇	Н	H	3-NO ₂	4-Br	205-207
20	592	i-C ₃ H ₇	Н	H	3-NO ₂	2-F	199-201
	593	i-C ₃ H ₇	Н	Н	3-N0 ₂	3-F	228-230
	594	i-C₃H₁	Н	Н	3-NO ₂	4-F	250-252
25	595	i-C3H7	Н	Н	3-NO ₂	4-I	187-189
	596	i-C3H7	H	Н	3-NO ₂	4-NO ₂	201-203
	597	i-C3H7	H	Н	3-NO ₂	3-CN	220-222
30	598	i-C3H7	H	Н	3-NO ₂	4-CN	226-228
	599	i-C ₃ H ₇	Н	Н	3-NO ₂	2-CH ₃	227-228
	600	i-C ₃ H ₇	Н	H	3-NO2	3-CH₃	195-197
35	601	i-C3H7	Н	H	3-NO ₂	4-CH ₃	196-198
	602	i-C₃H₁	Н	H	3-NO ₂	2-C ₂ H ₅	189-191
40	603	i-C₃H7	H	H	3-NO ₂	2-C ₃ H ₇ -i	190-192
	604	i-C ₃ H ₇	Н	H	3-N0 ₂	4-C ₃ H ₇ -i	221-223
	605	i-C ₃ H ₇	H	н	3-N0 ₂	4-C ₄ H ₉ -n	193-195
45	606	i-C ₃ H ₇	H	Н	3-NO ₂	4-CF 3	192-194
	607	i-C ₃ H ₇	H	Н	3-NO ₂	3-CF ₃	220-222
	608	i-C ₃ H ₇	Н	н	3-NO ₂	2-CF ₃	215-217
50	609	i-C ₃ H ₇	H	H	3-N0 ₂	4-CF ₂ CF ₂ CF ₃	184-185
i							

Table 1 (Cont'd)

5	No	R1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	610	i-C ₃ H ₇	H	Н	3-NO ₂	4-CF(CF ₃) ₂	243-244
	611	i-C3H7	Н	Н	3-NO2	4-(CF ₂) ₃ CF ₃	220-221
	612	i-C ₃ H ₇	н	Н	3-NO2	2-0СН 3	172-174
15	613	i-C ₃ H ₇	H	H	3-NO2	3-0CH ₃	201-203
	614	i-C ₃ H ₇	H	H	3-NO2	4-0CH ₃	221-223
20 .	615	i-C ₃ H ₇	H	H	3-NO2	3-0-C3H7-i	198-200
	616	i-C ₃ H ₇	H	H	3-NO2	3-0CHF 2	188-190
	617	i-C ₃ H ₇	H	H	3-NO2	4-0CHF 2	222-224
25	618	i-C3H7	H	H	3-NO2	4-0CF ₃	234-236
	619	i-C ₃ H ₇	H	H	3-NO ₂	4-0CF 2CHF0C 3F 7-n	138-140
	620	i-C ₃ H ₇	H	H	3-NO ₂	4-C00CH ₃	192-194
30	621	i-C ₃ H ₇	H	Н	3-NO2	3-SCH₃	205-207
	622	i-C ₃ H ₇	Н	Н	3-NO2	2-SCH₃	201-203
	623	i-C ₃ H ₇	Н	Н	3-NO2	3-SCF ₃	203-205
35	624	i-C ₃ H ₇	Н	Н	3-NO ₂	4-SCH ₂ CF ₃	155-156
	625	i-C3H7	H	Н	3-NO ₂	4-SCHF ₂	183-185
40	626	i-C ₃ H ₇	H	Н	3-NO ₂	4-SCF 2 CHF 2	235-237
	627	i-C ₃ H ₇	Н	H	3-NO ₂	4-SCF 2 CF 3	190-192
	628	i-C ₃ H ₇	Н	H	3-NO ₂	4-SCF ₂ CBrF ₂	228-230
45	629	i-C ₃ H ₇	Н	H	3-NO ₂	4-SCF(CF ₃) ₂	242-243
	630	i-C ₃ H ₇	Н	н	3-NO ₂	4-S(CF ₂) ₃ CF ₃	229-230
	631	i-C ₃ H ₇	H	н	3-NO ₂	4-SO(CF ₂) ₃ CF ₃	190-193
50	632	i-C ₃ H ₇	H	H	3-NO ₂	4-0-Ph	228-230

Table 1 (Cont'd)

5	No	R1	R2	Rз	Хn	Ym	Physical Properties (melting point: °C
10	633	i-C ₃ H ₇	Н	H	3-NO2	2,4-Cl ₂	202-204
	634	i-C ₃ H ₇	H	H	3-NO2	2,5-Cl ₂	230-232
	635	i-C3H7	H	H	3-NO2	2,6-Cl ₂	210-212
15	636	i-C ₃ H ₇	H	H	3-NO2	3,4-Cl ₂	227-229
	637	i-C3H7	H	H	3-NO2	3,5-Cl ₂	194-196
20	638	i-C3H7	H	H	3-NO ₂	2,3-F ₂	184-186
	639	i-C3H7	H	H	3-NO ₂	2,4-F ₂	210-212
	640	i-C₃H₁	H	Н	3-NO2	2,5-F ₂	191-193
25	641	i-C ₃ H ₇	H	H	3-NO2	2,6-F ₂	173-175
	642	i-C ₃ H ₇	Н	Н	3-NO2	3,4-F ₂	241-243
	643	i-C3H7	Н	Н	3-NO ₂	3-C1-4-F	203-205
30	644	i-C3H7	Н	Н	3-NO2	2,3,4-Cl ₃	203-205
	645	i-C ₃ H ₇	Н	Н	3-NO ₂	2,3,4-F ₃	202-204
	646	i-C3H7	H	Н	3-NO ₂	2,3,4,5,6-F ₅	192-194
35	647	i-C ₃ H ₇	Н	Н	3-NO2	2,3-(CH ₃) ₂	200-202
	648	i-C3H7	H	Н	3-NO2	2,4-(CH ₃) ₂	201-203
40	649	i-C ₃ H ₇	Н	Н	3-NO ₂	2,5-(CH ₃) ₂	221-223
40	650	i-C3H7	H	Н	3-NO ₂	2,6-(CH ₃) ₂	234-236
	651	i-C ₃ H ₇	H	Н	3-NO ₂	3,4-(CH ₃) ₂	195-197
45	652	i-C ₃ H ₇	H	Н	3-NO ₂	2,4,6-(CH ₃) ₃	229-231
	653	i-C3H7	H	Н	3-NO ₂	2,6-(C ₂ H ₅) ₂	258-260
	654	i-C₃H₁	H	Н	3-NO ₂	3,5-(CF ₃) ₂	225-227
50	655	i-C ₃ H ₇	Н	Н	3-NO ₂	3-C1-4-CH ₃	208-210

Table 1 (Cont'd)

5	No	Rı	R2	Бз	Xn	Ym	Physical Properties (melting point: °C
10	656	i-C ₃ H ₇	Н	Н	3-NO ₂	2-C1-4-CH ₃	195-197
	657	i-C ₃ H ₇	Н	Н	3-NO2	2-F-4-C1-5-CH3	193-195
	658	i-C ₃ H ₇	Н	Н	3-NO2	3-C1-4-0CHF ₂	222-224
15	659	i-C ₃ H ₇	Н	Н	3-NO2	3,5-Cl ₂ -4-OCHF ₂	218-220
	660	i-C3H7	H	Н	3-NO ₂	2-C1-4-CF ₃	217-219
20	661	i-C₃H₁	Н	Н	3-NO ₂	2-C1-5-CF ₃	193-195
	662	i-C₃H₁	Н	Н	3-NO2	2,6-Cl ₂ -4-CF ₃	226-228
	663	i-C3H7	Н	Н	3-NO2	2-CH ₃ -3-Cl	198-200
25	664	i-C3H7	H	H	3-NO ₂	2-CH3-4-Cl	235-237
	665	i-C3H7	Н	Н	3-NO2	2-CH3-5-C1	218-219
	666	i-C3H7	Н	Н	3-NO ₂	2-CH ₃ -6-Cl	248-250
30	667	i-C ₃ H ₇	Н	H	3-NO ₂	2-C ₂ H ₅ -4-Cl	235-237
	668	i-C ₃ H ₇	Н	H	3-NO ₂	2-CH ₃ -4,5-Cl ₂	196-198
	669	i-C ₃ H ₇	H	Н	3-NO ₂	2,3-(CH ₃) ₂ -4-Cl	226-228
35	670	i-C ₃ H ₇	Н	Н	3-NO ₂	2,4-(CH ₃) ₂ -3-Cl	203-205
	671	i-C ₃ H ₇	н	н	3-NO ₂	2-CH3-4-Br	214-216
40	672	i-C ₃ H ₇	Н	Н	3-NO ₂	2-CH3-5-Br	191-193
10	673	i-C ₃ H ₇	Н	Н	3-NO ₂	2-CH ₃ -4-I	227-227
	674	i-C ₃ H ₇	H	н	3-NO ₂	2-CH3-3-F	199-201
45	675	i-C ₃ H ₇	H	H	3-NO ₂	2-CH3-4-F	226-228
i	676	i-C3H7	Н	H	3-NO ₂	2-CH ₃ -5-F	213-215
	677	i-C ₃ H ₇	н	H	3-NO ₂	2-C ₂ H ₅ -5-F	191-193
50	678	i-C ₃ H ₇	Н	H	3-NO ₂	3-CF ₃ -4-Cl	215-217

Table 1 (Cont'd)

5	No	R 1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	679	i-C3H7	H	Н	3-NO2	2-CF ₃ -4-Cl	208-210
	680	i-C ₃ H ₇	H	Н	3-NO2	3-CH ₃ -4-Br	199-201
	681	i-C3H7	H	H	3-NO2	2-CH ₃ -3-CF ₃	221-222
15	682	i-C3H7	H	H	3-NO2	2-CH3-4-CF3	236-237
	683	i-C₃H7	H	H	3-NO2	2-CH3-4-CF2CF3	218-219
20	684	i-C ₃ H ₇	Н	H	3-NO2	2-CH3-4-CF2CF2CF3	188-189
20	685	i-C ₃ H ₇	Н	Н	3-NO2	2-CH ₃ -4-CF(CF ₃) ₂	248-250
	686	i-C3H7	Н	Н	3-NO2	2-CH ₃ -4-(CF ₂) ₃ CF ₃	225-226
25	687	i-C₃H₁	Н	H	3-NO ₂	2-СН з - 3-ОСН з	198-200
	688	i-C3H7	Н	Н	3-NO ₂	2-СН з -4-ОСН з	208-210
	689	i-C ₃ H ₇	Н	Н	3-NO ₂	2,3-(CH ₃) ₂ -4-0CH ₃	253-255
30	690	i-C3H7	Н	H	3-NO2	2-CH3-4-0-C3H7-i	233-234
	691	i-C₃H₁	Н	Н	3-NO ₂	3-CF ₃ -5-0CH ₃	214-216
	692	i-C ₃ H ₇	H	H	3-NO ₂	2-CF 3-4-0CHF 2	201-203
35	693	i-CaH7	Н	Н	3-NO ₂	3-CF 3-4-0CHF 2	231-232
!	694	i-C3H7	Н	Н	3-NO ₂	2,4-(CH ₃) ₂ -3-0CH ₃	201-203
	695	i-C ₃ H ₇	Н	Н	3-NO ₂	2-CH ₃ -3-0CHF ₂	200-202
40	696	i-C3H7	Н	Н	3-NO ₂	2-CH ₃ -4-0CHF ₂	186-188
	697	i-C ₃ H ₇	Н	Н	3-NO2	2-CH ₃ -4-0CH ₂ CF ₃	241-243
	698	i-C3H7	Н	H	3-NO ₂	2-CH3-4-OCF2CBrF2	229-230
45	699	i-C3H7	Н	Н	3-NO ₂	2-CH3-4-OCH2CF2CHF2	199-200
	700	i-C3H7	Н	Н	3-NO ₂	2-CH3-4-OCF2CBrFCF3	224-226
_	701	i-C3H7	Н	Н	3-NO ₂	2-CH ₃ -4-OCH ₂ CHFOCF ₃	208-210
50							

Table 1 (Cont'd)

5		<u> </u>	T	1	T		Physical
	No	R 1	R2	R3	Xn	Ym	Properties
			İ				(melting
10			 	-			point: °C
10	702	i-C3H7	H	H	3-NO ₂	3-0CH ₃ -4-0CHF ₂	242-243
	703	i-C ₃ H ₇	H	H	3-NO ₂	2-C1-4-CF(CF ₃) ₂	198-200
15	704	i-C ₃ H ₇	H	H	3-NO ₂	2-C1-4-OCF ₃	188-190
	705	i-C ₃ H ₇	H	H	3-NO ₂	2-Br-4-0CF ₃	202-203
	706	i-C ₃ H ₇	H	Н	3-NO2	2-CH ₃ -4-NO ₂	201-203
20	707	i-C ₃ H ₇	H	H	3-NO2	2-C1-5-N0 ₂	193-195
	708	i-C₃H₁	H	H	3-NO2	2-CH3-5-NO2	197-199
	709	i-C ₃ H ₇	Н	H	3-NO2	2,3-(CH ₃) ₂ -4-NO ₂	207-209
25	710	i-C3H7	Н	H	3-NO2	2-CH ₃ -4-0CF ₃	184-186
	711	i-C ₃ H ₇	Н	H	3-NO2	2-CH ₃ -4-0CBrF ₂	217-218
	712	i-C3H7	Н	H	3-NO ₂	2-CH3-4-OCF2CHF2	205-207
30	713	i-C3H7	н	H	3-NO2	2-CH3-3-OCF2CHC1F	164-166
	714	i-C ₃ H ₇	Н	H	3-NO2	2-CH3-4-OCF2CHC1F	192-193
	715	i-C ₃ H ₇	Н	H	3-NO2	2-CH ₃ -4-OCF ₂ CCl ₂ F	212-213
35	716	i-C ₃ H ₇	H	Н	3-NO2	2-CH3-4-OCF2CHFCF3	198-199
	717	i-C ₃ H ₇	H	Н	3-NO2	2-CH ₃ -3-C1-4-0CHF ₂	236-238
40	718	i-C ₃ H ₇	H	Н	3-NO ₂	2-CH ₃ -4-OCF ₂ CHF ₂ -5-Cl	233-234
	719	i-C ₃ H ₇	H	H	3-NO ₂	2-CH ₃ -4-SCH ₃	214-216
	720	i-C ₃ H ₇	H	Н	3-NO ₂	2,3-(CH ₃) ₂ -4-SCH ₃	254-256
45	721	i-C₃H ₇	H	H	3-NO2	2-CH3-4-SC3H7-i	209-211
	722	i-C ₃ H ₇	Н	Н	3-NO ₂	2-CH ₃ -4-SCHF ₂	225-227
	723	i-C ₃ H ₇	Н	H	3-NO ₂	2-CH ₃ -4-N(CH ₃) ₂	215-217
50	724	i-C ₃ H ₇	H	н	3-NO ₂	2-CH ₃ -4-(3-CF ₃ -Ph0)	174-175

Table 1 (Cont'd)

5				γ	1		
·	No	R1	R2	R3	Xn	V-	Physical
	NO	n.	п-	l "	All	Ym	Properties (melting
							point: °C
10	725	i-C₃H₁	Н	Н	3-NO ₂	2-CH ₃ -4-(F ₅ -PhO)	242-244
	726	i-C ₃ H ₇	Н	H	3-NO ₂	2-CH ₃ -4-(2-C1-	191-192
						4-CF ₃ -Ph0)	
15	727	i-C3H7	Н	H	3-NO2	2-CH ₃ -4-(4-Cl-PhS)	165-167
	728	i-C3H7	H	H	3-NO2	2-CH ₃ -4-(5-CF ₃ -	216-218
20						2-Pyi-0)	
	729	i-C₃H7	Н	Н	3-N0 ₂	2-CH ₃ -4-(3-Cl-	236-238
						5-CF ₃ -2-Pyi-0)	
25	730	i-C3H7	Н	H	3-NO ₂	4-(3-C1-5-CF ₃ -	190-192
						2-Pyi-S)	
	731	i-C₃H7	H	Н	3-NO ₂	2-CH ₃ -4-P=0(OC ₂ H ₅) ₂	128-130
30	732	i-C ₃ H ₇	H	H	3-NO2	2-CH ₃ -4-P=S(OCH ₃) ₂	128-130
	733	i-C ₃ H ₇	H	H	3-NO2	-3-0CH₂0-4-	229-231
	734	i-C ₃ H ₇	Н	Н	3-N0 ₂	3-CH2CH2CH2-4	209-211
35	735	i-C ₃ H ₇	H	H	3-NO2	2-CH2CH2CH2-3	226-228
	736	i-C ₃ H ₇	H	Н	3-NO2	3-N=C(CF ₃)-NH-4	162-164
40	737	i-C ₃ H ₇	H	H	3-NO2	3-N=C(CF ₃)-N(CH ₃)-4	186-188
	738	i-C ₃ H ₇	H	H	5-NO ₂	2-CH3-5-Cl	226-228
	739	i-C ₃ H ₇	Н	Н	6-N0 ₂	2-CH ₃ -5-Cl	247-249
45	740	i-C ₃ H ₇	Н	н	6-NO ₂	2-C1-4-CF ₃	Crystals
	741	i-C ₃ H ₇	н	Н	6-NO2	2-C1-4-CF ₂ CF ₂ CF ₃	192-193
	742	i-C3H7	Н	H	6-NO2	2-CH ₃ -4-CF ₃	239-240
50	743	i-C3H7	н	H	6-NO ₂	2-CH ₃ -4-OCF ₂ CHFCF ₃	252-253

Table 1 (Cont'd)

5		т			Y	<u> </u>	
	No	R 1	R2	Rз	Xn	Ym	Physical Properties (melting
10	744	i-C ₃ H ₇	Н	Н	3-CN	2-CH ₃ -4-C1	point: °C
							162-164
45	745	i-C₃H₁	H	H	6-CN	2-CH ₃ -4-C1	Crystals
15	749	i-C₃H₁	H	H	3-CH 3	4-0CF ₃	180-182
	750	i-C₃H7	H	Н	3-CH 3	2-CH3-4-C1	169-171
20	751	i-C₃H₁	Н	Н	3-CH₃	2-CH ₃ -4-OCHF ₂	192-193
	752	i-C ₃ H ₇	H	Н	5-CH₃	2-CH₃-5-C1	193-195
	753	i-C₃H₁	H	Н	6-C2H5	2-CH ₃ -4-Cl	180-182
25	754	i-C₃H₁	Н	н	3-CF 3	H	202-204
	755	i-C₃H₁	H	Н	3-CF 3	2-CH ₃ -5-C1	196-198
30	756	i-C3H7	Н	H	3-CF 3	2-CH ₃ -3-C1	216-218
30	757	i-C ₃ H ₇	Н	H	3-CF 3	2,6-(C ₂ H ₅) ₂	238-239
	758	i-C ₃ H ₇	H	H	3-CF 3	2-CH ₃ -4-Cl	207-209
35	759	i-C ₃ H ₇	H	H	3-CF 3	2-CH3-4-OCHF2	212-213
	760	i-C ₃ H ₇	Н	Н	5-CF 3	2,6-(C2H5)2	240-241
	761	i-C3H7	H	H	5-CF ₃	2-CH ₃ -4-C1	203-205
40	762	i-C ₃ H ₇	H	Н	5-CF 3	3-CF ₃ -5-0CH ₃	209-210
	763	i-C3H7	Н	Н	5-CF 3	2-CH ₃ -4-0CHF ₂	196-197
45	764	i-C₃H₁	Н	H	6-CF 3	Н	152-154
	765	i-C ₃ H ₇	Н	Н	6-CF ₃	2-CH ₃ -3-C1	158-160
	766	i-C ₃ H ₇	H	H	6-CF ₃	2-CH ₃ -5-C1	273-275
50							

Table 1 (Cont'd)

5	No	R 1	R2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	767	i-C ₃ H ₇	Н	Н	3-0CH3	4-0CF ₃	178-180
	768	i-C ₃ H ₇	H	H	3-0CH₃	2-CH3-4-Br	214-215
	769	i-C ₃ H ₇	Н	H	6-0CH ₃	4-0CF ₃	189-190
15	770	i-C3H7	H	Н	6-0CH3	2-CH3-5-Cl	155-157
	771	i-C ₃ H ₇	H	н	6-0CH ₃	2-CH ₃ -4-Br	195-197
20	772	i-C ₃ H ₇	H	Н	3-0CHF 2	2-CH ₃ -4-Cl	212-213
-	773	i-C3H7	Н	Н	3-0CHF 2	2-CH ₃ -5-Cl	198-200
	774	i-C3H7	Н	Н	3-0CHF 2	2-CH 3-4-OCHF 2	174-175
25	775	i-C3H7	H	Н	4-0CHF 2	2-CH3-5-Cl	215-217
	776	i-C ₃ H ₇	Н	Н	5-0CHF 2	2-CH ₃ -5-Cl	173-175
	777	i-C ₃ H ₇	Н	Н	6-0CHF 2	2-CH3-4-C1	224-226
30	778	i-C3H7	Н	Н	6-0CHF 2	2-CH3-5-C1	191-193
	779	i-C3H7	Н	Н	6-0CHF 2	2-CH 3-4-0CHF 2	199-200
	780	i-C3H7	Н	Н	3-SCH ₃	2-CH ₃ -3-Cl	191-193
35	781	i-C3H7	Н	Н	3-SCH₃	2-CH ₃ -4-Cl	188-190
	782	i-C3H7	Н	н	3-SCH₃	2-CH3-4-Br	185-187
40	783	i-C ₃ H ₇	Н	Н	3-SCH₃	2-CH3-4-0CHF2	159-161
40	784	i-C3H7	Н	Н	6-SCH₃	2-CH3-4-Br	201-202
	785	i-C ₃ H ₇	Н	Н	6-SCH₃	2-CH ₃ -3-Cl	207-209
45	786	i-C₃H₁	Н	Н	6-SCH ₃	2-CH3-4-Cl	204-206
	787	i-C ₃ H ₇	Н	Н	6-SCH ₃	2-CH3-4-0CHF2	212-214
	788	i-C ₃ H ₇	Н	H	3-SC ₃ H ₇ -i	2-CH ₃ -4-Cl	183-184
50	789	i-C ₃ H ₇	Н	Н	6-SC3H7-i	2-CH ₃ -4-Cl	228-229

Table 1 (Cont'd)

5		,	,				
10	No	R1	R ²	Кз	Xn	Ym	Physical Properties (melting point: °C
	790	i-C ₃ H ₇	Н	Н	3-SOCH 3	2-CH ₃ -4-Br	125-130
	791	i-C ₃ H ₇	Н	Н	3-SOCH3	2-CH ₃ -4-0CHF ₂	215-217
15	792	i-C ₃ H ₇	Н	Н	6-SOCH ₃	2-CH ₃ -4-Br	203-208
	793	i-C ₃ H ₇	Н	H	3-SOC ₃ H ₇ -i	2-CH ₃ -4-Cl	157-160
	794	i-C3H7	Н	H	6-SOC ₃ H ₇ -i	2-CH ₃ -4-Cl	170-173
20	795	i-C ₃ H ₇	Н	Н	3-S0 ₂ CH ₃	2-CH ₃ -4-OCHF ₂	211-213
	796	i-C3H7	Н	Н	3-S0 ₂ C ₃ H ₇ -i	2-CH ₃ -4-Cl	240-242
25	797	i-C₃H₁	Н	Н	3-SCH ₂ CF ₃	2-CH ₃ -4-0CHF ₂	184-186
	798	i-C3H7	Н	Н	6-SCH ₂ CF ₃	2-CH ₃ -4-0CHF ₂	239-241
	799	i-C₃H7	H	Н	3-SOCH2CF3	2-CH ₃ -4-0CHF ₂	198-200
30	800	i-C ₃ H ₇	Н	Н	6-SOCH ₂ CF ₃	2-CH ₃ -4-0CHF ₂	238-240
	801	i-C₃H7	H	Н	6-С≡СН	2-CH3-4-Cl	253-255
	802	i-C3H7	H	Н	6-COOCH3	2-CH3-4-Cl	149-151
35	803	i-C ₃ H ₇	H	H	3-CONHC ₃ H ₇ -i	2-CH3-4-C1	187-189
	804	i-C3H7	H	H	6-CONHC ₃ H ₇ -i	2-CH3-4-Cl	191-193
40	807	i-C ₃ H ₇	H	H	3-Ph	2-CH3-4-Cl	228-229
	808	i-C ₃ H ₇	H	Н	6-Ph	4-0CF 3	213-214
	809	i-C ₃ H ₇	H	H	6-Ph	2-CH3-4-Cl	254-256
45	810	i-C ₃ H ₇	Н	H	3-0-Ph	2-CH3-4-OCHF2	175-177
	811	i-C ₃ H ₇	Н	н	6-0-Ph	2-CH3-4-0CHF2	194-196
	812	i-C ₃ H ₇	н	Н	3-(4-C1-Ph0)	2-CH3-4-Br	204-206
50							

Table 1 (Cont'd)

5	No	R 1	R2	Rз		Xn	Ym	Physical Properties (melting point: °C
10	813	i-C ₃ H ₇	Н	н	3-S-Ph		2-CH ₃ -4-C1	204-206
	814	i-C3H7	H	Н	3-S-Ph		2-CH3-4-Br	193-194
	815	i-C ₃ H ₇	Н	Н	6-S-Ph		2-CH ₃ -4-Cl	211-213
15	816	i-C3H7	H	H	6-S-Ph		2-CH3-4-Br	193-194
	817	i-C ₃ H ₇	Н	H	3-S0-PI	ı	2-CH ₃ -4-Cl	201-203
20	818	i-C ₃ H ₇	H	H	3-SO ₂ -I	Ph	2-CH3-4-C1	189-191
	819	i-C ₃ H ₇	Н	н	3-CH=CI	I-CH=CH-4	2-CH3-4-0CHF2	158-160
	820	i-C3H7	Н	H	5-CH=CH	I-CH=CH-6	2-CH ₃ -4-0CHF ₂	154-155
25	821	i-C ₃ H ₇	Н	Н	3-СН=СН	I-CH=CH-4	2-CH3-5-C1	156-158
	822	i-C3H7	H	Н	4-CH=CH	І-СН=СН-5	2-CH3-5-C1	229-231
	823	i-C ₃ H ₇	Н	Н	5-CH=CH	I-CH=CH-6	2-CH3-5-C1	232-234
30				_ [_		
	824	i-C ₃ H ₇	СН₃		_ н	Н	4-CF 3	178-180
	825	i-C ₃ H ₇	СНз		Н	3-NO ₂	2-CH ₃ -4-OCHF ₂	148-149
35	826	i-C ₃ H ₇	СНз		Н	Н	2-CH3-4-C1	82-83
	827	i-C ₃ H ₇	Н		CH ₃	Н	2-CH3-4-C1	165-166
40	828	i-C ₃ H ₇	CH ₂ (ОСН₃	Н	н	2-CH₃-4-Cl	0il
	829	n-C4H9	H		Н	Н	4-CF 3	171-173
	830	n-C ₄ H ₉	Н		Н	3-NO2	2-CH ₃ -5-C1	172-174
45	831	i-C ₄ H ₉	Н		Н	3-NO ₂	2-CH3-5-C1	186-188
	832	i-C₄H ₉	H		Н	3-NO ₂	2-CH3-4-0CHF2	192-193
	833	i-C ₄ H ₉	H		H	H	4-CF ₃	149-151
50	834	i-C₄H ₉	СНз		Н	6-NO ₂	2-CH ₃ -4-0CHF ₂	135-137

Table 1 (Cont'd)

5	No	R1	R	2	Ra	Xn	Ym	Physical Properties (melting point: °C	
10	835	s-C ₄ H ₉	Н		H	Н	4-CF ₃	194-195	-
	836	s-C4H9	Н		H	3-C1	2-CH ₃ -4-0CHF ₂	203-205	
	837	s-C4H9	H		Н	6-C1	2-CH ₃ -4-OCHF ₂	213-215	
15	838	s-C4H9	Н		Н	3-NO ₂	2-CH ₃ -5-C1	205-207	
	839	s-C4H9	Н		Н	3-NO ₂	2-CH ₃ -4-OCHF ₂	228-229	
20	840	t-C ₄ H ₉	Н		Н	Н	Н	237-239	
· ·	841	t-C₄H9	Н		Н	Н	2-CH ₃ -5-Cl	200-202	
	842	t-C ₄ H ₉	Н		Н	3-NO ₂	2-CH3-5-C1	256-258	
25	843	t-C₄H ₀	Н		Н	3-NO ₂	2-CH3-4-0CHF2	172-173	
	844	CH ₂ C(CH ₃) ₃	H		Н	3-NO ₂	2-CH ₃ -4-0CHF ₂	226-227	
	845	CH(C ₂ H ₅) ₂		Н	Н	3-NO2	2-CH ₃ -4-0CHF ₂	245-246	
30	846	CH(CH3)CH(C	H ₃) ₂	Н	Н	3-NO ₂	2-CH3-4-0CHF2	245-247	
	847	n-C ₈ H ₁₇		Н	Н	3-NO2	2-CH3-5-C1	164-166	
	848	c-C3H5	Н		Н	Н	4-CF 3	195-197	ĺ
35	849	c-C3H5	Н		H	3-C1	2-CH3-4-0CHF2	156-158	ĺ
	850	c-C3H5	Н		H	6-C1	2-CH3-4-0CHF2	179-181	
40	851	c-C ₃ H ₅	Н		Н	3-NO ₂	2-CH ₃ -5-Cl	194-196	
	852	c-C ₃ H ₅	H		H	3-NO2	2-CH3-4-0CHF2	191-192	
	853	C-C4H7	H		Н	H	2-CH3-5-C1	205-207	
45	854	c-C4H7	H		Н	3-NO ₂	2-CH₃-5-C1	206-208	
	855	c-C4H7	Н		Н	3-NO ₂	2-СН₃-5-F	199-201	
	856	c-C ₅ H ₉	H		н	3-NO2	2-CH3-4-0CHF2	219-220	
50	857	c-C ₅ H ₉	H		H	H	4-CF 3	208-210	

Table 1 (Cont'd)

5	No	R:	R2	Вз	Xn	Ym	Physical Properties (melting point: °C
10	858	c-C ₅ H ₉	Н	Н	3-NO ₂	2-CH ₃ -5-Cl	200-202
	859	C-C6H11	Н	Н	3-NO ₂	2-CH ₃ -5-Cl	225-227
15	860	CH2-C3H5-C	Н	Н	3-NO ₂	2-CH ₃ -5-F	190-192
	861	CH2CH2C1	Н	Н	3-NO2	2-CH ₃ -5-F	179-181
	862	CH2CH2F	H	Н	3-NO ₂	2-CH ₃ -5-F	179-181
20	863	CH2CH2F	Н	Н	3-NO2	2-CH ₃ -4-0CHF ₂	190-191
	864	CH ₂ CF ₃	H	Н	H	2-CH ₃ -5-Cl	187-189
	865	CH 2 CH=CH 2	H	Н	Н	4-CF 3	161-163
25	866	CH 2 CH=CH 2	Н	Н	3-NO2	2-CH3-5-C1	175-177
	867	CH 2 CH=CH 2	Н	Н	3-NO2	2-CH ₃ -4-OCHF ₂	194-195
30	868	CH₂C≡CH	Н	Н	H	4-CF 3	185-188
	869	CH₂C≡CH	Н	Н	3-NO ₂	2-CH ₃ -5-Cl	191-193
	870	CH₂C≡CH	H	Н	3-NO ₂	2-CH ₃ -4-0CHF ₂	190-191
35	871	CH2CH2OCH3	Н	H	3-NO2	2-CH ₃ -5-Cl	165-167
	872	CH2CH2OCH3	Н	Н	3-NO ₂	2-CH3-4-0CHF2	165-167
	873	CH(CH ₃)CH ₂ OCH ₃	H	H	H	4-CF 3	252-253
40	874	CH(CH ₃)CH ₂ OCH ₃	H	Н	3-NO2	2-CH ₃ -4-0CHF ₂	153-155
	875	CH ₂ CH(OC ₂ H ₅) ₂	H	Н	3-NO2	2-CH ₃ -4-0CHF ₂	149-151
45	876	CH2-Ph	H	Н	H	4-CF 3	148-150
	877	CH2-Ph	Н	Н	3-NO ₂	2-CH3-5-C1	196-198
	878	CH(CH ₃)-Ph	н	Н	3-NO ₂	2-CH ₃ -5-Cl	168-170
50	879	CH(CH ₃)-Ph	Н	Н	3-NO ₂	2-CH ₃ -4-0CHF ₂	187-189

Table 1 (Cont'd)

5		<u> </u>	Τ	Γ	1	 	Physical
	No	R1	R2	Rз	Xn	Ym	Properties
					ļ		(melting
10		OT 07 0 10 1	-				point: °C
	880	CH ₂ CH ₂ 0-(2,4-	H	H	3-NO ₂	2-CH ₃ -5-Cl	126-128
		(CH ₃) ₂ -Ph)					
15	881	-CH2CH2CH2	CH 2 -	H	Н	4-CF 3	170-171
,,	882	-CH2CH2CH2	CH 2 -	H	6-NO2	2-CH ₃ -5-Cl	157-159
	883	-CH2CH2CH2	CH 2 -	H	6-NO2	2-CH ₃ -4-OCHF ₂	163-165
20	884	-CH2CH2OCH2	CH2-	H	H	4-CF ₃	167-168
	885	-CH2CH2OCH2	CH2-	H	6-N0 ₂	2-CH₃-5-Cl	192-194
	886	-CH2CH2OCH2	CH 2 -	H	6-NO ₂	2-CH3-4-OCHF2	186-188
25	887	-CH2CH(CH3)		H	6-NO2	3-CF ₃ -5-OCH ₃	164-165
		OCH(CH₃)	CH 2 −				
	888	CH2-3-Pyi	H	H	3-NO2	2-CH3-4-Br	180-182
30	889	i-C ₃ H ₇	H	H	Н	4-CF 2CF 3	155-157
	890	i-C ₃ H ₇	H	H	3-NO2	4-CF 2CF 3	223-225
35	891	i-C ₃ H ₇	H	H	3-F	4-CF 2CF 3	199-201
33	892	i-C ₃ H ₇	H	H	6-F	4-CF 2CF 3	213-215
	893	i-C ₃ H ₇	Н	H	3-C1	4-CF 2CF 3	214-216
40	894	i-C3H7	Н	H	6-C1	4-CF ₂ CF ₃	225-227
	895	i-C3H7	Н	H	3-I	4-CF ₂ CF ₃	208-210
	896	i-C3H7	н	H	6-I	4-CF 2CF 3	224-226
45	897	i-CaH7	н	н	H	2-CH ₃ -4-0S0 ₂ -	135-137
						(4-CH ₃ -Ph)	
	898	i-C ₃ H ₇	H	H	3-NO ₂	2-CH ₃ -4-0S0 ₂ -	208-210
50						(4-CH3-Ph)	*

Table 1 (Cont'd)

5	No	R 1	R ²	R3	Xn	Ym	Physical Properties (melting
40		<u> </u>	ļ	-	ļ. <u>.</u>		point: °C
10	899	i-C ₃ H ₇	H	H	3-C1	2-CH ₃ -4-0S0 ₂ -	187-189
						(4-CH ₃ -Ph)	
15	900	i-C ₃ H ₇	H	H	6-C1	2-CH ₃ -4-0S0 ₂ -	218-220
70						(4-CH₃-Ph)	
	901	i-C ₃ H ₇	Н	Н	3-F	2-F-4-0-(4-CF ₃ -2-	137-139
20						Cl-Ph)	
	902	i-C ₃ H ₇	Н	H	6-F	2-F-4-0-(4-CF ₃ -2-	155-157
						C1-Ph)	
25	903	i-C ₃ H ₇	н	H	3-C1	2-F-4-0-(4-CF ₃ -2-	119-121
						Cl-Ph)	
	904	i-C3H7	Н	Н	6-C1	2-F-4-0-(4-CF ₃ -2-	154-156
30						C1-Ph)	
	905	i-C ₃ H ₇	Н	Н	3-F	2-CH ₃ -4-SCF ₂ CF ₃	140-142
	906	i-C₃H₁	Н	Н	6-F	2-CH3-4-SCF2CF3	162-164
35	907	i-C ₃ H ₇	Н	Н	3-C1	2-CH3-4-SCF2CF3	172-173
	908	i-C ₃ H ₇	Н	Н	6-C1	2-CH ₃ -4-SCF ₂ CF ₃	193-195
	909	i-C ₃ H ₇	Н	Н	3-I	2-CH3-4-SCF2CF3	207-209
40	910	i-C ₃ H ₇	Н	Н	6-I	2-CH3-4-SCF2CF3	196-198
	911	i-C ₃ H ₇	Н	Н	3-C1	4-CH=C(Cl)CF₃	196.3-208.2
45	912	i-C ₃ H ₇	Н	Н	6-C1	4-CH=C(Cl)CF ₃	202.8-209.4
10	913	i-C ₃ H ₇	H	Н	3-C1	4-CH=CBr ₂	209.8-214.8
	914	i-C₃H₁	Н	Н	6-C1	4-CH=CBr ₂	207.7-213.9
50	915	i-C3H7	H	Н	3-C1	4-CH=CCl ₂	120.1
						L	

Table 1 (Cont'd)

5	No	R 1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	916	i-C3H7	Н	Н	6-C1	4-CH=CCl ₂	199.7
	917	i-C ₃ H ₇	Н	Н	3-I	4-CH=C(C1)CF ₃	196.6
45	918	i-C ₃ H ₇	Н	Н	6-I	4-CH=C(C1)CF ₃	203.3
15	919	i-C ₃ H ₇	Н	Н	3-1	2-C ₂ H ₅ -4-I	195.5
	920	i-C ₃ H ₇	Н	Н	6-I	2-C ₂ H ₅ -4-I	242.3
20	921	C ₂ H ₅	Н	Н	Н	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	171-173
	922	i-C3H7	Н	H	Н	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	185-186
	923	t-C₄H ₉	Н	H	Н	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	166-167
25	924	i-C ₃ H ₇	Н	H	3-C1	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	260-261
	925	i-C₃H₁	H	H	3-I	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	269-271
	926	t-C₄H ₉	H	H	3-C1	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	221-222
30	927	t-C₄H ₉	Н	H	Н	2-CH ₃ -4-Cl	216-218
	928	t-C4H9	Н	Н	Н	4-CF 3	220-221
	929	t-C₄H ₉	Н	H	Н	4-0CF 3	178-179
35	930	t-C4H9	H	H	Н	2-CH ₃ -4-0CF ₃	184-185
	931	t-C ₄ H ₉	Н	H	Н	2-CH ₃ -4-CF ₂ CF ₃	223-224
40	932	t-C ₄ H ₉	н	H	3-C1	2-CH3-4-CF2CF3	219-220
	933	t-C ₄ H ₉	Н	СНз	H	4-0CF ₃	155-158
	934	t-C ₄ H ₉	Н	H	3-C1	4-CF ₃	228-229
45	935	t-C4H9	H	Н	6-C1	4-CF 3	253-255
	936	t-C ₄ H ₉	Н	Н	3-C1	4-0CF ₃	268-270
	937	t-C₄H ₉	H	Н	3-C1	2-CH3-4-C1	242-244
50	938	t-C4H9	н	H	6-C1	2-CH3-4-C1	262-264
Į							

Table 1 (Cont'd)

5	No	R 1	R2	R3	Xn	Ym	Physical Properties (melting
		}	<u> </u>	ļ			point: °C
10	939	t-C₄H,	Н	H	3-1	4-CF ₃	268-269
	940	t-C₄H₀	H	Н	3-1	4-0CF ₃	263-265
15	941	t-C ₄ H ₉	Н	H	3-I	2-CH3-4-C1	218-220
15	942	t-C₄H ₉	Н	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	205-207
	943	t-C₄H ₉	Н	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	216-217
20	944	t-C₄H ₉	Н	Н	3-C1	2-CH ₃ -4-0CF ₃	260-262
	945	n-C ₄ H ₉	Н	Н	3-1	2-CH ₃ -4-CF ₂ CF ₃	173.1-178.5
	946	n-C4H9	Н	Н	6-1	2-CH ₃ -4-CF ₂ CF ₃	181.8-187.7
25	947	n-C5H11	Н	Н	3-1	2-CH ₃ -4-CF ₂ CF ₃	140.2-151.4
	948	n-C ₅ H ₁ 1	Н	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	168.7-171.3
	949	n-C ₆ H ₁₃	Н	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	135.5-143.9
30	950	n-C6H13	H	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	167.1-169.9
	951	i-C3H7	H	H	3-I	2-C ₂ H ₅ -4-I	254.8-273.8
	. 952	i-C ₃ H ₇	Н	H	3-I	2-n-C3H7-4-I	179.7
35	953	i-C ₃ H ₇	Н	H	3-CH ₃	2-CH ₃ -4-CF ₂ CF ₃	184-186
	954	i-C ₃ H ₇	Н	H	6-CH ₃	2-CH ₃ -4-CF ₂ CF ₃	177-179
40	955	t-C ₄ H ₉	H	H	3-CH ₃	2-CH ₃ -4-CF ₂ CF ₃	198-200
40	956	t-C ₄ H ₉	H	H	6-CH₃	2-CH ₃ -4-CF ₂ CF ₃	236-237
•	957	t-C ₄ H ₉	H	H	3-I	2-CH ₃ -4-0CF ₃	208-210
45	958	t-C ₄ H ₉	н	H	6-I.	2-CH ₃ -4-0CF ₃	253-255
	959	n-C ₃ H ₇	н	H	1-8	2-CH ₃ -3-Cl	190-192
	960	n-C ₃ H ₇	н	H	6-I	2-CH ₃ -3-Cl	159-161
50	961	n-C ₃ H ₇	H	H	6-I	2-C ₂ H ₅ -3-Cl-6-C ₂ H ₅	225-228

Table 1 (Cont'd)

5	No	R1	R2	Rз	Хn	Ym	Physical Properties (melting point: °C
10	962	i-C3H7	H	Н	3-NO2	4-0C0CF ₃	185-187
	963	i-C ₃ H ₇	H	H	3-C1	4-0C0CF ₃	Paste
15	964	i-C ₃ H ₇	Н	Н	3-I	4-0C0CF ₃	Paste
15	965	i-C ₃ H ₇	Н	H	3-I	2-i-C ₃ H ₇ -4-I	132.5
	966	i-C ₃ H ₇	H	H	3-I	2-n-C4H8-4-I	194.2-198.3
20	967	i-C ₃ H ₇	Н	H	3-I	2-CH ₃ -4-Br-6-CH ₃	119.1
	968	i-C₃H₁	Н	H	3-C1	4-C0 ₂ CH(CF ₃) ₂	168-170
	969	i-C ₃ H ₇	H	H	3-I	4-C0 ₂ CH(CF ₃) ₂	193-195
25	970	i-C₃H₁	Н	H	3-NO ₂	4-C0 ₂ CH(CF ₃) ₂	215-217
	971	i-C ₃ H ₇	Н	H	3-C1	2-CH ₃ -4-C≡C-	123-125
						(2,4-Cl ₂ -Ph)	
30	972	i-C3H7	Н	H	3-I	2-CH ₃ -4-C≡C-	138-140.
						(2,4-Cl ₂ -Ph)	
	973	i-C ₃ H ₇	H	H	3-C1	3-0CF 2CF 2-4	125-128
35	974	i-C₃H ₇	H	H	3-I	3-0CF 2CF 2-4	123-126
	975	i-C ₃ H ₇	H	Н	H	3-0CF 2CF 20-4	152-154
40	976	i-C ₃ H ₇	н	H	3-NO ₂	3-0CF 2CF 20-4	247-248
	977	i-C ₃ H ₇	H	Н	3-C1	3-0CF 2CF 20-4	224-226
	978	i-C3H7	H	H	H	4-C(CF ₃) ₂ OH	87-89
45	979	i-C ₃ H ₇	H	H	3-NO ₂	4-C(CF ₃) ₂ OH	205-207
	980	i-C3H7	Н	Н	3-C1	4-C(CF ₃) ₂ OH	187-189
	981	CH2CH2OCH3	Н	н	3-I	2-CH ₃ -4-CF ₂ CF ₃	145.3-151.7
50	982	CH2CH2OCH3	н	н	6-I	2-CH ₃ -4-CF ₂ CF ₃	166.7-169.4

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	983	CH2CH2OC2H5	H	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	146.5-150.3
	984	CH2CH2OC2H5	H	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	157.3-160.4
15	985	(CH ₂) ₃ OCH ₃	H	H	3-I	2-CH3-4-CF2CF3	151.9-155.8
15	986	(CH ₂) ₃ OCH ₃	H	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	156.5-158.8
	987	CH 2 CH=CH 2	Н	H	3-I	2-CH3-4-CF2CF3	157.5
20	988	CH 2 CH=CH 2	H	H	6-I	2-CH3-4-CF2CF3	164.6-171.3
	989	CH₂C≡CH	H	H	3-I	2-CH3-4-CF2CF3	153.6-158.4
	990	CH ₂ C≡CH	Н	H	6-I	2-CH3-4-CF2CF3	171.5-178.1
25	991	c-C ₅ H ₉	H	H	3-I	2-CH3-4-CF2CF3	212.9
	992	c-C ₅ H ₉	H	H	6-I	2-CH3-4-CF2CF3	205.2
	993	C-C6H11	Н	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	219.7-224.3
30	994	C-C6H11	H	H	6-I	2-CH3-4-CF2CF3	239.0-244.4
	995	i-C ₃ H ₇	Н	H	Н	4-SCF 3	182-184
	996	i-C ₃ H ₇	H	H	3-NO ₂	4-SCF ₃	228-229
35	997	i-C ₃ H ₇	H	H	3-C1	4-SCF 3	229-231
	998	i-C ₃ H ₇	H	H	3-I	4-SCF 3	226-227
40	999	i-C ₃ H ₇	H	H	H	4-SOCF 3	175-178
+0	1000	i-C3H7	H	Н	3-NO ₂	4-SOCF 3	202-205
	1001	i-C3H7	Н	Н	3-C1	4-SOCF 3	242-244
45	1002	i-C3H7	Н	Н	3-I	4-SOCF 3	229-231
	1003	i-C3H7	Н	Н	3-I	3-0CF 2CF 20-4	163-165
	1004	i-C3H7	н	Н	1-6	4-C(CF ₃) ₂ 0H	227-229
50	1005	i-C ₄ H ₉	н	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	200.4-206.8

Table 1 (Cont'd)

5	No	R1	R2	Ra	Xn	Ym	Physical Properties (melting point: °C
10	1006	i-C ₄ H ₉	Н	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	179.2-181.8
•	1007	s-C4H9	Н	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	226.0-230.9
15	1008	s-C4H9	Н	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	216.1-218.0
	1009	s-C ₅ H ₁₁	Н	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	215.3-218.2
	1010	s-C ₅ H ₁₁	Н	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	191.4-210.5
20	1011	CH(C ₂ H ₅) ₂	H	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	234.8-236.9
	1012	CH(C ₂ H ₅) ₂	H	H	1-6	2-CH ₃ -4-CF ₂ CF ₃	253.7-255.7
	1013	CH(C ₂ H ₅)CH ₂ O	H	H	3-1	2-CH ₃ -4-CF ₂ CF ₃	177
25		-CH 3					
	1014	CH(C ₂ H ₅)CH ₂ O	H	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	198.3-201.0
30		-СН 3					
	1015	i-C5H11	Н	H	3-I	2-CH3-4-CF2CF3	190.0-192.5
	1016	i-C ₅ H ₁₁	H	H	6-I	2-CH3-4-CF2CF3	187.8
35	1017	i-C3H7	H	Н	3-I	2-C ₂ H ₅ -4-CF ₂ CF ₃	232.5-235.8
	1018	t-C4H9	Н	Н	H	2-CH ₃ -4-OCHF ₂	138-140
	1019	t-C ₄ H ₉	Н	Н	3-C1	2-CH ₃ -4-OCHF ₂	206-208
40	1020	t-C ₄ H ₉	Н	Н	3-I	2-CH ₃ -4-OCHF ₂	204-206
	1021	t-C ₄ H ₉	H	Н	H	2-C1-4-OCF ₃	162-164
45	1022	t-C ₄ H ₉	Н	H	3-C1	2-C1-4-OCF ₃	189-191
	1023	t-C ₄ H ₉	Н	H	3-1	2-C1-4-0CF ₃	188-190
	1024	C-C ₃ H ₅	H	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	156.0-165.0
50	1025	C-C ₃ H ₅	H	H	1-6	2-CH ₃ -4-CF ₂ CF ₃	173.2-176.4
ļ						· · · · · · · · · · · · · · · · · · ·	

Table 1 (Cont'd)

5		···					
	No	R1	R2	Rз	Xn	Ym	Physical Properties
							(melting
10							point: °C
70	1026	CH ₂ CH(CH ₃)	H	H	3-1	2-CH ₃ -4-CF ₂ CF ₃	148.6
		-C ₂ H ₅					
15	1027	CH ₂ CH(CH ₃)	H	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	157.8
15		-C2H5				·	
	1028	CH2-c-C6H11	H	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	186.8-188.7
20	1029	CH ₂ (4-t-C ₄ H ₉	H	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	226.0-231.2
		-c-C ₆ H ₁₁)					
	1030	CH2(4-t-C4H9	H	H	I-9	2-CH ₃ -4-CF ₂ CF ₃	215.4
25		-c-C ₆ H ₁₁)					
	1031	CH(CH₃)CH2O	H	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	187.2-189.9
		-СН ₃					
30	1032	CH(CH ₃)CH ₂ O	Н	Н	1-6	2-CH3-4-CF2CF3	169.7-176.1
		−СН з					
	1033	CH(CH ₃)CH	Н	Н	3-I	2-CH3-4-CF2CF3	208.3-212.7
35		-(CH ₃) ₂					
	1034	CH(CH ₃)CH	H	Н	6-I	2-CH3-4-CF2CF3	219.3-223.0
		-(CH ₃) ₂					
40	1035	C ₂ H ₅	C ₂ H ₅	Н	3-I	2-CH3-4-CF2CF3	131.3
	1036	C2H5	C ₂ H ₅	Н	6-I	2-CH3-4-CF2CF3	137
45	1037	t-C4He	H	Н	Н	2-CH ₃ -4-CF(CF ₃) ₂	172-175
	1038	t-C₄H ₉	H	Н	3-C1	2-CH ₃ -4-CF(CF ₃) ₂	241-243
	1039	t-C₄H ₉	H	Н	3-I	2-CH ₃ -4-CF(CF ₃) ₂	238-240
50	1040	CH 2 CF 3	H	Н	3-I	2-CH3-4-CF2CF3	166.1-175.5

Table 1 (Cont'd)

5	No	R1	R²	R3	Xn	Ym	Physical Properties (melting point: °C
10 .	1041	CH2CF3	Н	Н	6-I	2-CH ₃ -4-CF ₂ CF ₃	184.7-202.5
	1042	i-C ₃ H ₇	СН з	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	201.4
4.5	1043	i-C₄H ₉	CH 3	Н	3-I	2-CH ₃ -4-CF ₂ CF ₃	183.5-189.0
15	1044	n-C3H7	n-C3H7	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	142.6-145.4
	1045	CH2CH=CH2	CH₂CH	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	100.2-105.6
20			=CH ₂				
	1046	CH2CH2O	CH2CH2O	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	84.0-87.3
		-C2H5	-C ₂ H ₅				
25	1047	CH2CH2	CH2CH2	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	172.7-177.3
	1048	C2H5	C2H5	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	119.1
	1049	t-C₄H ₉	H	H	H	2-CH ₃ -4-OCBrF ₂	195-197
30	1050	t-C ₄ H ₉	H	H	3-C1	2-CH ₃ -4-OCBrF ₂	198-200
	1051	t-C₄H ₉	H	H	3-I	2-CH ₃ -4-OCBrF ₂	196-198
	1052	t-C₄H ₉	H	Н	H	4-C(CF ₃) ₂ OH	123-125
35	1053	t-C₄H ₉	H	Н	3-C1	4-C(CF ₃) ₂ OH	185-187
	1054	t-C ₄ H ₉	H	Н	3-I	4-C(CF ₃) ₂ OH	203-205
40	1055	i-C ₃ H ₇	H	н	3-I	2,4-F ₂	236-237
	1056	C ₂ H ₅	H	Н	3-I	2-CH ₃ -4-0CF ₂	176-178
						-CHF 2	,
45	1057	C ₂ H ₅	Н	Н	6-I	2-CH ₃ -4-0CF ₂	207-209
						-CHF 2	
	1058	n-C3H7	Н	H	3-I	2-CH ₃ -4-0CF ₂	185-187
50						-CHF 2	

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ym	Physical Properties (melting point: °C
,	1059	n-C3H7	Н	Н	6-I	2-CH ₃ -4-OCF ₂	215-217
	:					-CHF 2	
15	1060	t-C ₄ H ₉	Н	H	Н	2-CH ₃ -4-OCF ₂	197-198
			;			-CHF 2	
20	1061	t-C₄H ₉	H	H	3-C1	2-CH ₃ -4-OCF ₂	192-194
20						-CHF 2	
	1062	t-C ₄ H ₉	Н	H	3-İ	2-CH ₃ -4-OCF ₂	217-218
25	<u> </u>					-CHF 2	
	1063	i-C ₃ H ₇	H	H	3-C1	2-CH ₃ -4-0-(3,5	186-188
			i			-(CH ₃ O) ₂ -2-Pym)	
30	1064	i-C3H7	H	H	3-1	2-CH ₃ -4-0-(3,5	201-202
						-(CH ₃ O) ₂ -2-Pym)	
35	1065	t-C ₄ H ₉	H	H	H	3-0CF 2CF 20-4	156-158
	1066	t-C ₄ H ₉	H	H	3-C1	3-0CF 2CF 20-4	240-241
	1067	t-C ₄ H ₉	H	Н	3-I	3-0CF 2 CF 2 O-4	252-253
40	1068	СН 3 .	СНэ	H	3-I	2-CH3-4-CF2CF3	148.7
	1069	n-C3H7	СНз	H	3-1	2-CH3-4-CF2CF3	129.3
45	1070	CH2CH2O	CH2CH2	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	164.7
	1071	i-C ₃ H ₇	i-C ₃ H ₇	H	H	2-CH3-4-CF2CF3	169.1
	1072	i-C ₃ H ₇	i-C ₃ H ₇	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	201.2
50	1073	C2H5	H	H	3-I	2-CH ₃ -4-CF(CF ₃) ₂	194-195
:				·			

Table 1 (Cont'd)

10	No	R :	R2	R3	Xn	Ym	Physical Properties (melting point: °C
	1074	C ₂ H ₅	H	Н	6-I	2-CH ₃ -4-CF(CF ₃) ₂	218-220
	1075	n-C3H7	H	H	3-1	2-CH ₃ -4-CF(CF ₃) ₂	188-190
15	1076	n-C3H7	H	Н	1-6	2-CH ₃ -4-CF(CF ₃) ₂	201-203
	1077	i-C ₃ H ₇	H	Н	H	4-S0 ₂ CF ₃	184-186
20	1078	i-C ₃ H ₇	Н	Н	3-C1	4-S0 ₂ CF ₃	239-241
20	1079	i-C ₃ H ₇	Н	Н	3-I	4-S0 ₂ CF ₃	225-227
	1080	t-C ₄ H ₉	H	Н	3-I	4-S0 ₂ CF ₃	230-232
25	1081	i-C ₃ H ₇	i-C3H7	Н	3-1	2-CH3-4-CF2CF3	Paste
	1082	CH2CH2CH2	CH 2 CH 2	Н	3-I	2-CH3-4-CF2CF3	140.0-146.8
	1083	CH2CH2CH(CH3)CH2	Н	3-1	2-CH3-4-CF2CF3	171.4
30			-CH ₂ -				
1	1086	i-C3H7	H	Н	H	2-CH ₃ -4-0CF ₂ CF ₂	138-140
35		•				-Ph	
	1087	i-C3H7	Н	Н	3-C1	2-CH ₃ -4-0CF ₂ CF ₂	160-162
						-Ph	
40	1088	i-C3H7	H	Н	3-1	2-CH ₃ -4-OCF ₂ CF ₂	209-211
						-Ph	
45	1089	i-C3H7	Н	Н	3-	2-CH ₃ -4-OCF ₂ CF ₂	190-192
					NO ₂	-Ph	

Table 1 (Cont'd)

5	No	R1	R 2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1090	i-C ₃ H ₇	H	Н	Н	2-CH3-4-SCH2CH2	190-192
			ľ			-CF=CF 2	
15	1091	i-C ₃ H ₇	H	H	н	2-CH ₃ -4-SOCH ₂ CH ₂	149-153
						-CF=CF ₂	
	1092	i-C₃H7	H	Н	Н	2-CH3-4-SO2CH2	183-185
20						-CH ₂ CF=CF ₂	
	1093	i-C ₃ H ₇	H	Н	3-C1	2-CH ₃ -4-SCH ₂ CH ₂	168-170
						-CF=CF ₂	
25	1094	i-C ₃ H ₇	Н	H	3-C1	2-CH3-4-SOCH2CH2	164-167
						-CF=CF 2	
	1095	i-C3H7	H	H	3-C1	2-CH ₃ -4-SO ₂ CH ₂	181-183
30						-CH2CF=CF2	
	1096	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-SCH ₂ CH ₂	193-195
35						-CF=CF 2	
	1097	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-SOCH ₂ CH ₂	182-186
	1000		**		0.7	-CF=CF ₂	000 040
40	1098	i-C₃H₁	H	H	3-I	2-CH ₃ -4-SO ₂ CH ₂	208-210
	1000	: O II		**	**	-CH ₂ CF=CF ₂	040 040
	1099	i-C ₃ H ₇	H 	H	H	3-0CF ₂ 0-4	216-218
45	1100	i-C3H7	H	H	3-	3-0CF 20-4	227-229
					NO 2		
	1101	i-C ₃ H ₇	H	H	3-C1	3-0CF ₂ 0-4	243-245
50	1102	i-C ₃ H ₇	H	H	3-I	3-0CF 20-4	229-231

Table 1 (Cont'd)

5	No	R1	R 2	Rз	Xn	Ym	Physical Properties (melting
							point: ℃
10	1103	t-C ₄ H ₉	Н	H	H	3-0CF ₂ 0-4	209-211
	1104	t-C₄H₀	Н	Н	3-C1	3-0CF 20-4	206-208
15	1105	t-C₄H₀	Н	Н	3-1	3-0CF ₂ 0-4	228-230
15	1106	i-C ₃ H ₇	Н	н	Н	4-SCBrF ₂	175-177
	1107	i-C3H7	Н	H	H	4-SOCBrF ₂	158-161
20	1108	i-C ₃ H ₇	Н	H	3-NO ₂	4-SCBrF ₂	180-182
	1109	i-C ₃ H ₇	H	H	3-NO2	4-SOCBrF2	195-198
	1110	i-C ₃ H ₇	H	Н	3-C1	4-SCBrF ₂	156-158
25	1111	i-C ₃ H ₇	H	H	3-C1	4-SOCBrF2	218-220
	1112	i-C ₃ H ₇	- Н	H	3-1	4-SCBrF ₂	206-208
	1113	i-C3H7	H	H	3-I	4-SOCBrF ₂	158-160
30	1114	t-C ₄ H ₉	H	H	3-C1	4-SCBrF ₂	210-212
	1115	t-C ₄ H ₉	Н	H	3-I	4-SCBrF ₂	219-220
	1116	C2H5	C ₂ H ₅	H	3-I	2-CH3-4-CF2CF3	179.8-183.7
35	1117	CH2CH2CH2		H	3-1	2-CH ₃ -4-CF ₂ CF ₃	170.7
		-CH ₂	CH2CH2				
40	1118	C2H5	C ₂ H ₅	H	3-NO ₂	2-CH ₃ -4-0CF ₃	161.9
10	1119	C ₂ H ₅	C ₂ H ₅	H	3-NO2	2-CH ₃ -4-CF(CF ₃) ₂	169.1
	1120	СНз	СН₃	СНз	3-I	2-CH3-4-CF2CF3	141.9-146.6
45	1121	i-C ₃ H ₇	СН₃	СНз	3-I	2-CH3-4-CF2CF3	Paste
	1122	C ₂ H ₅	C2H5	СНз	3-I	2-CH ₃ -4-CF ₂ CF ₃	Paste
	1123	i-C ₃ H ₇	H	H	Н	4-SCF 3	135-137
50	1124	i-C ₃ H ₇	H	H	3-NO2	4-SCF ₃	187-189

Table 1 (Cont'd)

5	No	R1	R 2	R 3	Хn	Ym	Physical Properties (melting point: °C
10	1125	i-C ₃ H ₇	H	H	3-C1	4-SCF ₃	192-194
	1126	i-C3H7	H	H	3-I	4-SCF 3	194-196
15	1127	t-C₄H₀	H	H	3-1	4-SCF 3	195-197
15	1128	C2H5	C ₂ H ₅	H	3-I	4-SCF ₃	173-175
	1129	C ₂ H ₅	C ₂ H ₅	H	3-I	3-0CF 20-4	128-130
20	1130	C2H5	C ₂ H ₅	H	3-I	4-C(CF ₃) ₂ OH	152-154
	1131	C ₂ H ₅	C ₂ H ₅	H	3-NO2	2-CH ₃ -4-OCF ₃	178.7-182.6
	1132	C ₂ H ₅	C2H5	H	3-NO2	2-CH ₃ -4-OCF ₂ CHF ₂	160.8-165.0
25	1133	C2H5	C2H5	H	3-NO2	2-C1-4-CF ₂ CF ₃	91.9-95.2
	1134	C2H5	C ₂ H ₅	H	3-NO2	2-F-4-CF 2CF 3	162.6-166.8
	1135	C2H5	C ₂ H ₅	H	3-NO2	2-CH₃-4-Cl	188.8-190.8
30	1136	C ₂ H ₅	C2H5	H	3-NO2	4-0CF 3	185.7-187.9
	1137	C ₂ H ₅	C2H5	H	6-NO2	2-CH 3-4-OCF 2 CHF 2	111.2
	1138	C2H5	C ₂ H ₅	H	6-NO2	2-CH3-4-C1	149.7
35	1139	C2H5	C ₂ H ₅	H	6-NO2	4-0CF ₃	173.4
	1140	CH2CH(CH3)CH2	H	6-I	2-CH ₃ -4-CF ₂ CF ₃	166.4
40		-CH(C	H ₃)CH ₂				
40	1141	t-C ₄ H ₉	H	H	3-I	2-CH ₃ -4-CF ₃	197-198
	1142	i-C ₃ H ₇	H	Н	3-I	3-N=C(CF ₂ CF ₃)0-4	214-216
45	1143	t-C₄H ₉	H	H	3-I	3-N=C(CF ₂ CF ₃)0-4	253-254
	1144	C ₂ H ₅	C2H5	Н	3-I	2-CH3-4-CF3	160-161
	1145	i-C ₃ H ₇	Н	H	Ή	3-0CHFCF 20-4	102-104
50	1146	i-C ₃ H ₇	н	H	3-NO2	3-0CHFCF 20-4	190-192

Table 1 (Cont'd)

5	No	R1	R2	R 3	Xn	Ym	Physical Properties (melting point: °C
10	1147	i-C3H7	Н	H	3-C1	3-0CHFCF 20-4	123-125
	1148	i-C3H7	Н	H	3-I	3-0CHFCF 20-4	218-220
15	1149	t-C4H9	H	H	Н	3-0CHFCF 20-4	165-167
	1150	t-C4H9	н	H	3-I	3-0CHFCF 20-4	240-241
	1151	C ₂ H ₅	C2H5	H	3-I	3-0CHFCF 20-4	193-195
20	1152	t-C5H11	Н	H	3-F	2-CH ₃ -4-CF ₂ CF ₃	223.3
	1153	t-C5H11	Н	H	3-F	2-CH ₃ -4-	222
						CF(CF ₃) ₂	
25	1154	t-C ₅ H ₁₁	Н	H	3-F	2-CH ₃ -4-OCF ₃	193.6-195.8
	1155	t-C5H11	H	H	3-F	2-CH ₃ -4-0CHF ₂	165.5-174.0
	1156	n-C3H7	n-C3H7	H	3-I	2-CH ₃ -4-OCF ₃	132.2-135.0
30	1157	n-C3H7	n-C3H7	H	3-1	2-CH ₃ -4-OCHF ₂	81.4-87.8
	1158	n-C3H7	n-C3H7	H	3-I	2-CH ₃ -4-	116.3
						OCF 2 CHF 2	
35	1159	i-C ₃ H ₇	C 2H 5	H	3-I	2-CH ₃ -4-CF ₂ CF ₃	124.4
	1160	i-C3H7	C ₂ H ₅	H	3-I	4-0CF ₃	137.3-144.1
40	1161	i-C ₃ H ₇	- Н	H	3-I	3-0CF 2CHF0-4	161-163
	1162	i-C₃H₁	Н	H	3-NO ₂	3-0CF 2CHF0-4	238-240
	1163	i-C₃H₁	Н	H	3-C1	3-0CF 2CHF0-4	243-245
45	1164	i-C ₃ H ₇	Н	H	3-I	3-0CF 2CHF0-4	192-194
	1165	t-C ₄ H ₉	Н	H	H	3-0CF 2CHF0-4	205-207
	1166	t-C4H9	Н	H	3-I	3-0CF 2CHF0-4	238-240
50	1167	C ₂ H ₅	C ₂ H ₅	Н	3-I	3-0CF 2CHF0-4	195-197

55 .

Table 1 (Cont'd)

5	No	Rı	R2	R3	V.,	V_	Physical
	NO		n.	K.	Xn	Ym	Properties (melting
10	4400				 		point: ℃
	1168	i-C ₃ H ₇	. Н	Н	3-I	2-CH ₃ -4-SOCF ₃	148-152
	1169	t-C ₄ H ₉	H	Н	1-8	2-CH ₃ -4-SOCF ₃	165-168
15	1173	i-C ₃ H ₇	H	H	3-I	3-N=C(4-CF ₃ -Ph)	253-255
15						-0-4	
	1174	t-C ₄ H ₉	H	H	3-I	3-N=C(4-CF ₃ -Ph)	251-253
20					į	-0-4	
	1175	C ₂ H ₅	C ₂ H ₅	H	3-I	3-N=C(4-CF ₃ -Ph)	231-233
						-0-4	
25	1176	i-C ₃ H ₇	Н	Н	3-I	3-0-C(2-CF ₃ -Ph)	242-244
						=N-4	
	1177	t-C4H9	Н	H	3-I	3-0-C(2-CF ₃ -Ph)	229-231
30						=N-4	
	1178	C ₂ H ₅	C ₂ H ₅	H	3-I	3-0-C(2-CF ₃ -Ph)	203-205
						=N -4	
35	1179	C ₂ H ₅	C ₂ H ₅	C ₂ H ₅	3-1	2-CH3-4-CF2CF3	Paste
	1180	i-C ₃ H ₇	Н	H	3-I	3-0-C(CF ₂ CF ₃)	130-132
40						=N-4	
40	1181	t-C ₄ H ₉	Н	Н	3-I	3-0-C(CF ₂ CF ₃)	205-207
						=N-4	
45	1182	C ₂ H ₅	C ₂ H ₅	Н	3-I	3-0-C(CF ₂ CF ₃)	188-190
						=N-4	
	1183	i-C₃H₁	H	H	3-CF 3	2-CH ₃ -4-OCF ₃	222-224
50	1184	i-C3H7	Н	Н	3-CF 3	2-CH ₃ -4-CF ₂ CF ₃	219-221

Table 1 (Cont'd)

5	No	R1	R²	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1185	C ₂ H ₅	C ₂ H ₅	Н	3-CF 3	2-CH ₃ -4-OCF ₃	192-194
	1186	C2H5	C ₂ H ₅	H	3-CF 3	2-CH3-4-CF2CF3	218-220
	1187	i-C3H7	H	H	3-C1	2-F-4-0CF ₃	126-128
15	1188	i-C3H7	Н	H	3-I	2-F-4-0CF ₃	220-222
	1189	t-C4H9	H	H	3-I	2-F-4-0CF ₃	198-200
· 20	1190	C 2 H 5	C ₂ H ₅	Н	3-I	2-F-4-0CF ₃	129-131
	1191	i-C ₃ H ₇	Н	H	3-0CF ₃	2-CH3-4-CF2CF3	190-192
	1192	t-C ₄ H ₉	H	H	3-0CF ₃	2-CH3-4-CF2CF3	205-207
25	1193	C2H5	C 2H 5	H	3-0CF ₃	2-CH3-4-CF2CF3	146-148
	1202	i-C ₃ H ₇	Н	H	4-I	2-CH3-4-CF2CF3	197-199
	1203	i-C ₃ H ₇	Н	H	5-I	2-CH3-4-CF2CF3	201-203
30	1204	i-C ₃ H ₇	Н	H	4-I	2-CH 3-4-OCHF 2	241-243
	1205	i-C ₃ H ₇	H	H	5-I	2-CH3-4-0CHF2	214-216
	1206	i-C ₃ H ₇	H	H	3-CF 3	2-CH ₃ -4-OCF ₂ CHF ₂	195-197
35	1207	i-C ₃ H ₇	H	H	3-CF 3	2-CH ₃ -4-CF(CF ₃) ₂	227-229
	1208	i-C ₃ H ₇	H	H	Н	2-C ₂ H ₅ -4-0CF ₃	160-162
40	1209	i-C ₃ H ₇	H	H	3-C1	2-C ₂ H ₅ -4-0CF ₃	205-207
40	1210	i-C ₃ H ₇	H	H	3-I	2-C ₂ H ₅ -4-0CF ₃	241-243
	1211	t-C ₄ H ₉	H	H	3-I	2-C ₂ H ₅ -4-0CF ₃	224-225
45	1212	C ₂ H ₅	C ₂ H ₅	H	3-I	2-C ₂ H ₅ -4-0CF ₃	141-143
	1221	i-C ₃ H ₇	H	Н	3,4-Cl ₂	2-CH3-4-0CF3	199-200
	1222	i-C3H7	Н	Н	3,4-Cl ₂	2-CH3-4-CF2CF3	208-209
50	1223	i-C3H7	H	H	3,4-Cl ₂	2-CH ₃ -4-CF(CF ₃) ₂	228-229
					,		

Table 1 (Cont'd)

55

5			·				
10	No	R 1	R²	R 3	Xn	Ym	Physical Properties (melting point: °C
	1224	i-C3H7	Н	H	3,5-Cl ₂	2-CH3-4-0CF3	228-230
	1225	i-C3H7	Н	H	3,5-Cl ₂	2-CH3-4-CF2CF3	219-220
15	1226	i-C3H7	H	H	3,5-Cl ₂	2-CH ₃ -4-CF(CF ₃) ₂	211-212
	1227	i-C3H7	Н	H	3-C1-4-F	2-CH3-4-0CF3	184-186
20	1228	i-C3H7	H	H	3-C1-4-F	2-CH ₃ -4-CF ₂ CF ₃	178-180
20	1229	i-C3H7	Н	H	3-C1-4-F	2-CH ₃ -4-CF(CF ₃) ₂	200-201
	1230	t-C4H9	H	H	3-CF 3	2-CH3-4-0CF3	209-210
25	1231	t-C₄H ₉	н	Н	3-CF 3	2-CH3-4-CF2CF3	210-211
	1232	t-C ₄ H ₉	H	H	3-CF 3	2-CH ₃ -4-CF(CF ₃) ₂	242-243
	1233	i-C₃H7	Н	Н	3-0CF ₃	2-CH ₃ -4-OCF ₃	219-220
30	1234	t-C₄H9	Н	H	3-0CF ₃	2-CH ₃ -4-0CF ₃	222-223
	1235	C ₂ H ₅	C ₂ H ₅	Н	3-0CF ₃	2-CH ₃ -4-0CF ₃	125-126
35	1236	i-C₃H₁	Н	H	3-0CF ₃	2-CH ₃ -4-CF(CF ₃) ₂	235-236
	1237	t-C₄H _B	Н	H	3-0CF ₃	2-CH ₃ -4-CF(CF ₃) ₂	220-222
	1238	C ₂ H ₅	C 2H 5	H	3-0CF 3	2-CH ₃ -4-CF(CF ₃) ₂	156-157
40	1245	i-C₃H₁	Н	Н	3-CN	2-CH3-4-CF2CF3	168-170
	1246	i-C3H7	Н	Н	4-I	2-CH3-4-0CF3	238-240
45	1247	i-C₃H₁	Н	Н	5-I	2-CH3-4-0CF3	205-206
	1248	i-C3H7	Н	Н	4-I	2-CH3-4-OCF2CHF2	222-223

Table 1 (Cont'd)

							Ť
5	No	R1	R2	R³	Xn	Ym	Physical Properties (melting point: °C
10	1249	i-C ₃ H ₇	Н	Н	5-I	2-CH ₃ -4-0CF ₂ CHF ₂	203-204
	1250	i-C ₃ H ₇	H	Н	4-I	2-CH ₃ -4-CF(CF ₃) ₂	215-216
15	1251	i-C ₃ H ₇	Н	Н	5-I	2-CH ₃ -4-CF(CF ₃) ₂	216-217
15	1256	i-C₃H7	H	H	3-C1	2-CH3-4-CF2CF3	235-236
					-4-F		
20	1257	t-C ₄ H ₉	H	H	3-C1	2-CH3-4-CF2CF3	225-226
					-4-F		
	1258	C ₂ H ₅	C ₂ H ₅	H	3-C1	2-CH ₃ -4-CF ₂ CF ₃	155-156
25					-4-F		
	1259	i-C3H7	Н	H	3-C1	2-CH ₃ -4-0CF ₃	229-231
		,			-4-F		
30	1260	t-C₄H ₉	H	H	3-C1	2-CH ₃ -4-0CF ₃	237-238
					-4 - F		
	1261	C ₂ H ₅	C ₂ H ₅	H	3-C1	2-CH ₃ -4-0CF ₃	140-141
35					-4-F		
	1262	i-C ₃ H ₇	н	Н	3-C1	2-CH ₃ -4-	264-265
40			ļ		-4-F	CF(CF ₃) ₂	
	1263	t-C₄H ₉	Н	H	3-C1	2-CH ₃ -4-	253-154
3					-4-F	CF(CF ₃) ₂	
45	1264	C2H5	C ₂ H ₅	H	3-C1	2-CH3-4-	158-159
					-4-F	CF(CF ₃) ₂	
	1266	i-C₃H₁	Н	H	3,4-	2-CH ₃ -4-	162-164
50					Br ₂	CF 2 CF 3	
							·

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ym	Physical Properties (melting
							point: °C
10	1277	i-C ₃ H ₇	Н	Н	4-C1	2-CH ₃ -4-	185-186
		1	ĺ			CF 2 CF 3	
	1278	t-C ₄ H ₉	Н	H	4-Cl	2-CH ₃ -4-	206-207
15						CF 2 CF 3	
	1280	C2H5	C ₂ H ₅	H	4-C1	2-CH ₃ -4-	163-164
					•	CF 2 CF 3	
20	1281	C2H5	C ₂ H ₅	Н	4-C1	2-CH ₃ -4-	193-194
					- 6−I	CF 2 CF 3	
25	1283	i-C ₃ H ₇	H	H	3,4-F ₂	2-CH3-4-OCF3	194-195
25	1284	t-C ₄ H ₉	Н	H	3,4-F ₂	2-CH ₃ -4-OCF ₃	216-217
	1285	C2H5	C ₂ H ₅	H	3,4-F ₂	2-CH3-4-OCF3	156-157
30	1287	i-C ₃ H ₇	H	H	4,5-F ₂	2-CH3-4-OCF3	195-196
•	1288	t-C ₄ H ₉	H	H	4,5-F ₂	2-CH ₃ -4-OCF ₃	223-224
	1290	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-0C	226-227
35						-(CF ₂ CF ₃)=C	
		a				-(CF ₃) ₂	•
	1291	i-C ₃ H ₇	H	H	3-C1	2-CH ₃ -4-0C	204-205
40						-(CF 2CF 3)=C	
						-(CF ₃) ₂	
	1292	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-0C	198-199
45						-(OCH ₃)=C	
						-(CF ₃) ₂	
:	1293	i-C ₃ H ₇	Н	H	3-C1	2-CH ₃ -4-OC	179-180
50						-(OCH ₃)=C	
						-(CF ₃) ₂	

Table 1 (Cont'd)

5	No	Rı	R2	Ra	Xn	Ym	Physical Properties (melting point: °C
10	1294	CH(CH₃)CH₂OH	Н	H	H	2-CH ₃ -4-C ₂ F ₅	73-74
	1295	i-C3H7	Н	H	6-C1	2-0CH₃-5-Ph	120
15	1296	i-C3H7	Н	Н	3-C1	2-0CH3-5-Ph	195
15	1297	n-C 3H 7	H	Н	6-C1	2-0CH3-5-Ph	200
	1298	CH(CH3)CH2OH	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	195
20	1299	CH(C2H5)CH2OH	Н	H	H	2-CH3-4-C2F5	78
	1300	CH(CH₃)CH₂OH	Н	Н	3-I	2-CH ₃ -4-C ₂ F ₅	98-99
	1301	i-C3H7	H	H	3-C1	2-CH ₃ -4-C≡C	210
25						-C₄H₀-t	
	1302	i-C ₃ H ₇	Н	н	6-C1	2-CH ₃ -4-C≡C	205
, .						-C ₄ H ₉ -t	
30	1303	n-C ₃ H ₇	Н	Н	3-1	2-CH ₃ -4-C ₂ F ₅	200
	1304	n-C ₃ H ₇	H	Н	6-I	2-CH ₃ -4-C ₂ F ₅	195
	1305	i-C ₃ H ₇	H	H	3-I	$2-CH_3-4-C \equiv C$	205
35						-C₄H₃-t	
·	1306	i-C ₃ H ₇	Н	H	6-I	$2-CH_3-4-C\equiv C-$	170
40		-				-C4H9-t	
	1307	CH2-Ph	H	H	3-C1	2-CH ₃ -4-C ₂ F ₅	175
	1308	CH2-Ph	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	175
45	1309	CH ₂ -(2-Cl-Ph)	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	170
	1310	CH ₂ -(2-C1-Ph)	Н	Н	6-C1	2-CH3-4-C2F5	210
	1311	CH ₃	Н	Н	3-I	2-CH3-4-C2F5	190
50	1312	CH ₃	Н	н	1-6	2-CH ₃ -4-C ₂ F ₅	200

Table 1 (Cont'd)

5	No	R1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	1313	C2H5	H	Н	3-I	2-CH3-4-C2F5	182
	1314	C2H5	H	Н	6-I	2-CH3-4-C2F5	205
	1315	CH2CH(OH)CH3	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	187
15	1316	CH(C ₂ H ₅)CH ₂ OH	H	H	3-C1	2-CH3-4-C2F5	208
	1317	C(CH ₃) ₂ CH ₂ OH	H	H	3-C1	2-CH ₃ -4-C ₂ F ₅	181-182
20	1318	CH2CH(OH)C2H5	H	Н	3-C1	2-CH3-4-C2F5	171-172
	1319	CH2CH2-Ph	H	Н	3-C1	2-CH3-4-C2F5	150
	1320	CH2CH2-Ph	Н	Н	6-C1	2-CH3-4-C2F5	190
25	1321	CH(CH ₃)-Ph	Н	Н	3-C1	2-CH3-4-C2F5	160
	1322	CH(CH ₃)-Ph	H	H	6-C1	2-CH3-4-C2F5	190
	1323	i-C3H7	H	Н	3-C1	2-CH3-4	220
30						-CH ₂ CH ₂ C(CH ₃) ₃	
	1324	i-C3H7	Н	Н	6-C1	2-CH ₃ -4	205
	'					-CH ₂ CH ₂ C(CH ₃) ₃	
35	1325	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-C≡C-Ph	215
	1326	i-C3H7	Н	Н	6-C1	2-CH ₃ -4-C≡C-Ph	230
40	1327	0-n-C ₃ H ₇	Н	H	3-C1	2-CH ₃ -4-C ₂ F ₅	165
	1328	0-n-C ₃ H ₇	H	H	6-C1	2-CH3-4-C2F5	150
	1329	0-CH2CH=CHC1	Н	Н	3-C1	2-CH3-4-C2F5	150
45		(E)					
	1330	i-C3H7	H	Н	3-C1	2-CH ₃ -4-CN	230
	1331	(CH ₂) ₃ -Ph	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	112
50	1332	(CH ₂) ₃ -Ph	Н	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	105
		w.					

Table 1 (Cont'd)

5	No	R1	R2	R³	Xn	Ym	Physical Properties (melting point: °C
10	1333	CH ₂ (4-Cl-Ph)	H	H	3-C1	2-CH ₃ -4-C ₂ F ₅	198
	1334	CH2(4-C1-Ph)	Н	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	156
*	1335	CH ₂ (3-Cl-Ph)	Н	H	3-C1	2-CH3-4-C2F5	168
15	1336	CH ₂ (3-C1-Ph)	Н	H	6-C1	2-CH3-4-C2F5	177
	1337	CH ₂ (2-CH ₃ -Ph)	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	152
20	1338	CH ₂ (2-CH ₃ -Ph)	H	Н	6-C1	2-CH3-4-C2F5	147
	1339	CH ₂ (3-CH ₃ -Ph)	H	Н	3-C1	2-CH3-4-C2F5	Crystals
	1340	CH ₂ (3-CH ₃ -Ph)	Н	Н	6-C1	2-CH3-4-C2F5	173
25	1341	CH ₂ (4-CH ₃ -Ph)	Н	Н	3-C1	2-CH3-4-C2F5	175
	1342	CH ₂ (4-CH ₃ -Ph)	Н	H	6-C1	2-CH ₃ -4-C ₂ F ₅	Crystals
	1343	CH ₂ (2-CH ₃ 0-Ph)	Н	Н	3-C1	2-CH3-4-C2F5	Crystals
30 .	1344	CH ₂ (2-CH ₃ 0-Ph)	Н	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	176
	1345	$CH_2(3-CH_3O-Ph)$	H	H	3-C1	2-CH ₃ -4-C ₂ F ₅	73
	1346	CH ₂ (3-CH ₃ 0-Ph)	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	86
35	1347	CH ₂ (4-CH ₃ 0-Ph)	Н	H	3-C1	2-CH ₃ -4-C ₂ F ₅	169
	1348	CH ₂ (4-CH ₃ 0-Ph)	H	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	168
	1349	CH ₂ (2,4-Cl ₂ -Ph)	H	н	3-C1	2-CH ₃ -4-C ₂ F ₅	169
40	1350	CH ₂ (2,4-Cl ₂ -Ph)	H	н	6-C1	2-CH ₃ -4-C ₂ F ₅	205
	1351	CH ₂ (3,4-Cl ₂ -Ph)	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	179
45	1352	CH ₂ (3,4-Cl ₂ -Ph)	н	н	6-C1	2-CH ₃ -4-C ₂ F ₅	192
	1353	CH ₂ (2,3-Cl ₂ -Ph)	н	H	3-C1	2-CH ₃ -4-C ₂ F ₅	179
	1354	CH ₂ (2,3-Cl ₂ -Ph)	н	H	6-C1	2-CH ₃ -4-C ₂ F ₅	208
50	1355	CH ₂ -2-Pyi	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	143
· [

Table 1 (Cont'd)

5	No	R:	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	1356	(CH ₂) ₂ (2-Cl-Ph)	H	H	3-C1	2-CH ₃ -4-C ₂ F ₅	141
	1357	(CH ₂) ₂ (2-C1-Ph)	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1358	(CH ₂) ₂ (3-C1-Ph)	H	H	3-C1	2-CH ₃ -4-C ₂ F ₅	117
15	1359	$(CH_2)_2(3-C1-Ph)$	H	H	6-C1	2-CH3-4-C2F5	Paste
	1360	(CH ₂) ₂ (4-Cl-Ph)	H	Н	3-C1	2-CH3-4-C2F5	118
20	1361	(CH ₂) ₂ (4-C1-Ph)	H	Н	6-Cl	2-CH ₃ -4-C ₂ F ₅	138
	1362	CH(CH ₃)(2-Cl-Ph)	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1363	CH(CH ₃)(2-C1-Ph)	H	Н	6-C1	2-CH3-4-C2F6	197
25	1364	CH(CH ₃)(3-C1-Ph)	H	H	3-C1	2-CH3-4-C2F5	100
	1365	CH(CH ₃)(3-C1-Ph)	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	Crystals
	1366	CH(CH ₃)(4-Cl-Ph)	H	Н	3-C1	2-CH3-4-C2F5	195
30	1367	CH(CH ₃)(4-Cl-Ph)	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1368	$(CH_2)_2O(2-C1-Ph)$	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	162
	1369	(CH ₂) ₂ 0(2-Cl-Ph)	H	H	6-C1	2-CH3-4-C2F5	160
35	1370	(CH ₂) ₂ 0(3-Cl-Ph)	H	H	3-C1	2-CH3-4-C2F5	115
	1371	(CH ₂) ₂ 0(3-Cl-Ph)	Н	H	6-C1	2-CH3-4-C2F5	172
40	1372	(CH ₂) ₂ 0(4-Cl-Ph)	H	н	3-C1	2-CH3-4-C2F5	185
40	1373	(CH ₂) ₂ 0(4-Cl-Ph)	H	Н	6-C1	2-CH3-4-C2F5	148
	1374	(CH ₂) ₂ 0-Ph	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	154
45	1375	(CH ₂) ₂ 0-Ph	H	Н	6-C1	2-CH3-4-C2F5	183
•	1376	(CH ₂) ₂ NH-Ph	Н	Н	3-C1	2-CH3-4-C2F5	104
	1377	(CH ₂) ₂ NH-Ph	H	H	6-C1	2-CH3-4-C2F5	Paste
50	1378	CH(CH ₃)CH ₂ OH	H	H	6-C1	2-CH3-4-C2F5	192

Table 1 (Cont'd)

5	No	K1	R ²	R³	Xn	Ym	Physical Properties (melting point: °C
10	1379	CH(Ph)CH2OH	Н	H	H	2-CH3-4-C2F5	100-101
	1380	CH(4-t-C ₄ H ₉ -Ph)	Н	Н	Н	2-CH3-4-C2F5	107-108
		-CH₂OH					
15	1381	C(CH ₃) ₂ CH ₂ OH	Н	Н	H	2-CH3-4-C2F5	227
	1382	i-C ₃ H ₇	Н	Н	3-C1	2-F-4-C ₂ F ₅	190
20	1383	i-C3H7	H	Н	3-C1	2-C1-4-C ₂ F ₅	180
	1384	i-C ₃ H ₇	Н	H	3-C1	2-CF3-4-C2F5	235
	1385	i-C3H7	Н	Н	3-I	2-F-4-C ₂ F ₅	190
25	1386	i-C ₃ H ₇	H	H	3-I	2-C1-4-C ₂ F ₅	200
	1387	i-C ₃ H ₇	H	H	3-I	2-CF ₃ -4-C ₂ F ₅	255
	1388	i-C3H7	H	H	1-8	2-0CH ₃ -4-C ₂ F ₅	152
30	1389	i-C3H7	H	Н	3-I	2-CH3-4-CN	215
	1390	2-Fur	H	H	3-C1	2-CH3-4-C2F5	178
	1391	2-Fur	H	Н	6-C1	2-CH3-4-C2F5	149
35	1392	2-TetFur	H	H	3-C1	2-CH3-4-C2F5	153
	1393	2-TetFur	H	H	6-Cl	2-CH3-4-C2F5	130
40	1394	CH2-4-Pyi	H	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	88
	1395	CH2-4-Pyi	H	н	6-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1396	(CH ₂) ₃ OH	Н	Н	H	2-CH3-4-C2F5	83-84
45	1397	(CH ₂) ₂ OH	Н	н	H	2-CH ₃ -4-C ₂ F ₅	136
	1398	CH2CH(OH)CH2Ph	Н	н	Н	2-CH ₃ -4-C ₂ F ₅	77-78
;	1399	(CH ₂) ₃ OH	Н	H	3-C1	2-CH ₃ -4-C ₂ F ₅	188
50	1400	CH2-Ph	Н	H	3-I	2-CH ₃ -4-C ₂ F ₅	172
Į							

Table 1 (Cont'd)

5							Physical
	No	R1	R2	Ŗз	Хn	Ym	Properties
							(melting
.10							point: °C
	1401	CH2-Ph	H	H	6-I	2-CH ₃ -4-C ₂ F ₅	212
*	1402	CH ₂ (2-Cl-Ph)	H	H	3-I	2-CH ₃ -4-C ₂ F ₅	136
15	1403	CH ₂ (2-Cl-Ph)	H	Н	6-I	2-CH ₃ -4-C ₂ F ₅	214
	1404	CH ₂ (2-CH ₃ -Ph)	Н	H	3-I	2-CH ₃ -4-C ₂ F ₅	100
_	1405	CH ₂ (2-CH ₃ -Ph)	Н	H	6-I	2-CH ₃ -4-C ₂ F ₅	185
20	1406	CH ₂ -Ph	СНз	H	3-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1407	CH2-Ph	CH2-Ph	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	136
25	1408	CH2-Ph	CH2-Ph	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1409	i-C ₃ H ₇	H	H	3-I	2-C ₂ F ₅ -4-Br	250
	1410	i-C ₃ H ₇	H	H	1-8	2-C ₂ F ₅ -4-C ₂ F ₅	245
30	1411	CH₂C≡CH	Н	H	H	2-CH ₃ -4-C ₂ F ₅	133-135
	1412	CH(4-Ph-Ph)CH ₂	Н	H	3-C1	2-CH3-4-C2F5	112
35 .	:	-ОН					
	1414	$C(CH_3)_2C\equiv CH$	Н	Н	Н	2-CH3-4-C2F5	207
	1415	C(CH ₃) ₂ CH ₂ OH	H	H	6-C1	2-CH3-4-C2F5	231
40	1416	CH(4-Cl-Ph)CH ₂	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	225
		-ОН					
45	1417	C(CH ₃) ₂ -Ph	H	Н	3-C1	2-CH3-4-C2F5	190
45	1418	C(CH ₃) ₂ CH ₂ -Ph	Н	Н	3-C1	2-CH3-4-C2F5	192
:	1419	CH ₂ -3-Pyi	H	Н	3-C1	2-CH3-4-C2F5	Paste
50	1420	CH ₂ -3-Pyi	Н	Н	6-C1	2-CH3-4-C2F5	Paste

Table 1 (Cont'd)

5	No	B1	R2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1421	CH2-Ph	Н	Н	3-C1	2-CH ₃ -4-0CHF ₂	187
	1422	CH2-Ph	Н	Н	6-C1	2-CH 3-4-0CHF 2	198
	1423	CH2-(2-Cl-Ph)	H	Н	3-C1	2-CH3-4-0CHF2	178
15	1424	CH ₂ -(2-Cl-Ph)	Н	H	6-C1	2-CH3-4-0CHF2	192
	1425	CH2-(2-CH3-Ph)	H	Н	3-C1	2-CH3-4-0CHF2	183
20	1426	CH2-(2-CH3-Ph)	H	Н	6-C1	2-CH 3-4-0CHF 2	192
	1427	t-C ₄ H ₉	H	Н	3-I	2-F-4-C ₂ F ₅	220
	1428	t-C ₄ H ₉	H	Н	3-I	2-Cl-4-C ₂ F ₅	187
25	1429	t-C ₄ H ₉	H	Н	3-I	2-CF ₃ -4-C ₂ F ₅	240
	1430	CH ₂ -Ph	H	н	3-I	2-CH ₃ -4-0CHF ₂	176
	1431	CH2-Ph	H	Н	6-I	2-CH3-4-0CHF2	196
30	1432	CH2-(2-Cl-Ph)	H	Н	3-I	2-CH 3-4-0CHF 2	189
	1433	CH ₂ -(2-C1-Ph)	H	Н	6-I	2-CH3-4-0CHF2	227
	1434	CH ₂ -(2-CH ₃ -Ph)	H	H	3-I	2-CH ₃ -4-0CHF ₂	215
35	1435	CH ₂ -(2-CH ₃ -Ph)	H	H	6-I	2-CH3-4-0CHF2	209
	1436	CH2-Ph	СН з	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	Paste
40	1437	CH2-Ph	СНз	H	3-C1	2-CH ₃ -4-0CHF ₂	Paste
40	1438	CH2-Ph	СНз	H	3-I	2-CH ₃ -4-C ₂ F ₅	175
	1439	CH2-Ph	СНз	Н	6-I	2-CH ₃ -4-C ₂ F ₅	Paste
45	1440	CH2-Ph	СНз	H	3-I	2-CH ₃ -4-0CHF ₂	Paste
	1441	CH(C ₂ H ₅)CH ₂ OH	Н	H	6-C1	2-CH ₃ -4-C ₂ F ₅	213
	1442	(R)-C*H(Ph)	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	105-107
50		-CH 2 OH					

Table 1 (Cont'd)

5	No	R 1	R2	R 3	Xn	Ym	Physical Properties (melting point: °C
	1443	(R)-C*H(Ph)	H	H	6-C1	2-CH3-4-C2F5	145-146
		-CH₂OH					
15	1445	(S)-C+H(CH₃)	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	93-95
		-CH₂OH					
	1446	(S)-C*H(CH ₃)	Н	H	6-Cl	2-CH3-4-C2F5	93-95
20		-СН 2 ОН					
	1447	t-C ₄ H ₉	Н	Н	3-C1	4-C ₂ F ₅	275
25	1448	t-C4H9	Н	Н	3-C1	2-F-4-C ₂ F ₅	225
	1449	t-C₄H ₉	H	Н	3-C1	2-C1-4-C ₂ F ₅	200
	1450	n-C ₃ H ₇	Н	Н	3-I	2-CH ₃ -4-OCHF ₂	181
30	1451	n-C3H7	H	Н	6-I	2-CH3-4-OCHF2	233
	1452	c-C3H5	H	Н	3-I	2-CH3-4-OCHF2	182
35	1453	c-C3H5	H	Н	6-I	2-CH3-4-0CHF2	231
	1454	s-C4H9	H	Н	3-I	2-CH3-4-0CHF2	225
	1455	s-C4H9	H	Н	6-I	2-CH3-4-0CHF2	244
40	1456	CH₂C≡CH	Н	Н	1-8	2-CH 3-4-0CHF 2	196
	1457	CH2-Ph	C ₂ H ₅	Н	3-C1	2-CH3-4-C2F5	Paste
	1458	(R)-C*H(CH₃)	H	Н	3-C1	2-CH3-4-OCHF2	136
45		-Ph					
	1459	(S)-C+H(CH ₃)	H	H	3-C1	2-CH3-4-OCHF2	136
50		-Ph					

Table 1 (Cont'd)

5	No	R1	R2	R3	Xn	Ym	Physical Properties
10							(melting point: °C
	1460	(R)-C*H(CH ₃)	H	H	3-C1	2-CH3-4-C2F5	94-95
		-CH 2 OH				·	
15	1461	$(R)-C+H(CH_3)$	H	H	6-C1	2-CH ₃ -4-C ₂ F ₅	94-95
		-CH 2 OH					
20	1464	C(CH ₃) ₂ CH ₂ OH	H	H	3-I	2-CH3-4-C2F5	118
	1465	CH(CH ₃)CH ₂ OH	H	Н	6-I	2-CH3-4-C2F5	130-131
	1466	$C(CH_3)_2C\equiv CH$	Н	Н	3-C1	2-CH3-4-C2F5	210-211
25	1467	C(CH ₃) ₂ C≡CH	Н	H	6-C1	2-CH ₃ -4-C ₂ F ₅	230
	1468	CH ₂ (2-F-Ph)	Н	Н	3-C1	2-CH3-4-C2F5	187
30	1469	CH ₂ (2-F-Ph)	Н	H	6-C1	2-CH ₃ -4-C ₂ F ₅	165
30	1470	CH2-Ph	Н	H	3-F	2-CH3-4-C2F5	158
	1471	CH2-Ph	Н	Н	6-F	2-CH3-4-C2F5	134
35	1472	s-C4H9	H	Н	3-I	2-F-4-C ₂ F ₅	200
	1473	s-C4H9	Н	Н	3-I	2-C1-4-C ₂ F ₅	205
	1474	i-C ₃ H ₇	Н	H	3-I	2-F-4-n-C3F7	165
40	1475	t-C4H9	Н	Н	3-I	2-C ₂ H ₅ -4-C ₂ F ₅	235
	1476	CH2CH(OH)Ph	Н	Н	3-C1	2-CH3-4-C2F5	108
45	1477	CH2CH(OH)Ph	Н	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	105
	1478	C(CH ₃) ₂ C≡CH	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	105
	! 						

Table 1 (Cont'd)

.	No	R:	R²	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1479	C(CH ₃) ₂ C≡C	Н	Н	3-C1	2-CH3-4-C2F5	110
		-2-Thi					
	1480	C(CH ₃) ₂ C≡C-Ph	H	Н	6-C1	2-CH3-4-C2F5	194
15	1481	(R)-C*H(CH ₃)	H	Н	3-I	2-CH3-4-C2F5	103-105
		-CH 2 OH					
20	1482	(S)-C*H(CH₃)	Н	Н	3-I	2-CH3-4-C2F5	103-105
		-CH 2 OH					
	1483	(R)-C+H(CH₃)	Н	Н	6-I	2-CH3-4-C2F5	173-174
25		-CH 2 OH			4		
	1484	C(CH ₃) ₂ (4-Cl	Н	H	3-C1	2-CH3-4-C2F5	218
		-Ph)					
30	1485	C(CH ₃) ₂ (3-C1	Н	Н	3-C1	2-CH3-4-C2F5	128
		-Ph')				•	
	1486	CH2-Ph	Н	H	3-C1	2-F-4-C ₂ F ₅	162
35	1487	CH2-Ph	H	Н	3-C1	2-C1-4-C ₂ F ₅	153
	1488	C ₂ H ₅	H	Н	3-C1	2-F-4-C ₂ F ₅	135
40	1489	C2H5	H	H	3-C1	2-C1-4-C ₂ F ₅	125
	1490	C ₂ H ₅	Н	H	3-C1	2-F-4-n-C ₃ F ₇	128
	1491	n-C3H7	H	H	3-C1	2-F-4-C ₂ F ₅	153
45	1492	n-C3H7	Н	H	3-C1	2-C1-4-C ₂ F ₅	147
	1493	n-C3H7	H	H	3-C1	2-F-4-n-C ₃ F ₇	142
	1494	i-C ₃ H ₇	H	Н	3-C1	2-F-4-n-C ₃ F ₇	142
50	1495	i-C ₃ H ₇	H	H	3-C1	2-C ₂ H ₅ -4-C ₂ F ₅	213
		1.3					

Table 1 (Cont'd)

<i>5</i>	No	R:	R2	Кз	Xn	Ym	Physical Properties (melting point: °C
10	1496	t-C4H9	Н	Н	3-C1	2-F-4-n-C ₃ F ₇	172
	1497	t-C ₄ H ₉	Н	Н	3-C1	2-C ₂ H ₅ -4-C ₂ F ₅	194
15	1498	s-C4H9	Н	Н	3-C1	2-F-4-C ₂ F ₅	209
	1499	s-C4H9	Н	Н	3-C1	2-C1-4-C ₂ F ₅	194
	1500	s-C ₄ H ₉	H	Н	3-C1	2-F-4-n-C ₃ F ₇	182
20	1501	s-C ₄ H ₉	H	H	3-C1	2-C ₂ H ₅ -4-C ₂ F ₅	212
	1502	C2H5	Н	Н	3-I	2-F-4-C ₂ F ₅	135
	1503	C2H5	Н	Н	3-I	2-C1-4-C ₂ F ₅	155
25	1504	t-C ₄ H ₉	Н	Н	3-I	2-F-4-n-C ₃ F ₇	180
*	1505	t-C ₄ H ₉	H	H	3-F	2-CH ₃ -4-C ₂ F ₅	220
30	1506	t-C ₄ H ₉	H	H	3-F	2-CH ₃ -4-0CHF ₂	186
	1507	t-C ₄ H ₉	H	H	3-F	2-F-4-C ₂ F ₅	214
	1508	t-C ₄ H ₉	H	Н	3-F	2-C1-4-C ₂ F ₅	222
35	1509	t-C ₄ H ₉	H	Н	3-F	2-F-4-n-C ₃ F ₇	179
	1510	C ₂ H ₅	H	H	3-F	2-F-4-C ₂ F ₅	125
	1511	C ₂ H ₅	H	H	6-F	2-F-4-C ₂ F ₅	155
40	1512	n-C ₃ H ₇	H	H	3-F	2-F-4-C ₂ F ₅	130
	1513	n-C3H7	H	H	6-F	2-F-4-C ₂ F ₅	170
45	1514	i-C ₃ H ₇	H	H	3-F	2-F-4-C ₂ F ₅	190
45	1515	i-C ₃ H ₇	Н	H	6-F	2-F-4-C ₂ F ₅	180
	1516	i-C ₃ H ₇	H	H	3-F	2-C1-4-C ₂ F ₅	210
50	1517	i-C ₃ H ₇	H	Н	6-F	2-C1-4-C ₂ F ₅	160

Table 1 (Cont'd)

5	No	R1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	1518	(S)-C+H(CH ₃)	H	H	6-I	2-CH ₃ -4-C ₂ F ₅	173-174
		-CH 2 OH			:		
15	1519	C(CH ₃) ₂ CH ₂ OH	H	Н	3-1	2-CH3-4-0CF3	205
,,	1520	C(CH3)2CH2OH	H	H	6-I	2-CH3-4-0CF3	248
	1521	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-(4-CF ₃ 0	247-250
20						-Ph)	
•	1522	i-C ₃ H ₇	H	H _.	3-I	2-CH ₃ -4-(4-CF ₃	243-246
						-Ph)	
25	1523	CH ₂ (2-CF ₃ -Ph)	H	Н	3-C1	2-CH3-4-C2F5	183
	1524	n-C ₃ H ₇	H	Н	3-I	2-F-4-n-C ₃ F ₇	145
	1525	C2H5	C ₂ H ₅	Н	3-F	2-CH3-4-C2F5	135
30	1526	C ₂ H ₅	C ₂ H ₅	Н	3-F	2-CH3-4-i-C3F7	150
	1527	C ₂ H ₅	C ₂ H ₅	Н	3-F	2-CH ₃ -4-0CF ₃	125
35	1528	C ₂ H ₅	C ₂ H ₅	Н	3-F	2-CH3-4-OCHF2	110
55	1529	C ₂ H ₅	C ₂ H ₅	H	3-F	2-CH3-4-OCF2CHF2	155
	1530	C ₂ H ₅	C ₂ H ₅	Н	3-F	2-F-4-C ₂ F ₅	130
40	1531	C ₂ H ₅	C ₂ H ₆	Н	3-F	2-C1-4-C ₂ F ₅	110
	1532	C ₂ H ₅	C ₂ H ₅	Н	3-I	2-CH3-4-i-C3F7	142
	1533	C2H5	C ₂ H ₅	Н	3-I	2-CH3-4-OCF3	142
45	1534	C ₂ H ₅	C ₂ H ₅	Н	3-I	4-0CF 3	142
	1535	C ₂ H ₅	C ₂ H ₅	Н	3-C1	2-CH3-4-C2F5	150
·	1536	C2H5	C ₂ H ₅	Н	3-C1	2-CH ₃ -4-OCF ₃	123
50	1537	C ₂ H ₅	C ₂ H ₅	Н	3-C1	2-CH3-4-i-C3F7	147

Table 1 (Cont'd)

5	No	R1	R 2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1538	C2H5	C ₂ H ₅	Н	3-C1	2-CH3-4-0CHF2	92
	1539	C ₂ H ₅	C ₂ H ₅	Н	3-C1	2-CH3-4-OCF2CHF2	135
15	1540	C2H5	C ₂ H ₅	Н	3-C1	2-C1-4-C ₂ F ₅	110
15	1541	C2H5	C ₂ H ₅	Н	3-C1	2-F-4-C ₂ F ₅	113
	1542	C2H5	C ₂ H ₅	н	3-C1	2-CH3-C1	142
20	1543	C ₂ H ₅	C ₂ H ₅	Н	3-C1	2-C ₂ H ₅ -4-C ₂ F ₅	101
	1544	C2H5	C ₂ H ₅	Н.	3-C1	4-0CF 3	138
	1545	C ₂ H ₅	C ₂ H ₅	Н	3-C1	4-CF 3	188
25	1546	C2H5	C ₂ H ₅	н	3-F	2-CH3-4-Cl	135
	1547	C ₂ H ₅	C ₂ H ₅	Н	3-F	4-CF 3	175
	1548	C ₂ H ₅	C ₂ H ₅	Н	3-F	4-0CF ₃	155
30	1549	C ₂ H ₅	C ₂ H ₅	Н	3-F	2-C ₂ H ₅ -4-C ₂ F ₅	80
	1550	C ₂ H ₅	C ₂ H ₅	H	3-NO2	2-CH3-4-C2F5	185
	1551	C ₂ H ₅	C ₂ H ₅	H	6-NO2	2-CH3-4-C2F5	145
35	1552	t-C₄H ₉	H	H	3-I	3-CH3-4-C2F5	215
	1553	CH2-Ph	CH 3	СНз	3-C1	2-CH3-4-C2F5	Paste
40	1554	CH(CH ₃)-Ph	H	СНз	3-C1	2-CH ₃ -4-C ₂ F ₅	Paste
	1555	C2H5	C ₂ H ₅	H	3-I	2-CH ₃ -4-0CHF ₂	138-139
	1556	C ₂ H ₅	C ₂ H ₅	H	3-I	2-CH ₃ -4-0CF ₂ CHF ₂	136
45	1557	C2H5	C ₂ H ₅	H	3-I	2-CH3-4-Cl	179
	1558	C2H5	C ₂ H ₅	H	3-I	4-CF 3	187
	1559	C ₂ H ₅	C ₂ H ₅	H	3-I	2-C ₂ H ₅ -4-C ₂ F ₅	106
50	1560	C ₂ H ₅	C ₂ H ₅	H	3-I	2-C1-4-C ₂ F ₅	103-105

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ym	Physical Properties (melting
							point: ℃
10	1561	C ₂ H ₅	C ₂ H ₅	Н	3-I	2-CH ₃ -4-C ₂ F ₅	115
	1562	t-C ₄ H ₉	H	Н	3-I	2-Br-4-C ₂ F ₅	185
	1563	i-C ₃ H ₇	Н	H	3-I	3-CH ₃ -4-C ₂ F ₅	240
15	1564	i-C ₃ H ₇	Н	Н	Н	4-0-(2-Pym)	246
	1565	C(CH ₃) ₂	Н	H	3-I	2-CH3-4-C2F5	193
		−CH 2 CH 3					
20	1566	C(CH ₃) ₂	Н	H	3-I	2-CH ₃ -4-0CF ₃	180
		-CH 2 CH 3					
	1567	C(CH ₃) ₂ CH ₂ CH ₃	H	Н	3-I		178-179
25	1568	C(CH ₃) ₂ CH ₂ CH ₃	H	Н	3-I	2-CH 3-4-0CHF 2	176-177
	1569	C(CH ₃) ₂ CH=CH ₂	H	Н	3-C1	2-CH3-4-C2F5	223-224
30	1570	C(CH ₃) ₂ CH≡C	, H	Н	3-C1	2-CH3-4-C2F5	92-93
		-(4-CH₃-Ph)					
	1571	C(CH ₃)₂CH≡C	H	Н	3-C1	2-CH3-4-C2F5	96-97
35		-(2,4-Cl ₂ -Ph)					
	1572	C(CH ₃)₂CH≡C	H	Н	3-C1	2-CH3-4-C2F5	88-89
		-(4-CH ₃ O-Ph)					·
40	1573	n-C3H7	C ₂ H ₅	Н	3-I	2-CH ₃ -4-C ₂ F ₅	93
	1574	n-C ₃ H ₇	C ₂ H ₅	H	3-I	2-CH ₃ -4-OCF ₃	109
	1575	n-C ₃ H ₇	C ₂ H ₅	Н	3-I	2-CH ₃ -4-OCHF ₂	102
45	1576	CH2(4-CF30-Ph)	H	Н	3-C1	2-CH3-4-C2F5	172
·	1577	CH2(4-CF30-Ph)	H	Н	6-C1	2-CH3-4-C2F5	193
	1578	CH ₂ (3-Cl-Ph)	СН₃	H	3-C1	2-CH3-4-C2F5	Paste
50	1579	CH2(2-F-Ph)	СН₃	H	3-C1	2-CH3-4-C2F5	115
:	1580	i-C3H7	H	H	3-I	2-Br-4-C ₂ F ₅	190

Table 1 (Cont'd)

5	No	R1	<u>R</u> 2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1581	n-C3H7	C2H5	H	3-F	2-CH ₃ -4-C ₂ F ₅	120
	1582	n-C 3H 7	C ₂ H ₅	Ħ	3-F	4-0CF ₃	115
	1583	n-C3H7	C ₂ H ₅	H	3-F	4-0CHF 2	85
15	1584	n-C3H7	C ₂ H ₅	H	3-F	2-C1-4-C ₂ F ₅	75
	1585	$C(CH_3)_2CH \equiv C$	H	H	3-C1	2-CH3-4-C2F5	102-103
20		-(4-CF ₃ -Ph)					
	1586	$C(CH_3)_2CH \equiv C$	H	H	3-C1	2-CH3-4-C2F5	115-117
		-(2,6-Cl ₂ -Ph)				:	
25	1587	$C(CH_3)_2CH \equiv C$	H	H	3-C1	2-CH3-4-C2F5	169
		-2-Pyi	:				
	1588	$C(CH_3)_2CH\equiv CH$	H	H	3-C1	2-CH 3-4-OCHF 2	191-192
30	1589	C(CH ₃) ₂ CH=CH ₂	H	Н	6-C1	2-CH ₃ -4-C ₂ F ₅	242
	1590	C(CH₃)₂CH≡C	H	Н	3-C1	2-CH3-4-C2F5	134-135
		-3-Pyi					
35	1591	i-C ₃ H ₇	H	Н	H	2-CH ₃ -4-(2,6	165
						-(CH ₃ O) ₂ -Ph)	
40	1592	i-C ₃ H ₇	H	H	H	2-CH ₃ -4-(3,5	150
40						-(CH3O)2-Ph)	
	1593	C ₂ H ₅	C ₂ H ₅	H	H	2-CH₃-4-(3,5	Paste
45					•	-(CH ₃ O) ₂ -Ph)	
	1594	i-C ₃ H ₇	Н	Н	3-C1	2-F-4-(0CF ₂ 0)-5	195
	1595	i-C ₃ H ₇	H	H	3-I	2-F-4-(0CF ₂ 0)-5	208
50	1596	t-C ₄ H ₈	H	Н	3-I	2-F-4-(0CF ₂ 0)-5	202

Table 1 (Cont'd)

5							
5	No	R1	R 2	Rз	Хn	Ym	Physical Properties
		·					(melting point: °C
10	1597	i-C ₃ H ₇	Н	Н	3-C1	2-CH ₃ -4-(OCHFCF ₂	211
						-0)-5	
15	1598	i-C ₃ H ₇	H	H	1-6	2-CH ₃ -4-(0CHFCF ₂	212
						-0)-5	
	1599	t-C4H9	Н	H	3-I	2-CH ₃ -4-(OCHFCF ₂	217
20						-0)-5	
	1600	i-C ₃ H ₇	H	Н	3-I	2-C1-4-(OCHFCF 2	210
					ı	-0)-5	
25	1601	i-C3H7	H	Н	3-I	2-C1-4-(OCF 2CHF	214
						-0)-5	
	1602	C(CH₃)₂C≡CH	H	Н	3-C1	2-CH3-4-0CF3	178-180
30	1603	C(CH₃)₂CHBr	H	H	3-C1	2-CH3-4-C2F5	130-131
		-CH ₂ Br					
35	1604	C(CH ₃) ₂ CH=CH	H	Н	3-C1	2-CH3-4-C2F5	90-93
		-Ph(E)					
	1605	C(CH ₃) ₂ CH ₂ Br	H	Н	3-I	2-CH3-4-0CF3	139-141
40	1606	(S)-C+H	Н	Н	3-I	2-CH3-4-C2F5	105-107
		-(CH ₃)-CH ₂ Br					
	1607	(R)-C+H	H	Н	3-I	2-CH3-4-C2F5	105-107
45		-(CH ₃)-CH ₂ Br			•		
	1608	i-C ₃ H ₇	Н	Н	3-I	3-C1-4-C ₂ F ₅	145
50	1609	t-C4H9	H	Н	3-I	3-C1-4-C ₂ F ₅	260
							

Table 1 (Cont'd)

5	No	Вī	R²	R3	Xn	Ym	Physical Properties (melting point: °C
10	1610	i-C ₃ H ₇	Н	Н	3-I	2-CH3-4-C2F5-5-CH3	210
	1611	t-C ₄ H ₉	H	Н	3-I	2-CH ₃ -4-C ₂ F ₅ -5-CH ₃	215
	1612	i-C ₃ H ₇	Н	Н	3-I	2,3-(CH ₃) ₂ -4-C ₂ F ₅	210
15	1613	t-C ₄ H ₉	H	Н	3-I	2,3-(CH ₃) ₂ -4-C ₂ F ₅	220
	1614	C2H5	C ₂ H ₅	Н	3-I	2-CH ₃ -4-(4-F-Ph)	130-133
20	1615	C2H5	C2H5	Н	3-I	2-CH ₃ -4-(4-Cl-Ph)	173-175
	1616	i-C₃H₁	H	н	H	2-CH ₃ -4-0-(2-Thz)	149
	1617	i-C3H7	H	Н	3-I	Mixture of 2-CH ₃ -4-	235
25						$(4-(2-CH_3-Thz))$ and	
						2-CH3-5-(4-(2-CH3-	
						Thz)) (1:1)	
30	1618	i-C3H7	H	Н	3-I	2-CH ₃ -4-0-(2-Pym)	239
	1619	C ₂ H ₅	C ₂ H ₅	Н	3-I	2-CH ₃ -4-(4-CF ₃ -Ph)	112-115
	1620	i-C ₃ H ₇	H	H	3-I	4-CF ₂ CF ₂ 0-5	239
35	1621	i-C ₃ H ₇	H	Н	3-C1	4-CF ₂ CF ₂ 0-5	243
	1622	i-C ₃ H ₇	H	H	3-I	2-C1-4-0CF ₂ 0-5	226
40	1623	i-C ₃ H ₇	H	H	3-C1	2-C1-4-0CF ₂ 0-5	223
40	1624	t-C4H9	H	H	3-I	2-C1-4-0CF ₂ 0-5	221
	1625	i-C3H7	H	Н	3-I	2-C1-4-0CF 2CF 2O	241
45	1626	i-C3H7	H	H	3-I	2-C1-3-0CF 2CF 20-4	219
	1627	C(CH ₃) ₂ CH ₂ Cl	Н	H	3-I	2-CH3-4-0CF3	160
	1628	C(CH ₃) ₂ C≡C	H	H	3-C1	2-CH3-4-C2F5	78-80
50		-3-Thi					
ļ							

Table 1 (Cont'd)

5	No	R1	R2	R3	Xn	Ym	Physical Properties (melting
10	1000	0/01/ \ 0 - 01/					point: °C
	1629	C(CH ₃) ₂ C≡CH	H	Н	3-I	2-CH3-4-C2F5	187-188
	1630	i-C ₃ H ₇	H	H	3-I	$2-CH_3-4-(3,5-(CH_30)_2$	199
15						-Ph)	
	1631	i-C ₃ H ₇	H	H	H	3-0CH ₂ 0-4	195
	1632	i-C3H7	Н	Н	Н	2-F-4-Cl	177
20	1633	C(CH ₃) ₂ C≡C	H	H	3-C1	2-CH3-4-C2F5	92-93
		-(4-CF ₃ 0-Ph)					
	1634	C(CH ₃) ₂ C≡CH	H	Н	3-I	2-CH ₃ -4-0CF ₃	188-189
25	1635	C(CH ₃) ₂ C≡CH	Н	Н	3-I	2-CH3-4-0CHF2	175-176
	1636	i-C ₃ H ₇	H	Н	3-I	4-N=(n-C ₃ F ₇)C-O-5	182
	1637	i-C ₃ H ₇	Н	Н	3-I	4-0-C(n-C ₃ F ₇)=N-5	250
30	1638	i-C ₃ H ₇	Н	Н	3-C1	4-0-C(n-C ₃ F ₇)=N-5	168
	1639	t-C₄H ₉	Н	Н	3-I	4-0-C(n-C ₃ F ₇)=N-5	248
	1640	i-C ₃ H ₇	H	Н	3-I	2,3-(CH ₃) ₂ -4-C ₂ F ₅	195
35	1641	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-OC(CF ₃)=N-5	229
	1642	i-C3H7	H	Н	3-C1	2-C1-3-0CF 2CF 2O-4	188
40	1643	i-C ₃ H ₇	H	H	3-C1	2-C1-4-0CF2CF20-5	203
**	1644	t-C₄H ₀	H	H	3-1	2-C1-3-0CF 2CF 2O-4	189
	1645	t-C ₄ H ₉	H	H	3-I	2-C1-4-0CF 2CF 20-5	234
45	1646	C(CH ₃) ₂ CH ₂ Cl	H	H	3-I	2-CH ₃ -4-C ₂ F ₅	168-169
	1647	C(CH3)2CH2Br	Н	Н	3-I	2-CH ₃ -4-C ₂ F ₅	167-168
	1648	C(CH ₃) ₂ C≡C	Н	Н	3-1	2-CH3-4-C2F5	90
50		-Naph					1

Table 1 (Cont'd)

-		,					
5	No	R1	R 2	R3	Xn	Ym	Physical Properties (melting point: °C
10	1649	$C(CH_3)_2C\equiv C$ -(5-Br-2-Pyi)	H	H	3-C1	2-CH3-4-C2F5	105-106
15	1650	$C(CH_3)_2C\equiv C$ -(2,4-F ₂ -Ph)	Н	Н	3-C1	2-CH ₃ -4-C ₂ F ₅	103-105
. 20	1651	(S)-C+H(CH₃) -CH₂F	Н	H	3-C1	2-CH ₃ -4-C ₂ F ₅	135
	1652	(S)-C*H -(CH ₃)-CH ₂ Br	Н	H	3-C1	2-CH ₃ -4-C ₂ F ₅	193-198
25	1653	i-C ₃ H ₇	Н	H	3-I	2-CH ₃ -4-C ₂ F ₅ -5 -Cl	210
30	1654	t-C₄H₀	Н	Н	3-I	2-CH3-4-C2F5-5 -Cl	200
35	1655	i-C ₃ H ₇	Н	Н	3-I	2-CH ₃ -4-C ₂ F ₅ -5 -CH ₃	190
	1656	t-C₄H₀	Н	Н	3-I	2-CH ₃ -4-C ₂ F ₅ -5 -CH ₃	195
40	1657	i-C3H7	Н	H	H	3-(2-CH ₃ -4-Thz)	211
	1658	i-C3H7	H	H	3-I	3-(2-CF ₃ -4-Thz)	122
45	1659	i-C ₃ H ₇	Н	Н	3-I	$3-(2-CH_3-4-0xa)$	102
	1660	i-C ₃ H ₇	H	Н	3-I	2-I-4-0CF ₂ 0-5	252
	1661	i-C ₃ H ₇	H	H	3-C1	2-CH ₃ -4-0CF ₂ 0-5	218
50	1662	t-C4H9	H	H	3-I	2-CH ₃ 0-4-C ₂ F ₅	135

Table 1 (Cont'd)

5	No	R1	R2	R 3	Xn	Ym	Physical Properties (melting
10							point: °C
	1663	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇ -5-F	235
	1664	t-C ₄ H ₉	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇ -5-F	230
15	1665	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇ -5-Cl	210
	1666	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-CF ₂ CF ₂ 0-5	198
	1667	i-C ₃ H ₇	H	H	3-I	2-CH3-4-i-C3F7	270
20	1668	t-C ₄ H ₉	H	H	3-1	2-CH ₃ -4-i-C ₃ F ₇	290
	1669	i-C ₃ H ₇	H	H	3-I	2-F-4-i-C ₃ F ₇	205
	1670	t-C ₄ H ₉	H	H	3-I	2-F-4-i-C ₃ F ₇	210
25	1671	i-C ₃ H ₇	H	H	3-I	2-SCH3-4-i-C3F7	205
	1672	t-C₄H _₽	H	H	3-I	2-SCH3-4-i-C3F7	205
30	1673	i-C ₃ H ₇	H	H	3-I	2,4-(CH ₃) ₂ -4-i-C ₃ F ₇	240
	1674	t-C ₄ H ₉	H	H	3-I	2,4-(CH ₃) ₂ -4-i-C ₃ F ₇	245
	1675	i-C ₃ H ₇	H	Н	3-I	4-(2-CH ₃ -4-Thz)	217
35	1676	i-C ₃ H ₇	H	Н	3-I	4-(2-CH ₃ -4-0xa)	212
	1677	i-C ₃ H ₇	H	H	3-I	4-(2-i-C ₃ H ₇ -4-Thz)	199
	1678	i-C ₃ H ₇	H	Н	3-NO2	4-(2-CH ₃ -4-Thz)	230
40	1679	i-C ₃ H ₇	H	Н	3-I	2-C1-3-OCF 2CHF0-4	188
	1680	i-C ₃ H ₇	H	Н	3-I	2-C1-3-OCHFCF20-4	191
	1681	i-C ₃ H ₇	H	H	3-I	Mixture of 2-Cl-3-	199
45				•		OCHFCF20-4-5-Cl and	
						2-C1-3-OCHFCF20-4-6-	
50		•		:		Cl (1:1)	

Table 1 (Cont'd)

	No	Rı	R2	R3	Xn	Ym	Physical Properties (melting point: °C
10	1682	i-C ₃ H ₇	H	H	3-I	2-C1-3-N=C(CF ₃)-0-4	265
	1683	t-C4H9	Н	Н	3-I	2-C1-3-N=C(CF ₃)-0-4	259
	1684	i-C₃H7	Н	Н	3-I	2-Br-4-0CF ₂ CHF0-5	185
15	1685	i-C₃H₁	Н	Н	3-I	Mixture of 2,3-Br ₂ -4-	250
						OCF ₂ CHFO-5; 2,5-Br ₂ -3-	
20						OCHFCF 20-4; and 2,6-	
						Br ₂ -3-0CF ₂ CHF0-4(1:1:1)	
	1686	i-C₃H₁	H	H	3-I	Mixture of 2,3-Br ₂ -4-	228
25						OCHFCF20-5; 2,5-Br2-3-	
						OCF 2CHF0-4; and 2,6-	
						Br ₂ -3-0CF ₂ CHF0-4(1:1:1)	
30	1689	i-C ₃ H ₇	H	Н	3-I	2,3-(CH ₃) ₂ -4-i-C ₃ F ₇	270
	1690	t-C4H9	H	H	3-I	2,3-(CH ₃) ₂ -4-i-C ₃ F ₇	280
	1691	i−C₃H7	H	H	3-I	2-i-C ₃ H ₇ -4-i-C ₃ F ₇	240
35	1692	t-C ₄ H ₉	Н	H	3-I	2-i-C ₃ H ₇ -4-i-C ₃ F ₇	245
	1693	i-C ₃ H ₇	Н	H	3-I	2-0C ₂ H ₅ -4-i-C ₃ F ₇	195
40	1694	t-C₄H9	H	H	3-I	2-0C ₂ H ₅ -4-i-C ₃ F ₇	210
40	1695	i-C3H7	Н	H	3-I	3-F-4-i-C ₃ F ₇	265
	1696	t-C₄H ₉	H	H	3-I	3-F-4-i-C ₃ F ₇	285
45	1697	i-C ₃ H ₇	H	H	3-I	3-Cl-4-i-C ₃ F ₇	295
	1698	i-C3H7	Н	Н	3-I	2-Br-4-i-C ₃ F ₇ -5-CH ₃	240
	1699	i-C ₃ H ₇	Н	Н	3-I	2-Br-4-i-C ₃ F ₇	240
50	1700	i-C ₃ H ₇	H	Н	3-I	2-SCH ₃ -4-C ₂ F ₅	200
	1703	i-C ₃ H ₇	Н	H	3-I	4-(2-c-C ₃ H ₅ -4-Thz)	198

Table 1 (Cont'd)

5	No	R1	R 2	R3	Xn	Ym	Physical Properties (melting point: °C
,,	1714	i-C ₃ H ₇	H	Н	3-I	2-C ₂ H ₅ -4-i-C ₃ F ₇	220
	1715	i-C3H7	Н	Н	3-I	2-0CH ₃ -4-i-C ₃ F ₇	190
15	1716	i-C ₃ H ₇	Н	H	3-I	2,6(CH ₃) ₂ -4-i-C ₃ F ₇	275
	1717	i-C3H7	Н	Н	3-I	2,6-(CH ₃) ₂ -4-C ₂ F ₅	250
	1722	i-C ₃ H ₇	Н	Н	3-I	2-Cl-4-i-C ₃ F ₇	220
20	1723	t-C ₄ H ₉	Н	H	3-I	2-Cl-4-i-C ₃ F ₇	210
	1726	i-C ₃ H ₇	H	H	3-I	2-(CH ₂) ₄ -3-4-i-C ₃ F ₇	260
25	1727	t-C ₄ H ₉	H	H	3-I	2-(CH ₂) ₄ -3-4-i-C ₃ F ₇	272
	1732	i-C3H7	Н	H	3-I	2-C1-3-0CF ₂ CF ₂ 0-4	245
	1733	i-C ₃ H ₇	H	H	3-I	2-C1-3-OCHFCF20-4	190
30	1737	i-C ₃ H ₇	H	H	3-I	4-C(CH ₃)=NOCH ₃	190
	1742	i-C ₃ H ₇	H	H	3-I	2-0CF ₂ 0-3	190
<i>35</i>	1743	i-C ₃ H ₇	H	H	3-I	2-0CF ₂ 0-3-6-C1	213
	1744	i-C3H7	H	Н	3-I	2-0CF ₂ 0-3-4-C1	202
	1745	i-C3H7	H	H	3-I	2-0CF ₂ 0-3-4,6-Cl ₂	228
40	1746	i-C ₃ H ₇	H	H	3-I	2-0CF ₂ 0-3-4-i-C ₃ F ₇	175
	1747	t-C ₄ H ₉	H	H	3-I	2-0CF ₂ 0-3-4-C1	235
45	1748	t-C ₄ H ₉	H	H	3-I	2-0CF ₂ 0-3-4,6-Cl ₂	243
45	1749	i-C ₃ H ₇	H	H	3-I	4-C(CH ₃)=NOCH ₂ -Ph	205
	1750	i-C3H7	H	H	3-I	4-C(CH ₃)=NOCH ₂	Decomp.
50						−CH=CH 2	

Table 1 (Cont'd)

5 .		Т					5 1
	No	R1	R ²	Rз	Xn	Ym	Physical Properties
				"	****		(melting
10							point: °C
	1751	СН₃	СНз	H	H	2-CH3-4-C1	149
	1752	C2H5	C ₂ H ₅	H	H	2-CH3-4-C1	172
15	1753	n-C3H7	n-C3H7	H	H	2-CH ₃ -4-Cl	126
	1762	i-C3H7	H	H	3-I	3-C(i-C ₃ F ₇)=NN	Paste
						-(i-C ₃ F ₇)-4	
20	1763	i-C ₃ H ₇	Н	H	3-I	4-i-C ₃ H ₇ -2-N=CH-S-3	200
	1764	i-C ₃ H ₇	H	H	3-I	3-S-C(i-C ₃ H ₇)=N-4	218
<i>25</i>	1765	i-C3H7	Н	H	3-I	4-(2-CF ₃ -4-Thz)	105
	1766	i-C3H7	H	H	3-I	3-SCH3-4-i-C3F7	160
	1767	i-C3H7	Н	H	3-I	2-Ph-4-i-C ₃ F ₇	240
30	1768	i-C ₃ H ₇	Н	H	3-I	2-0Ph-4-i-C ₃ F ₇	180
	1769	i-C ₃ H ₇	Н	H	3-I	2-0CH ₃ -4-i-C ₃ F ₇	265
35	1770	(CH ₂) ₂ -3-Pyi	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	Amorphous
	1771	(CH ₂) ₂ -3-Pyi	Н	H	6-I	2-CH ₃ -4-i-C ₃ F ₇	Amorphous
	1772	(CH ₂) ₂ -3-Pyi	Н.	H	3-I	2-CH ₃ -4-OCF ₃	169-173
40	1773	CH(CH₃)-2-Pyi	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	Amorphous
	1774	CH(CH₃)-2-Pyi	H	H	6-I	2-CH3-4-i-C3F7	Amorphous
45	1775	CH(CH₃)-2-Pyi	H	H	3-I	2-CH ₃ -4-OCF ₃	158-161
70	1776	CH(CH₃)-2-Pyi	Н	H	6-I	2-CH ₃ -4-OCF ₃	213-216
	1777	CH(CH₃)-2-Pyi	Н	H	3-I	2-CH ₃ -4-C ₂ F ₅	149-152
50	1778	CH(CH₃)-2-Pyi	Н	H	6-I	2-CH ₃ -4-C ₂ F ₅	194-196
ĺ							

Table 1 (Cont'd)

5	
10	
15	
20	

No	R1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
1780	N(Ph)COCF₃	Н	Н	3-I	2-CH3-4-C2F5	239-241
1799	CH(CH3)-2-Fur	Н	H	3-1	2-CH3-4-i-C3F7	191
1800	CH(CH₃)-2-Thi	Н	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	159
,1801	i-C3H7	Н	Н	3-CF 3	2-CH3-4-C2F5	210-212
1802	i-C ₃ H ₇	H	H	3-C1-6-	2-CH ₃ -4-C ₂ F ₅	236-237
				CF 3 S		
1803	i-C3H7	Н	Н	3-CF 3 SO	2-CH3-4-C2F5	186-187
1804	i-C3H7	H	Н	6-CF ₃ S0	2-CH3-4-C2F5	206-208
1805	i-C3H7	H	Н	3-CF 3 SO	2-CH3-4-i-C3F7	211-213
1815	i-C3H7	H	Н	3-I	2-CH3-4-s-C4F9	190
1816	i-C3H7	Н	Н	3-I	2-0H-4-i-C ₃ F ₇	155
1824	i-C3H7	Н	Н	3-1	2-N=C(CF ₃)0-3	132
					-4-i-C ₃ F ₇	
1825	i-C3H7	H	H	3-I	2-N=C(CF ₃)0-3	145
1826	t-C₄H9	H	H	3-1	2-N=C(CF ₃)0-3	110
					-4-i-C ₃ F ₇	
1827	t-C ₄ H ₉	H	Н	3-I	2-N=C(CF ₃)0-3	120
1829	(CH ₂) ₂ NH-CO ₂ C ₂ H ₅	H	Н	3-I	2-CH3-4-i-C3F7	155
1830	(CH ₂) ₂ NHCO ₂ CH ₂ Ph	H	H	3-I	2-CH ₃ -4-C ₂ F ₅	155
1831	(CH ₂) ₂ CH=CF ₂	H	Н	3-I	2-CH3-4-C2F5	180

Table 1 (Cont'd)

5	No	R1	R2	Ra	Xn	Ym	Physical Properties (melting point: °C
10	1838	i-C ₃ H ₇	Н	Н	Н	2-CH ₃ -4-OCF ₂ CF ₃	POZZO. C
	1839	i-C₃H7	H	Н	3-NO ₂	2-CH ₃ -4-OCF ₂ CF ₃	
	1840	i-C₃H7	H	Н	3-F	2-CH ₃ -4-OCF ₂ CF ₃	
15	1841	i-C3H7	H	Н	3-C1	2-CH3-4-OCF2CF3	
	1842	i-C₃H₁	Н	Н	3-Br	2-CH ₃ -4-OCF ₂ CF ₃	
20	1843	i-C ₃ H ₇	H	Н	3-C1-4-F	2-CH3-4-OCF2CF3	
20	1844	i-C ₃ H ₇	Н	Н	3,4-Cl ₂	2-CH3-4-OCF2CF3	
	1845	i-C3H7	H	Н	3-I	4-0CF 2CF 3	
25	1846	i-C3H7	Н	Н	3-I	2-C1-4-0CF 2CF 3	400
	1847	i-C3H7	H	Н	3-I	2-CH ₃ -4-OCF ₂ CF ₃	
	1848	t-C₄H9	H	Н	3-I	2-CH ₃ -4-OCF ₂ CF ₃	
30	1849	C2H5	C ₂ H ₅	Н	3-I	2-CH ₃ -4-OCF ₂ CF ₃	:
	1850	i-C ₃ H ₇	H	H	3-I	2-C ₂ H ₅ -4-0CF ₂ CF ₃	
	1851	i-C ₃ H ₇	H	Н	Н	2-CH ₃ -4-0-n-C ₃ F ₇	
35	1852	i-C3H7	Н	Н	3-NO ₂	2-CH ₃ -4-0-n-C ₃ F ₇	
•	1853	i-C ₃ H ₇	Н	Н	3-F	2-CH3-4-0-n-C3F7	
40	1854	i-C3H7	H	Н	3-C1	2-CH ₃ -4-0-n-C ₃ F ₇	·
40	1855	i-C ₃ H ₇	H	Н	3-Br	2-CH ₃ -4-0-n-C ₃ F ₇	
	. 1856	t-C ₄ H ₉	H	Н	3-C1-4-F	2-CH3-4-0-n-C3F7	
45	1857	C2H5	C ₂ H ₅	H	3,4-Cl ₂	2-CH3-4-0-n-C3F7	<u> </u>
	1858	i-C3H7	H	H	3-1	4-0-n-C ₃ F ₇	
	1859	i-C3H7	H	H	3-I	2-C1-4-0-n-C3F7	
50	1860	i-C3H7	H	H	3-I	2-CH ₃ -4-0-n-C ₃ F ₇	
	L						

Table 1 (Cont'd)

5	No	R 1	R ²	R3	Xn	Ym	Physical Properties (melting
10	1861	t-C4H9	Н	Н	3-1	2-CH ₃ -4-0-n-C ₃ F ₇	point: °C
	1862	C ₂ H ₅	C ₂ H ₅	H	3-I	2-CH ₃ -4-0-n-C ₃ F ₇	
	1863	i-C ₃ H ₇	H	Н	3-I	2-C ₂ H ₅ -4-OCF ₂ CF ₃	
15	1864	1-C3H7	H	Н	3-I	2-CH ₃ -4-C≡C-t-C ₄ F ₉	
	1865	1-C3H7	н	Н	3-1	2-CH ₃ -4-C≡C-CF ₃	
20	1866	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-C≡C-i-C ₃ F ₇	
	1867	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-CF=CF ₂	
	1868	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-CF=CFCF ₃	
25	1869	i-C₃H7	H	H	3-I	2-CH ₃ -4-C(CF ₃)=CF ₂	
	1870	i-C₃H7	H	H	3-I	2-CH ₃ -4-COCH ₃	
	1871	i-C3H7	H	H	3-I	2-CH ₃ -4-COCF ₃	195
30	1872	i-C ₃ H ₇	H	H	3-I	2-CH ₃ -4-COC ₂ F ₅	
	1873	i-C ₃ H ₇	H	Н	3-I	2-CH ₃ -4-COCF(CH ₃) ₂	
	1874	i-C ₃ H ₇	Н	Н	3-I	2-CH3-4-COOCH3	217
35	1875	i-C₃H₁	H	H	3-I	2-CH3-4-COOC2H5	
	1876	i-C3H7	Н	Н	3-I	2-CH ₃ -4-C(CH ₃)=NOCH ₃	218
	1877	i-C ₃ H ₇	Н	Н	3-I	2-CH ₃ -4-C(CH ₃)=NOC ₂ H ₅	
40	1878	i-C3H7	Н	Н	3-I	2-CH ₃ -4-C(CH ₃)=NO	
	}					-CH2CH=CH2	
45	1879	i-C₃H₁	Н	H	3-I	2-CH ₃ -4-C(CH ₃)=NO	
43						-CH₂C≡CH	
	1880	i-C3H7	Н	Н	3-1	2-CH ₃ -4-C(CH ₃)=NOCH ₂ -Ph	
50	1881	i-C3H7	Н	Н	3-I	2-CH3-4-CH2OH	

Table 1 (Cont'd)

5	No	R1	R2	Rз	Хn	Ym	Physical Properties (melting point: °C
10	1882	i-C ₃ H ₇	Н	Н	3-I	4-CH(OH)CH ₃	
	1883	i-C3H7	Н	Н	3-I	2-CH ₃ -4-CH(OH)CH ₃	
	1884	i-C3H7	Н	Н	3-I	2-CH ₃ -4-CH ₂ ON=C(CH ₃) ₂	
15	1885	i-C3H7	Н	H	3-I	2-CH ₃ -4-CH ₂ ON=C(Ph)	
						-i-C ₃ H ₇	
20	1886	i-C3H7	н	H	3-I	2-0CH ₂ 0-3-4-i-C ₃ F ₇	
,	1887	i-C ₃ H ₇	H	H	3-I	2-0CH ₂ CH ₂ 0-3-4-i-C ₃ F ₇	
	1888	i-C3H7	H	H	3-I	2-0CF ₂ CF ₂ 0-3-4-i-C ₃ F ₇	
25	1889	i-C3H7	Н	H	3-I	2-0CF ₂ CHF0-3-4-i-C ₃ F ₇	
	1890	i-C3H7	Н	H	3-I	2-0CHFCF20-3-4-i-C3F7	
	1891	i-C3H7	Н	H	3-I	2-SCH ₂ S-3-4-i-C ₃ F ₇	
30	1892	i-C3H7	Н	H	3-I	2-SCF ₂ S-3-4-i-C ₃ F ₇	
	1893	i-C3H7	Н	H	3-I	2-SCH2CH2S-3-4-i-C3F7	
	1894	i-C3H7	Н	H	3-I	2-SCF2CF2S-3-4-i-C3F7	
35	1895	i-C ₃ H ₇	H	H	3-I	2-CH ₂ OCH ₂ -3-4-i-C ₃ F ₇	
	1896	i-C3H7	H	H	3-I	2-CH ₂ SCH ₂ -3-4-i-C ₃ F ₇	
40	1897	i-C3H7	H	H	3-I	2-CF ₂ OCF ₂ -3-4-i-C ₃ F ₇	
***	1898	i-C ₃ H ₇	H	H	3-I	2-CF ₂ SCF ₂ -3-4-i-C ₃ F ₇	
	1899	i-C₃H₁	H	H	3-Br	2-CH ₃ -4-i-C ₃ F ₇	
45	1900	i-C ₃ H ₇	H	H	3-Br	2-CH ₃ -4-i-C ₃ F ₇	
					-4-Cl		
	1901	i-C₃H7	H	H	3-I-4-F	2-CH3-4-i-C3F7	
50	1902	i-C3H7	H	H	3-I-4-Cl	2-CH ₃ -4-i-C ₃ F ₇	

Table 1 (Cont'd)

5							Physical
	No	R 1	R2	R3	Χn	Ym	Properties
							(melting point: °C
10	1903	i-C ₃ H ₇	Н	Н	3-I-4-CF ₃	2-CH ₃ -4-i-C ₃ F ₇	point. C
	1904	i-C ₃ H ₇	н	Н	3-I-4-0CH ₃	2-CH ₃ -4-i-C ₃ F ₇	
	1905	i-C ₃ H ₇	H	H	3-I-4-Br	2-CH ₃ -4-i-C ₃ F ₇	
15							
	1906	i-C3H7	H	H	3-C1-4-CF ₃	2-CH3-4-i-C3F7	
•	1907	i-C ₃ H ₇	H	H	3-CF ₃ -4-Cl	2-CH ₃ -4-i-C ₃ F ₇	
20	1908	i-C ₃ H ₇	H	H	3-CF 3-4-F	2-CH ₃ -4-i-C ₃ F ₇	
	1919	i-C3H7	H	Н	3-CF ₃ -4-0CH ₃	2-CH ₃ -4-i-C ₃ F ₇	
	1910	i-C3H7	H	Н	3-N=CH-CH=CH-4	2-CH ₃ -4-i-C ₃ F ₇	
25	1911	i-C3H7	H	н	3-0CH ₂ 0-4	2-CH ₃ -4-i-C ₃ F ₇	·
	1912	i-C3H7	Н	н	3-0CH ₂ 0-4	2-CH3-4-C2F6	
	1913	i-C ₃ H ₇	H	H	3-0CH ₂ 0-4	2-CH ₃ -4-0CF ₃	
30	1914	i-C ₃ H ₇	Н	Н	3-0CF ₂ 0-4	2-CH ₃ -4-i-C ₃ F ₇	
	1915	i-C3H7	H	Н (3-0CF 20-4	2-CH3-4-C2F5	
	1916	i-C ₃ H ₇	H	Н	3-0CF ₂ 0-4	2-CH3-4-0CF3	
35	1917	i-C3H7	H	Н	3-0CH ₂ CH ₂ 0-4	2-CH ₃ -4-i-C ₃ F ₇	
	1918	i-C3H7	H	H	3-0CF 2CF 20-4	2-CH3-4-i-C3F7	
40	1919	i-C3H7	Н	Н	3-0CHFCF 20-4	2-CH3-4-i-C3F7	
	1920	i-C ₃ H ₇	Н	H	3-0CF 2CHF0-4	2-CH ₃ -4-i-C ₃ F ₇	
	1921	i-C ₃ H ₇	Н	H	3-0CH ₂ CH ₂ -4	2-CH3-4-i-C3F7	
45	1922	i-C ₃ H ₇	Н	Н	3-CH2CH2O-4	2-CH ₃ -4-i-C ₃ F ₇	
	1923	i-C ₃ H ₇	Н	Н	3-0CF 2CF 2-4	2-CH3-4-i-C3F7	
	1924	i-C ₃ H ₇	Н	Н	3-CF 2CF 20-4	2-CH ₃ -4-i-C ₃ F ₇	
50	1925	i-C ₃ H ₇	Н	Н	3-SOCH3	2-CH ₃ -4-i-C ₃ F ₇	

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ym	Physical Properties (melting point: °C
10	1926	i-C ₃ H ₇	Н	Н	3-S02CH3	2-CH ₃ -4-i-C ₃ F ₇	
	1927	i-C₃H7	H	H	3-CF 3 S	2-CH3-4-i-C3F7	222-223
15	1928	i-C ₃ H ₇	H	Н	6-CF 3 S	2-CH ₃ -4-i-C ₃ F ₇	219-221
	1929	t-C ₄ H ₉	H	Н	3-CF 3 S	2-CH3-4-i-C3F7	231
	1930	t-C ₄ H ₉	H	Н	6-CF 3 S	2-CH3-4-i-C3F7	245-247
20	1931	t-C4H9	H	H	3-CF 3 SO 2	2-CH3-4-i-C3F7	
	1932	t-C ₄ H ₉	Н	H	3-CF 3 SO 2	2-CH ₃ -4-C ₂ F ₅	
	1933	t-C ₄ H ₉	H	H	3-CF 3 SO 2	2-CH ₃ -4-0CF ₃	
25	1934	C2H5	C ₂ H ₅	Н	3-CF 3 SO 2	2-CH3-4-i-C3F7	
	1935	C2H5	C ₂ H ₅	Н	3-CONHCH₃	2-CH3-4-i-C3F7	
30	1936	C2H5	C ₂ H ₅	Н	3-CON(CH ₃) ₂	2-CH ₃ -4-i-C ₃ F ₇	
	1937	C2H5	C ₂ H ₅	H	3-COCH3	2-CH3-4-i-C3F7	
	1938	C2H5	C ₂ H ₅	Н	3-COC2H5	2-CH3-4-i-C3F7	
35	1939	C2H5	C ₂ H ₅	Н	3-C(CH ₃)=NOCH ₃	2-CH ₃ -4-i-C ₃ F ₇	
	1940	C2H5	C ₂ H ₅	Н	3-C(CH ₃)=NO	2-CH3-4-i-C3F7	
					-C ₂ H ₅		
40	1941	i-C3H7	Н	Н	3-C≡CH	2-CH3-4-C2F5	
	1942	i-C₃H₁	Н	H	3-С≡СН	2-CH ₃ -4-i-C ₃ F ₇	
45	1943	i-C₃H₁	Н	Н	3-C≡C-t-C ₄ H ₉	2-CH ₃ -4-C ₂ F ₅	195-202
**	1944	i-C₃H7	Н	H	3-C≡C-t-C ₄ H ₉	2-CH ₃ -4-i-C ₃ F ₇	-
	1945	i-C ₃ H ₇	Н	Н	3-C≡C-Ph	2-CH3-4-C2F5	179-183
50	1946	i-C ₃ H ₇	Н	Н	3-C≡C-Ph	2-CH3-4-i-C3F7	

Table 1 (Cont'd)

5							
10	No	R1	R2	R3	Xn	Ym	Physical Properties (melting point: °C
	1947	i-C ₃ H ₇	Н	H	3-C≡C	2-CH ₃ -4-C ₂ F ₅	
15	1948	i-C ₃ H ₇	H	Н	-CF 3 3-C≡C	2-CH ₃ -4-i-C ₃ F ₇	
20	1949	i-CaH7	H	Н	-CF ₃	2-CH3-4-C2F5	
20	1950	t-C ₄ H ₉	Н	Н	3-C ₂ F ₅	2-CH3-4-C2F5	
	1951	C2H5	C ₂ H ₅	Н	3-C ₂ F ₅	2-CH ₃ -4-C ₂ F ₅	
25	1952	i-C3H7	H	Н	3-C ₂ F ₅	2-CH ₃ -4-i-C ₃ F ₇	
	1953	t-C4H9	Н	Н	3-C ₂ F ₅	2-CH3-4-i-C3F7	
	1954	C2H5	C2H5	Н	3-C ₂ F ₅	2-CH ₃ -4-i-C ₃ F ₇	
30	1955	i-C ₃ H ₇	SN	Н	3-I	2-CH3-4-i-C3F7	
			$-(n-C_4H_9)_2$				
35	1956	i-C ₃ H ₇	SO ₂ CH ₃	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	·
	1957	i-C₃H7	CN	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	·
	1958	i-C₃H₁	СООСН3	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
40	1959	i-C₃H₁	C00C2H5	Н	3-I	2-CH3-4-i-C3F7	
	1960	i-C ₃ H ₇	COCH ₃	Н	3-1	2-CH3-4-i-C3F7	
45	1961	i-C ₃ H ₇	COC ₂ H ₅	Н	3-1	2-CH ₃ -4-i-C ₃ F ₇	
	1962	i-C ₃ H ₇	CO-Ph	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	1963	i-C3H7	NHCOCH 3	Н	3-1	2-CH3-4-i-C3F7	
50]

Table 1 (Cont'd)

5		·					,
	No	R1	R ²	Rз	Xn	Ym	Physical Properties (melting
							point: °C
10	1964	C2H5	C ₂ H ₅	SN(n	3-I	2-CH ₃ -4-i-C ₃ F ₇	
				-C4H9)2			
15	1965	C2H5	C ₂ H ₅	SO ₂ CH ₃	3 - I	2-CH3-4-i-C3F7	
	1966	C2H5	C ₂ H ₅	CN	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	1967	C2H5	C ₂ H ₅	COOCH ₃	3-I	2-CH3-4-i-C3F7	
20	1968	C ₂ H ₅	C ₂ H ₅	COOC 2H 5	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	1969	C ₂ H ₅	C ₂ H ₅	COCH3	3-I	2-CH3-4-i-C3F7	
	1970	C2H5	C ₂ H ₅	COC 2H 5	3-I	2-CH ₃ -4-i-C ₃ F ₇	Amorphous
25	1971	C ₂ H ₅	C ₂ H ₅	COPh	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	1972	C2H5.	C ₂ H ₅	NHCOCH 3	3-I	2-CH ₃ -4-i-C ₃ F ₇	
30	1973	(CH ₂) ₂ COO	H	H	3-I	2-CH3-4-i-C3F7	
		-СН з					:
	1974	(CH ₂) ₂ COO	H	Н	3-I	2-CH ₃ -4-C ₂ F ₅	
35		−СН з					
	1975	(CH ₂) ₂ COO	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	133.2
		-C ₂ H ₅					
40	1976	(CH ₂) ₂ COO	H	Н	3-I	2-CH3-4-C2F5	
	ľ	-C2H5		:			
45	1977	(CH ₂) ₂ COO	H	H	6-I	2-CH3-4-C2F6	163.5
		-C ₂ H ₅					
	1978	CH(CH ₃)CH ₂	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
50		-C00CH3				·	

Table 1 (Cont'd)

5	No	R1	R2	B 3	Xn	Ym	Physical Properties (melting point: °C
10	1979	CH(CH ₃)CH ₂ COOC ₂ H ₅	Н	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	point. C
15	1980	CH(CH3)CH2COO-i- C3H7	Н	H	3-1	2-CH ₃ -4-i-C ₃ F ₇	
	1981	(CH ₂) ₂ CONHCH ₃	Н	H	3-I	2-CH3-4-i-C3F7	
	1982	(CH ₂) ₂ CONHC ₂ H ₅	H	H	3-1 3-I	2-CH ₃ -4-1-C ₃ F ₇	
20	1983	CH(CH ₃)CH ₂ CONHCH ₃	н	Н	3-1 3-I	2-CH ₃ -4-1-C ₃ F ₇	
	1984						
		CH(CH ₃)CH ₂ CONHC ₂ H ₅	Н	H 	3-I	2-CH ₃ -4-i-C ₃ F ₇	
25	1985	CH(CH ₃)CH ₂ CONH-i-	H	H	3-I	2-CH3-4-i-C3F7	
		C3H7					
	1986	CH(CH3)CH2CON	Н	H	3-I	2-CH3-4-i-C3F7	
30		-(CH ₃) ₂					
	1987	CH(CH ₃)CH ₂ CON	H	H	3-I	2-CH3-4-i-C3F7	
		-(C ₂ H ₅) ₂					
35	1988	(CH ₂) ₂ NHCOOCH ₃	H	H	3-I	2-CH3-4-C2F5	
	1989	(CH ₂) ₂ NHCOOCH ₃	H	Н	3-I	2-CH3-4-i-C3F7	
	1990	(CH2)2NHCOOC2H5	H	H	3-I	2-CH3-4-C2F5	145
40	1991	(CH ₂) ₂ NHCOOC ₂ H ₅	Н	Н	3-I	2-CH ₃ -4-OCF ₃	210
	1992	CH(CH3)CH2NHCOOCH3	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	
45	1993	CH(CH₃)CH₂NHCOO	Н	Н	3-I	2-CH3-4-i-C3F7	
		-C ₂ H ₅					
	1994	(CH ₂) ₂ P(CH ₃) ₂	H	H	3-I	2-CH3-4-i-C3F7	
50	1995	CH(CH ₃)P(C ₂ H ₅) ₂	Н	Н	3-I	2-CH3-4-i-C3F7	

Table 1 (Cont'd)

5	No	R1	R2	Ra	Xn	Ym	Physical Properties
				! !			(melting point: °C
10	1996	(CH ₂) ₂ P(Ph) ₂	H	Н	3-I	2-CH3-4-i-C3F7	point. C
	1997	CH(CH ₃)CH ₂ P(CH ₃) ₂	H	H	3-I	2-CH3-4-i-C3F7	
15	1998	CH(CH ₃)CH ₂ P(C ₂ H ₅) ₂	Н	H	3-I	2-CH3-4-i-C3F7	
	1999	CH(CH ₃)CH ₂ P(Ph) ₂	Н	H	3-I	2-CH3-4-i-C3F7	
	2000	CH(CH ₃)(CH ₂) ₂ P	H	H	3-I	2-CH3-4-i-C3F7	
20		-(CH ₃) ₂					!
	2001	CH(CH ₃)(CH ₂) ₃ P	Н	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
		-(CH ₃) ₂					
25	2002	(CH ₂) ₂ PO(CH ₃) ₂	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	2003	$(CH_2)_2PO(OC_2H_5)_2$	Н	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	Amorphous
30	2004	CH(CH ₃)CH ₂ PO(OCH ₃) ₂	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	2005	(CH ₂) ₂ OPO(OCH ₃) ₂	Н	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
	2006	CH(CH ₃)CH ₂ PS(OCH ₃) ₂	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
35	2007	CH(CH₃)CH₂PS	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
		-(OC ₂ H ₅) ₂					
	2008	(CH ₂) ₂ OPO(OC ₂ H ₅) ₂	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	
40	2009	CH(CH ₃)CH ₂ OPO	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	
		-(OCH ₃) ₂					
45	2010	CH(CH ₃)CH ₂ OPO	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
		-(OC2H5)2					
	2011	(CH ₂) ₂ OPS(OCH ₃) ₂	H	Н	3-I	2-CH ₃ -4-i-C ₃ F ₇	
50	2012	(CH ₂) ₂ OPS(OC ₂ H ₅) ₂	Н	н	3-I	2-CH3-4-i-C3F7	

Table 1 (Cont'd)

5	No	R1	R2	Rз	Xn	Ym	Physical
	NO	r.	n-	n s	VII	1111	Properties (melting
10							point: °C
	2013	CH(CH3)CH2OPS	Н	H	3-I	2-CH3-4-i-C3F7	
		(OCH ₃) ₂		! 			
15	2014	CH(CH3)CH2OPS	H	H	3-I	2-CH ₃ -4-i-C ₃ F ₇	
		-(OC ₂ H ₅) ₂					
	2015	CH(CH ₃)-2-Pyi-N	H	H	3-I	2-CH3-4-C2F5	198-205
20		-0xide					
	2016	CH(CH ₃)-2-Pyi-N	Н	Н	3-1	2-CH ₃ -4-i-C ₃ F ₇	208-210
25		-0xide					
	2017	i-C3H7	Н	Н	3-I	2-CH ₃ -4-C(CF ₃)	
						=NOCH 3	
30	2018	i-C3H7	H	H	3-I	2-CH ₃ -4-C(CF ₃)	
				!		=NOCH 2 Ph	
35	2019	i-C3H7	H	Н	3-I	2-NCHCHCHCH-3	180
	·					-4-i-C ₃ F ₇	
	2020	i-C3H7	Н	Н	3-I	2-n-C3H7-4-i	225
40						-C ₃ F ₇	
	2021	i-C3H7	Н	Н	3-I	2-0-(2-Pyi)-4	158.3-159.8
45						-i-C ₃ F ₇	

[0083] The abbreviations in Table 1 stand for the following substituents:

Ph: phenyl group,

50

55

c-: alicyclic hydrocarbon group,

Pyi: pyridyl group, Pym: pyrimidinyl group, Fur: furyl group,

TetFur: tetrahydrofuryl group,

Thi: thienyl group, Thz: thiazolyl group,

Naph: naphthyl group, Oxa: oxazolyl group,

C*: asymmetric carbon atom

Table 2

10

5

15

20

 R^2 Z² R¹ R³ Z^1 No Xn Ym **Physical Properties** (melting point: °C 2-CH₃-4-CF₂CF₃ S 0 S-1 i-C₃H₇ Н Н 3-CI 162-164 S 0 S-2 t-C₄H₉ Н Н 3-CI 2-CH₃-4-CF₂CF₃ 141-143 S-3 Н 3-CI 2-CH₃-4-CF₂CF₃ S 0 c-C₃H₅ Н 138-139 S **S-4** Н 3-CI 2-CH₃-4-CF₂CF₃ 0 C_2H_5 C_2H_5 184-186 i-C₃H₇ S S-5 Н н 2-CH₃-4-CI 0 168-170 Н 2-CH₃-4-Cl 0 **S-6** i-C₃H₇ Н Н н S S-7 2-CH₃-4-i-C₃H₇ 0 S i-C₃H₇ Н Н 3-1 **S-8** Н 2-CH₃-4-i-C₃H₇ S S i-C₃H₇ Н S S-9 i-C₃H₇ Н Н 3-1 2-CH₃-4-i-C₃H₇ S

[0084] The ¹H-NMR data of the compounds obtained as paste (physical properties) are given in Table 3 below.

Table 3

30

25

No.	¹ H-NMR[CDCl ₃ /TMS, δ values (ppm)]
1122	1.2-1.4(m.6H), 2.4-2.5(m.3H), 3.1-3.9(m.7H), 6.6-7.9(m.6H)
1218	1.3(d.3H), 2.3(s.3H), 2.9-3.2(m.2H), 4.4(m.1H), 6.2(d.1H), 7.1-7.5(m.3H), 7.8(d.1H), 8.0(d.1H), 8.4(d.1H),8.5(s.1H).

[0085] Agricultural and horticultural insecticides containing the phthalic acid diamide derivative of the general formula (I) of the present invention as an active ingredient are suitable for controlling various insect pests such as agricultural insect pests, forest insect pests, horticultural insect pests, stored grain insect pests, sanitary insect pests, nematodes, etc., which are injurious to paddy rice, fruit trees, vegetables, other crops, flowers and ornamental plants, etc. They have a marked insecticidal effect, for example, on LEPIDOPTERA including summer fruit tortrix (Adoxophyes orana fasciata), smaller tea tortrix (Adoxophyes sp.), Manchurian fruit moth (Grapholita inopinata), oriental fruit moth (Grapholita molesta), soybean pod border (Leguminivora glycinivorella), mulberry leafroller (Olethreutes mori), tea leafroller (Caloptilia thevivora), Caloptilia sp. (Calopilia zachrysa), apple leafminer (Phyllonorycter ringoniella), pear barkminer (Spulerring astaurota), common white (Piers rapae crucivora), tabacco budworm (Heliothis sp.), codling moth (Laspey resia pomonella), diamondback moth (Plutella xylostella), apple fruit moth (Argyresthia conjugella), peach fruit moth (Carposina niponensis), rice stem borer (Chilo suppressalis), rice leafroller (Cnaphalocrocis medinalis), tabacco moth (Ephestia elutella), mulberry pyralid (Glyphodes pyloalis), yellow rice borer (Scirpophaga incertulas), rice skipper (Parnara guttata), rice armyworm (Pseudaletia separata), pink borer (Sesamia inferens), common cutworm (Spodoptera litura), beet armyworm (Spodoptera exigua), etc.; HEMIPTERA including aster leafhopper (Macrosteles fascifrons), green rice leafhopper (Nephotettix cincticeps), brown rice planthopper (Nilaparvata lugens), whitebacked rice planthopper (Sogatella furcifera), citrus psylla (Diaphorina citri), grape whitefly (Aleurolobus taonabãe), sweetpotato whitefly (Bemisia tabaci), greenhouse whitefly (Trialeurodes vaporariorum), turnip aphid (Lipaphis erysimi), green peach aphid (Myzus persicae), Indian wax scale (Ceroplastes ceriferus), cottony citrus scale (Pulvinaria aurantii), camphor scale (Pseudaonidia duplex), San Jose scale (Comstockaspis perniciosa), arrowhead scale (Unaspis yanonensis), etc.; COLEOPTERA including soybean beetle (Anomala rufocuprea), Japanese beetle (Popillia japonica), tabacco beetle (Lasioderma serricorne), powderpost beetle (Lyctus brunneus), twenty-eight spotted ladybird (Epilachna vigintiotopunctata), adzuki bean weevile (Callosobruchus chinensis), vegetable weevil (Listroderes costirostris), maize weevil (Sitophilus zeamais), boll weevil (Anthonomus gradis gradis), rice water weevil (Lissorhoptrus oryzophilus), cucurbit leaf beetle (Aulacophora femoralis), rice leaf beetle (Oulema oryzae), striped flea beetle (Phyllotreta striolata), pine

shoot beetle (<u>Tomicus piniperda</u>), Colorado potato beetle (<u>Leptinotarsa decemlineata</u>), Mexican bean beetle (<u>Epilachna varivestis</u>), corn rootworm (<u>Diabrotica sp.</u>), etc.; DIPTERA including melon fly (<u>Dacus(Zeugodacus</u>) cucurbitae), oriental fruit fly (<u>Dacus(Bactrocera</u>) <u>dorsalis</u>), rice leafminer (<u>Agnomyza oryzae</u>), onion maggot (<u>Delia antiqua</u>), seedcorn maggot (<u>Delia platura</u>), soybean pod gall midge (<u>Asphondylia sp.</u>), muscid fly (<u>Musca domestica</u>), house mosquito (<u>Culex pipiens pipiens</u>), etc.; and TYLENCHIDA including root-lesion nematode (<u>Pratylenchus sp.</u>), coffer root-lesion nematode (<u>Pratylenchus coffeae</u>), potato cyst nematode (<u>Globodera rostochiensis</u>), root-knot nematode (<u>Meloidogyne sp.</u>), citrus nematode (<u>Tylenchulus semipenetrans</u>), Aphelenchus sp. (<u>Aphelenchus avenae</u>), chrysanthemum foliar (<u>Aphelenchoides ritzemabosi</u>), etc.

[0086] The agricultural and horticultural insecticide containing the phthalic acid diamide derivative of the general formula (I) of the present invention as an active ingredient has a marked insecticidal effect on the above-exemplified insect pests, sanitary insect pests, and/or nematodes, which are injurious to paddy field crops, upland crops, fruit trees, vegetables, other crops, flowers and ornament plants, and the like. Therefore, the desired effect of the agricultural and horticultural insecticide of the present invention can be obtained by applying the insecticide to the paddy field water, stalks and leaves of fruit trees, vegetables, other crops, flowers and ornament plants, soil, etc. at a season at which the insect pests, sanitary pests or nematodes are expected to appear, before their appearance or at the time when their appearance is confirmed.

[0087] In general, the agricultural and horticultural insecticide of the present invention is used after being prepared into conveniently usable forms according to an ordinary manner for preparation of agrochemicals.

[0088] That is, the phthalic acid diamide derivative of the general formula (I) and, optionally, an adjuvant are blended with a suitable inert carrier in a proper proportion and prepared into a suitable preparation form such as a suspension, emulsifiable concentrate, soluble concentrate, wettable powder, granules, dust or tablets through dissolution, dispersion, suspension, mixing, impregnation, adsorption or sticking.

[0089] The inert carrier used in this invention may be either solid or liquid. As the solid carrier, there can be exemplified soybean flour, cereal flour, wood flour, bark flour, saw dust, powdered tobacco stalks, powdered walnut shells, bran, powdered cellulose, extraction residues of vegetables, powdered synthetic polymers or resins, clays (e.g. kaolin, bentonite, and acid clay), talcs (e.g. talc and pyrophyllite), silica powders or flakes (e.g. diatomaceous earth, silica sand, mica and white carbon, i.e. synthetic, high-dispersion silicic acid, also called finely divided hydrated silica or hydrated silicic acid, some of commercially available products contain calcium silicate as the major component), activated carbon, powdered sulfur, powdered pumice, calcined diatomaceous earth, ground brick, fly ash, sand, calcium carbonate powder, calcium phosphate powder and other inorganic or mineral powders, chemical fertilizers (e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, urea and ammonium chloride), and compost. These carriers may be used alone or as a mixture thereof.

[0090] The liquid carrier is that which itself has solubility or which is without such solubility but is capable of dispersing an active ingredient with the aid of an adjuvant. The following are typical examples of the liquid carrier and can be used alone or as a mixture thereof. Water; alcohols such as methanol, ethanol, isopropanol, butanol and ethylene glycol; ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone, diisobutyl ketone and cyclohexanone; ethers such as ethyl ether, dioxane, Cellosolve, dipropyl ether and tetrahydrofuran; aliphatic hydrocarbons such as kerosene and mineral oils; aromatic hydrocarbons such as benzene, toluene, xylene, solvent naphtha and alkylnaphthalenes; halogenated hydrocarbons such as dichloroethane, chloroform, carbon tetrachloride and chlorobenzene; esters such as ethyl acetate, diisopropyl phthalate, dibutyl phthalate and dioctyl phthalate; amides such as dimethylformamide, diethylformamide and dimethylacetamide; nitriles such as acetonitrile; and dimethyl sulfoxide.

[0091] The following are typical examples of the adjuvant, which are used depending upon purposes and used alone or in combination in some cases, or need not to he used at all.

[0092] To emulsify, disperse, dissolve and/or wet an active ingredient, a surfactant is used. As the surfactant, there can be exemplified polyoxyethylene alkyl ethers, polyoxyethylene alkylaryl ethers, polyoxyethylene higher fatty acid esters, polyoxyethylene resinates, polyoxyethylene sorbitan mono-laurate, polyoxyethylene sorbitan monoleate, alkylarylsulfonates, naphthalenesulfonic acid condensation products, ligninsulfonates and higher alcohol sulfate esters.

[0093] Further, to stabilize the dispersion of an active ingredient, tackify it and/or bind it, there may be used adjuvants such as casein, gelatin, starch, methyl cellulose, carboxymethyl cellulose, gum arabic, polyvinyl alcohols, turpentine, bran oil, bentonite and ligninsulfonates.

[0094] To improve the flowability of a solid product, there may be used adjuvants such as waxes, stearates and alkyl phosphates.

[0095] Adjuvants such as naphthalenesulfonic acid condensation products and polycondensates of phosphates may be used as a peptizer for dispersible products.

[0096] Adjuvants such as silicon oils may also be used as a defoaming agent.

[0097] The content of the active ingredient may be varied as required. In dusts or granules, the suitable content thereof is from 0.01 to 50% by weight. In emulsifiable concentrates or flowable wettable powders, it is also from 0.01 to 50% by weight.

[0098] The agricultural and horticultural insecticide of the present invention is used to control a variety of insect pests in the following manner. That is, it is applied to a crop on which the insect pests are expected to appear or a site where the appearance of the insect pests is undesirable, as it is or after being properly diluted with or suspended in water or the like, in an amount effective for control of the insect pests.

- [0099] The applying dosage of the agricultural and horticultural insecticide of the present invention is varied depending upon various factors such as a purpose, insect pests to be controlled, a growth state of a plant, tendency of insect pests appearance, weather, environmental conditions, a preparation form, an application method, an application site and an application time. It may be properly chosen in a range of 0.1 g to 10 kg (in terms of the active ingredient) per 10 ares depending upon purposes.
- 10 [0100] The agricultural and horticultural insecticide of the present invention may be used in admixture with other agricultural and horticultural disease or pest controllers in order to expand both spectrum of controllable diseases and insect pest species and the period of time when effective applications are possible or to reduce the dosage.

[0101] Typical examples of the present invention are described below, but they should not be construed as limiting the scope of the invention.

EXAMPLES

15

Example 1

20 (1-1) Production of 3-chloro-N-[4-(1,1,2,2-tetrafluoroethoxy)-2-methylphenyl]phthalimide

[0102] In 10 ml of acetic acid were dissolved 0.55 g of 3-chlorophthalic anhydride and 0.67 g of 4-(1,1,2,2-tetrafluoroethoxy)-2-methylaniline, and the reaction was carried out with heating under reflux for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure and the resulting residue was washed with an ether-hexane mixed solvent to obtain 1.1 g of the desired compound.

Physical property: m.p. 121 - 122°C. Yield: 95%.

(1-2) Production of 3-chloro-N¹-[4-(1,1,2,2-tetrafluoroethoxy)-2-methylphenyl]-N²-isopropylphthalic acid diamide (compound No. 141) and 6-chloro-N¹-[4-(1,1,2,2-tetrafluoroethoxy)-2-methylphenyl]-N²-isopropylphthalic acid diamide (compound No. 239)

[0103] In 10 ml of dioxane was dissolved 1.1 g of 3-chloro-N-[4-(1,1,2,2-tetrafluoroethoxy)-2-methylphenyl] phthalimide, followed by adding thereto 0.5 g of isopropylamine, and the reaction was carried out at 80°C for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure and the resulting residue was purified by a silica gel column chromatography using a hexane/ethyl acetate (2/1) mixed solvent as an eluent, to obtain 0.4 g of the desired compound (compound No. 141) having an Rf value of 0.5 to 0.7 and 0.5 g of the other desired compound (compound No. 239) having an Rf value of 0.2 to 0.4.

Compound No. 141:

[0104]

40

45 Physical property: m.p. 202 - 204°C. Yield: 31%.

Compound No. 239:

50 **[0105**]

Physical property: m.p. 199 - 201°C. Yield: 39%.

Example 2

10

20

25

35

45

50

- (2-1) Production of N-(4-trifluoromethoxyphenyl)-3-nitrophthalimide
- 5 [0106] In 50 ml of acetic acid were dissolved 5.97 g of 3-nitrophthalic anhydride and 5.31 g of 4-trifluoromethoxy-aniline, and the reaction was carried out with heating under reflux for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure and the resulting residue was washed with an ether-hexane mixed solvent to obtain 10.2 g of the desired compound.

Physical property: m.p. 149 - 150°C. Yield: 97%.

- (2-2) Production of 3-amino-N-(4-trifluoromethoxyphenyl)phthalimide
- [0107] In a pressure vessel were placed 10.0 g of N-(4-trifluoromethoxyphenyl)-3-nitrophthalimide, 100 ml of acetic acid and 0.5 g of 5% palladium carbon, and catalytic reduction with hydrogen was carried out at a hydrogen pressure of 5 kg/cm². After completion of the reaction, the catalyst was filtered off and the filtrate was concentrated under reduced pressure. The resulting residue was washed with an ether-hexane mixed solvent to obtain 9.0 g of the desired compound.

Physical property: m.p. 161 - 162°C. Yield: 98%.

(2-3) Production of 3-bromo-N-(4-trifluoromethoxyphenyl)phthalimide

[0108] In 20 ml of acetic acid was dissolved 1.6 g of 3-amino-N-(4-trifluoromethoxyphenyl)phthalimide, and a solution of 0.35 g of sodium nitrite in 5 ml of concentrated sulfuric acid was added dropwise while maintaining the temperature at 15°C or lower. The resulting mixture was stirred at 15°C or lower for another 20 minutes to obtain a diazonium salt. The diazonium salt was slowly added to a mixture of a solution of 0.86 g of cuprous bromide in 50 ml of hydrobromic acid and 10 ml of toluene which was maintained at 80°C. The resulting mixture was stirred until foaming ceased. After completion of the reaction, the organic layer was washed with an aqueous sodium thiosulfate solution and an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then distilled under reduced pressure to remove the solvent, and the resulting residue was purified by a silica gel chromatography to obtain 1.3 g of the desired compound.

Physical property: m.p. 117 - 118°C. Yield: 67%.

(2-4) Production of 3-bromo-N¹-(4-trifluoromethoxyphenyl)-N²-isopropylphthalic acid diamide (compound No. 262) and 6-bromo-N¹-(4-trifluoromethoxyphenyl)-N²-isopropylphthalic acid diamide (compound No. 302)

[0109] From 1.3 g of 3-bromo-N-(4-trifluoromethoxyphenyl)phthalimide, 0.5 g of the desired compound (compound No. 262) and 0.7 g of the other desired compound (compound No. 302) were obtained in the same manner as in Example 1-2.

Compound No. 262:

[0110]

Physical property: m.p. 208 - 210°C. Yield: 33%.

Compound No. 302:

55 [0111]

Physical property: m.p. 210 - 212°C. Yield: 47%.

Example 3

15

20

25

- (3-1) Production of N-(4-difluoromethoxy-2-methylphenyl)-3-nitrophthalimide
- [0112] In 100 ml of acetic acid were dissolved 5.8 g of 3-nitrophthalic anhydride and 5.2 g of 4-difluoromethoxy-2-methylaniline, and the reaction was carried out with heating under reflux for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure and the resulting residue was washed with an ether-hexane mixed solvent to obtain 10.2 g of the desired compound.

10 Physical property: m.p. 163 - 164°C. Yield: 98%.

- (3-2) Production of N¹-(4-difluoromethoxy-2-methylphenyl)-N²-isopropyl-3-nitrophthalic acid diamide (compound No. 696)
- [0113] In 100 ml of dioxane was dissolved 10 g of N-(4-difluoromethoxy-2-methylphenyl)-3-nitrophthalimide, followed by adding thereto 2.5 g of isopropylamine, and the reaction was carried out for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure and the resulting residue was washed with ether to obtain 4.0 g of the desired compound.

Physical property: m.p. 148 - 149°C. Yield: 86%.

- (3-3) Production of 3-amino-N1-(4-difluoromethoxy-2-methylphenyl)-N2-isopropylphthalic acid diamide
- [0114] In a pressure vessel were placed 5 g of N¹-(4-difluoromethoxy-2-methylphenyl)-N²-isopropyl-3-nitrophthalic acid diamide, 50 ml of acetic acid and 0.25 g of 5% palladium carbon, and catalytic reduction with hydrogen was carried out at a hydrogen pressure of 5 kg/cm². After completion of the reaction, the catalyst was filtered off and the filtrate was concentrated under reduced pressure. The resulting residue was washed with an ether-hexane mixed solvent to obtain 4.0 g of the desired compound.

Physical property: m.p. 148 - 149°C. Yield: 86%.

- 35 (3-4) Production of N¹-(4-difluoromethoxy-2-methylphenyl)-3-iodo-N²-isopropylphthalic acid diamide (compound No. 387)
- [0115] In 20 ml of acetic acid was dissolved 1.89 g of 3-amino-N¹-(4-difluoromethoxy-2-methylphenyl)-N²-isopropyl-phthalic acid diamide, and 1.5 g of concentrated sulfuric acid was added under ice-cooling. While maintaining the resulting solution at 15°C or lower, a solution of 0.35 g of sodium nitrite in 0.5 ml of water was added dropwise. The resulting solution was stirred at 15°C or lower for another 20 minutes to obtain a diazonium salt. The diazonium salt was slowly added to a mixture of 50 ml of an aqueous solution containing 1.0 g of potassium iodide and 50 ml of chloroform which was maintained at 40°C. The resulting mixture was stirred until foaming ceased. After completion of the reaction, the organic layer was washed with an aqueous sodium thiosulfate solution and an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then distilled under reduced pressure to remove the solvent, and the resulting residue was purified by a silica gel chromatography to obtain 0.8 g of the desired compound.

Physical property: m.p. 207 - 209°C. Yield: 33%.

Example 4

- (4-1) Production of 3-iodo-2-N-isopropyl-phthalamic acid
- [0116] A solution of 0.67 g of isopropylamine in 5 ml of acetonitrile was added dropwise to a solution of 1.37 g of 3-iodophthalic anhydride in 10 ml of acetonitrile under ice-cooling, and the reaction was carried out with stirring at room temperature for another 5 hours. After completion of the reaction, the crystals formed in the reaction solution were collected by filtration and washed with a small volume of acetonitrile to obtain 1.45 g of the desired compound.

Yield: 87%.

¹H-NMR [CDCl₃/TMS, δ values (ppm)]
1.23(6H, d), 4.35(1H, m), 5.80(1H, d), 6.85(1H, broad), 7.07(1H, t), 7.93(1H, d), 7.96(1H, d).

5 (4-2) Production of 6-iodo-N-isopropyl-phthalic acid isoimide

[0117] In 10 ml of toluene was dissolved 0.45 g of 3-iodo-2-N-isopropyl-phthalamic acid, followed by adding thereto 0.85 g of trifluoroacetic anhydride, and the reaction was carried out with stirring for 30 minutes. After completion of the reaction, the solvent was distilled off under reduced pressure to obtain 0.43 g of the desired compound as a crude product. The obtained desired compound was used in the subsequent reaction without purification.

Physical property: m.p. 87.5 - 88.5°C.

(4-3) Production of 3-iodo-N¹-(4-pentafluoroethyl-2-methylphenyl)-N²-isopropyl-phthalic acid diamide (compound No. 372)

[0118] In 10 ml of tetrahydrofuran was dissolved 0.43 g of the 6-iodo-N-isopropyl-phthalic acid isoimide obtained in 4-2, followed by adding thereto 0.30 g of 4-pentafluoroethyl-2-methylaniline, and the reaction was carried out with stirring for 1 hour. After completion of the reaction, the solvent was removed from the reaction solution by distillation under reduced pressure, and the resulting residue was washed with ether-n-hexane to obtain 0.70 g of the desired compound.

Physical property: m.p. 195 - 196°C. Yield: 95%.

25 Example 5

35

45

[0119] (5-1) Production of ethyl 6-nitro-N-(4-chloro-2-methylphenyl)-phthalamate

[0120] In 30 ml of tetrahydrofuran was dissolved 1.29 g of 3-nitro-2-ethoxycarbonylbenzoyl chloride, followed by adding thereto 0.71 g of 4-chloro-2-methylaniline and 0.56 g of triethylamine, and the reaction was carried out with stirring for 30 minutes. After completion of the reaction, the reaction solution containing the desired compound was poured into water and the desired compound was extracted with ethyl acetate. The extracted solution was dried over anhydrous magnesium sulfate and distilled under reduced pressure to remove the solvent, and the resulting residue was purified by a silica gel column chromatography to obtain 1.7 g of the desired compound.

Physical property: m.p. 164 - 165°C. Yield: 94%.

- (5-2) Production of 3-nitro-N1-(4-chloro-2-methylphenyl)-N2-isopropyl-phthalic acid diamide (compound No. 664)
- 40 [0121] In 20 ml of dioxane was dissolved 1.7 g of ethyl 6-nitro-N-(4-chloro-2-methylphenyl)-phthalamate, followed by adding thereto 1.5 g of isopropylamine, and the reaction was carried out with stirring at 80°C for 1 hour. After completion of the reaction, the solvent was removed from the reaction solution containing the desired compound, by distillation under reduced pressure, and the resulting residue was purified by a silica gel column chromatography to obtain 1.5 g of the desired compound.

Physical property: m.p. 202 - 204°C. Yield: 85%.

Example 6

50

(6-1) Production of N-isopropyl-3,4-dichlorophthalamic acid

[0122] In 30 ml of tetrahydrofuran was dissolved 2.32 g of N-isopropyl-3,4-dichlorobenzamide, and 21 ml of s-BuLi (0.96 M/L) was slowly added while maintaining the temperature at -70°C. The resulting mixture was stirred at -70°C for 30 minutes, after which the cooling bath was removed. An excess of carbon dioxide was introduced into the reaction solution, and the thus treated solution was stirred at room temperature for 30 minutes to carry out the reaction.

[0123] After completion of the reaction, the reaction solution was poured into water and acidified with diluted hydrochloric acid, and the desired compound was extracted with ethyl acetate. The extracted solution was dried over anhy-

drous magnesium sulfate and distilled under reduced pressure to remove the solvent, and the crystals thus obtained were washed with an ether-hexane mixed solvent to obtain 2.4 g of the desired compound.

Physical property: m.p. 155 - 156°C. Yield: 86.9%.

(6-2) Production of N-isopropyl-3,4-dichlorophthalic acid isoimide

[0124] In 10 ml of toluene was dissolved 0.41 g of N-isopropyl-3,4-dichlorophthalamic acid, followed by adding thereto 0.42 g of trifluoroacetic anhydride, and the reaction was carried out with stirring at room temperature for 30 minutes. After completion of the reaction, the solvent was distilled off under reduced pressure to obtain 0.39 g of the desired compound as a crude product. The obtained desired compound was used in the subsequent reaction without purification.

15 (6-3) Production of 3,4-dichloro-N¹-(4-pentafluoroethyl-2-methylphenyl)-N²-isopropylphthalic acid diamide (compound No. 1222)

[0125] In 10 ml of acetonitrile was dissolved 0.39 g of N-isopropyl-3,4-dichlorophthalic acid isoimide, followed by adding thereto 0.34 g of 4-pentafluoroethyl-2-methylaniline, and the reaction was carried out with stirring for 2 hours. After completion of the reaction, the reaction solution was maintained at 0°C for 10 minutes and the crystals precipitated were collected by filtration and washed with hexane to obtain 0.61 g of the desired compound.

Physical property: m.p. 208 - 209°C. Yield: 84.1%.

Example 7

25

30

40

5

Production of 3-chloro-2-isopropylaminothiocarbonyl-N-(pentafluoroethyl-2-methylphenyl)benzamide (compound No. S-1)

[0126] In 20 ml of tetrahydrofuran was dissolved 1.06 g of N-(pentafluoroethyl-2-methylphenyl)-3-chlorobenzamide, and 7 ml of s-BuLi (0.96 M/L) was slowly added while maintaining the temperature at -70°C. The resulting mixture was stirred at -70°C for 30 minutes, after which the cooling bath was removed. A solution of 0.33 g of isopropyl isothiocyanate in 5 ml of tetrahydrofuran was poured into the reaction solution, and the resulting solution was stirred at room temperature for 30 minutes to carry out the reaction.

[0127] After completion of the reaction, the reaction solution was poured into water and acidified with diluted hydrochloric acid, and the desired compound was extracted with ethyl acetate. The extracted solution was dried over anhydrous magnesium sulfate and distilled under reduced pressure to remove the solvent, and the crystals thus obtained were washed with an ether-hexane mixed solvent to obtain 1.2 g of the desired compound.

Physical property: m.p. 162 - 164°C. Yield: 86%.

[0128] Typical preparation examples and test examples of the present invention are described below but they should not be construed as limiting the scope of the invention.

[0129] In the preparation examples, parts are all by weight.

Formulation Example 1

50 [0130]

	Each compound listed in Table 1	50 parts
15	Xylene	40 parts
	Mixture of polyoxyethylene nonylphenyl ether and calcium alkylbenzenesulfonate	10 parts

[0131] An emulsifiable concentrate was prepared by mixing uniformly the above ingredients to effect dissolution.

Formulation Example 2

5 [0132]

10

Each compound listed in Table 1	3 parts
Clay powder	82 parts
Diatomaceous earth powder	15 parts

15 [0133] A dust was prepared by mixing uniformly and grinding the above ingredients.

Formulation Example 3

[0134]

20

25

Each compound listed in Table 1	5 parts
Mixed powder of bentonite and clay	90 parts
Calcium lignin sulfonate	5 parts

[0135] Granules were prepared by mixing the above ingredients uniformly, and kneading the resulting mixture together with a suitable amount of water, followed by granulation and drying.

Formulation Example 4

[0136]

35

40

Each compound listed in Table 1	20 parts
Mixture of kaolin and synthetic high-dispersion silicic acid	75 parts
Mixture of polyoxyethylene nonylphenyl ether and calcium alkylbenzenesulfonate	5 parts

[0137] A wettable powder was prepared by mixing uniformly and grinding the above ingredients.

Test Example 1

Insecticidal effect on diamondback moth (Plutella xylostella)

[0138] Adult diamondback moths were released and allowed to oviposit on a Chinese cabbage seedling. Two days after the release, the seedling having eggs deposited thereon was immersed for about 30 seconds in a liquid chemical prepared by diluting a preparation containing each compound listed in Table 1 as an active ingredient to adjust the concentration to 500 ppm. After air-drying, it was allowed to stand in a room thermostated at 25°C. Six days after the immersion, the hatched insects were counted. The mortality was calculated according to the following equation and the insecticidal effect was judged according to the criterion shown below. The test was carried out with triplicate groups of 10 insects.

	[Number of hatched insects] [Number of hatched insects]	
Corrected mortality (9/.) -	in untreated group in treated group.	X 100
Corrected mortality (%) =	L in untreated group J L in treated group J [Number of hatched insects in untreated group]	A 100

Criterion:

[0139]

10

15

Effect	Mortality(%)
Α	100
В	99 - 90
С	89 - 80
D	79 - 50

20

[0140] The results obtained are shown in Table 4.

Test Example 2

25 Insecticidal effect on common cutworm (Spodoptera Litura)

[0141] A piece of cabbage leaf (cultivar; Shikidori) was immersed for about 30 seconds in a liquid chemical prepared by diluting a preparation containing each compound listed in Table 1 as an active ingredient to adjust the concentration to 500 ppm. After air-drying, it was placed in a plastic Petri dish with a diameter of 9 cm and inoculated with second-instar larvae of common cutworm, after which the dish was closed and then allowed to stand in a room thermostated at 25°C. Eight days after the inoculation, the dead and alive were counted. The mortality was calculated according to the following equation and the insecticidal effect was judged according to the criterion shown in Test

Example 1. The test was carried out with triplicate groups of 10 insects.

[0142]

35

40

Corrected mortality (%) =
$$\frac{\begin{bmatrix} \text{Number of alive larvae} \\ \text{in untreated group} \end{bmatrix} - \begin{bmatrix} \text{Number of alive larvae} \\ \text{in treated group} \end{bmatrix}}{[\text{Number of alive larvae in untreated group}]} \times 100$$

[0143] The results obtained are shown in Table 4.

45 Test Example 3

Insecticidal effect on rice leafroller (Cnaphalocrocis medinalis)

[0144] The lamina of a rice plant at the 6 to 8 leaf stage was immersed for about 30 seconds in a liquid chemical prepared by diluting a preparation containing each compound listed in Table 1 as an active ingredient to adjust the concentration to 500 ppm. After air-drying, the lamina was placed in a plastic Petri dish with a diameter of 9 cm whose bottom had been covered with a wetted filter paper. The lamina was inoculated with third-instar larvae of rice leafroller, after which the dish was allowed to stand in a room thermostated at 25°C and having a humidity of 70%. Four days after the inoculation, the dead and alive were counted and the insecticidal effect was judged according to the criterion shown in Test Example 1. The test was carried out with triplicate groups of 10 insects.

[0145] The results obtained are shown in Table 4.

Table 4

	5	

No	Test Example	Test Example 2	Test Example 3
1	D .	D	A
2	A	С	
3	С	A	
4	A		D .
7	A		
8	A	A	A
9	A		A
10	A	D	D
11	A	С	С
12	A	D	
13	D		D
14	A		
15	A		Α
16	A		
17	A		D
18	D		A
20	A		

Table 4 (Cont'd)

5	No	Test Example		Test Example
		1	2	3
	22	Α	D	
10	23	A		D
	24	Α	j	D
	25	Α		Α
15	26	Α		D
	27	A	Α	С
20	28			Α
	29	A	В	Α
	30	A	Α	A
25	31	A		
	32	A		
	33	A		
30	34	. A	С	
	37	A		j
	41	A		A
35	42	A	D	Α
	43	В	D	
40	44			A
	45	Α		A
	46	Α		В
45	47	Α	D	Α .
	48	Α	В	Α
	49	Α	Α	Α
50	50	Α	. A	Α

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	51	A		A
10	52	Α	A	A
	53	\mathbf{A}^{\cdot}		Α
	55	A	В	A
15	56	Α	Α	A
	58	A	Α	A
	59	A		
20	60	A	Α	A
	61	A	В	A
25	62	A.	Α	A
-	63	A	В	A
	64	A	В	A
30	65	Α	A	A
	66	Α	A	В
	67	A	A	A
35	68	A		
	69	A		A
	70	A		Α
40	71			D
	73	A		
45	74	A		,
	75	Α		Α
	76	С		В
50	77	Α	С	A

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
	78	1 A	2 A	3 A
10	79	A	A	D
	81			A
	83	Α	Α	Α
15	84	Α		
	86	В		В
	87	· A		A
20	88	A		
·	89	Α	В	A
25	90	Α	Α	В
	91	Α	A	A
	92	A		
30	93	A	A	A
	98	A		С
	99	A		A
35	100	A	Α	Α
	101	Α		
40	102	Α	D	A
	103	Α	С	A
	109	A	A	С
45	110	A		Α .
• •	111	A	C	В
	112	A	A	A
50	113	Α	В	A

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	114	A	A	A
10	115	A	·C	A
	116	Α	D	A
	117	A	Α	A
15	118	Α	Α	A
	119	Α	Α	Α
	120	Α	D	A
20	121	A	Α	A
	122	Α	A	A
25	123	Α		A
25	124	A	Α	A
	125	Α	В	A
30	126	Α	Α	Α
	127	Α	Α	A
	128	Α	. D	A
35	129	Α	A	A
	130	Α	A	A
	132	A	A	A
40	133	A	A	A
	134	A		A
45	135	A	Α	A
	136	. A	A	A
	137	A		A
50	138	Α	A	A

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example 2	Test Example
139	A	A	3 A
140	A	A	A
141	A	A	A
142			
142	A A	A	В
]		A	A
144	A	A	Α
145	Α	Α	Α
146	A	Α	Α
147	A	С	
148	A	Α	A
149	A	Α	. A
150	A	A	Α
151	Α		
152	Α	Α	Α
153	A		D
157	A	Α	Α
158	A	A	Α
159	A	Α	Α
161	Α	D	Α
162	Α	Α	В
163	Α	Α	A .
164	Α	Α	
165	Α	В	С
167	A	A	·A

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	168	. A		
10	169	Α	D	
	170	A	D	В
·	171	A		D
15	172	Α	Α	D
	173	A	D	D
20	174	A		٠
20	175	A		
	176	A	D	A
25	177	A	A	A
	178	A		A
	179	A		
30	180	A	Α	Α
	181		Α	
	183	Α	• В	
35	185	Α		
	186	D		
40	187	A		D
	188	D		D
	189	A		
45	190	A		
ı	191	A		A
×3.	192	A		
50	193	A	D	

55

Table 4 (Cont'd)

5	No
	19
10	19
	19
	19
15	19
	19
	20
20	20
	20
25	20
	20
	20
30	20
	20
	21
35	21
	21
40	21
40	21
	21
45	21
	21
•	21
	1

No	Test Example	Test Example 2	Test Example 3
194	A		
195	A		
196	Α		D
197	A	A	Α
198	Α	С	Α
199	· A		
200	Α		Α
201	Α	В	A
202	A		
203	A		
206	Α		A
207	A		
208	A		
209	A		В
210	A		D
211	A		Α
212	A	D	¹ A
213	A	A	Α
214	A	, A	Α
215	A	D	
216	A		Α .
217	A		A
218	A		С
219	A	D	A

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	220	A		A
10	221	Α	Α	Α
	222	Α	В	Α
	223	A	Α	A
15	225	Α	В	Α
	226	Α		A
	227	A		j :
20	228		В	A
	229	Α	D	Α
<i>2</i> 5	230	Α	С	. A
	231		В	Α
	232	A		Α
30	233	A		
	234	, A		Α .
	235	Α		Α
35	236	A	A	Α
	237	A		A
40	238	A		A
•	239	A	A	A
	240	Α		
45	241	A	В	Α .
	242	A	В	A
	243	A	A	В
50	244	A	С	

Table 4 (Cont'd)

5	
10	
15	
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example
245	A	D	
246	Α	В	В
248	Α	С	
249	Α	D	Α
250	Α		D
251	· A		Α
252	Α		
253	A	A	С
254	Α	A	
255	Α		Α
256	Α		
257	Α		В
258	Α		A
259	Α		D
261	Α	A	D
262	Α	A	D
263	A		Α
264	_	D	Α
265	Α		
266	Α	A	A
267	Α .	A	Α .
268	Α	A	A
269	A	A	A
270	Α	· A	A

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	271	A	A	A
10	272	A	A	Α
	273	A	D	D
	274	A	A	A
15	275	A	D	A
	276	A	A	Α
	277	Α .	A	Α
20	278	A	Α	A
	279	A	A	A
<i>25</i>	281	A	Α	A
	282	A	A	A
	283	A	A	A
30	284	A	A	A
	285	A	D	A
	286	A	A	A
35	287	A	A	A
	288	A	A	A
	289	A	A	A
40	290	A	A	A
	291	A	A	A
45	292	A	A	A
	293	A	A	A
	294	A	A	A
50	295	D		
				}

55

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	296	A	A	Α
10	297	A	A	В
	298	A	A	A
	299	A	A	A
15	300	A		. A
	301	A	A	D
20	302	A		D
20	303	A		D .
	304	A		
25	305	A	A	A
	306	A	A	A
	307	A		D
30	308	A		
	309	A	A	
•	310	A		
35	311	A		D
	312	A	A	A
40	313	A	A	A
	314	A		A
	315	A		A
45	316	A	A	Α .
	318	A	В	A
	319	A	В	В
50	320	A		D
	1	1	1	1

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	321	Α	A	
10	322	A		В
	323	A	С	A
15	324	A		Α .
	325	Α	Α	A
	326	A	ý.	A
20	327	A		A
	328	A	Α	A
	329	A	Α	A
25	330	A		A
	332	A		A
30	333	A		D
	334	A	С	С
	335	Α		В
35	336	A		D
	337	A		A
	338	A	В	A
40	339	A	В	A
	340	A		A
	341	A		A
45	342	A		
	343	A		
	345	A	В	A
50	346	Α	С	Α

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	347	A	В	· C
10	348	A		A
	349			A
	350	A	Α	A
15	351	Α	A	Α
	352			Α
	353	A	Α ·	Α
20	354	A	Α	A
	355	A	С	A
25	356	A	Α	A
	360	· A	D	Α
	361	A	A	A
30	362	A	A	Α
	363	A	A	A
	364	A	A	D ·
35	365	A	A	A
	366	A	, A	A
	367	A	A	A
40	368	A	A	A
	369	A	A	A
45	370	A	A	Α .
	371	A	A	A
50	372	A	A	A
	373	A	A	Α
		I	I	

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
	374	A	A	A
10	375	A	A	Α
	376	A		Α
15	377	Α		A
	378	A	D	Α
	379	A	A	Α
20	380	A	A	A
	381	A	A	Α
25	382	A	В	Α
	383	A		Α
	384	A		С
30	385	A	В	Α
	386	A	Α	Α
	387	A	Α	Α
35	388	A	Α	В
	389	· A	Α	Α
	390	A	Α	Á
	391	A	Α	Α
	392	A	Α	Α
45	393	A	Α	Α
	394	Α	Α	Α
	395	Α	Α	Α
50	396	A	Α	Α
	397	Α	Α	Α

55

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
*	000	1	2	3
	398	A		
10	399	A	A	A
	400	A	D	A
45	402	A		
15	403	A	В	A
	404	A	A	A
20	406	Α	Α	A
20	407	A	A	A
*	408	A	В	A
25	409	A	A	A
	410	A	A	A
	411	A		A
30	412	A		С
	413	A		С
	414	A		A
35	415			A
•	416	A	A	A
	417	Α .	A	A
40	418			A
	419	A	A	A
45	420	Α		D
45	421	A	. В	A
	422	A		
50	424	A	A	
	767			

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	427	A	2	D
				D
,,	428	A	.	
	429	A	D	_
15	430	Α	D	D
	431	A	Α	
	432	Α		Α
20	433	A		Α
-	434	Α		
	435	A	В	Α
25	436	A	. B	Α
	437	A	С	A :
	438	A	В	Α
30	439	A	A	Α
	440	A	С	В
	441	A		В
35	442	A		•
	443	A		D
	444	A		Α
40	445		В	A
	446	A	A	A
45	447	A	В	C
	448	A		Α
	449	A		
50	450	A		С

55

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	451	A	Α	
10	452	A	A	Α
•	453	A	D	A
	454	A	. A	A
15	455	A	В	A
	456	A		A
	457	Α .	A	В
20	458	A		
	459	Α		·
25	460	A	В	
	461	A		
	462	A		
30	463	A		
	464	A		A
	465	A		
35	466	A		A
	467	· A		A
	468	A	Α	В
40	469	A	Α	D
	470	Α	С	С
45	471	Α	Α	Α .
	472	Α		В
	473	A	Α	Α
50	474	Α	В	Α

Table 4 (Cont'd)

5	No	Test Example	Test
	475	A	
10	476	A	Į į
	477	A	
	478	A	
15	479	A	
	480	A	I
20 .	488	A	_ A
20 .	489	A	1
	490	A	1
25	491	A	1
	492	A	1
	493	A	
30	494	A	
	495	A	4
	496	A	
35	498	A	
	499	A	.
40	500	A	:
₩	501	A	
	502	A	
45	503	A	
	504	A	
	505	A	
50	506	A	

	No	Test Example	Test Example 2	Test Example 3
	475	A	·	D
	476	Α	Α	Α
	477	Α		С
	478	Α		
	479	Α		Α
	480	Α	В	Α
	488	A	A	Α
١	489	A	A	Α
	490	A	A	A
	491	A	A	A
	492	A	A	A
l	493	A	A	A
	494	A		A
	495	A	A	A
I	496	A	A	A
	498	A	A	A
	499	A	A	A
	500	A	В	A
	501	A	A	A
	502	A	A	A
	503	A	В	Α :
	504	A	Α	Α
	505	A	A	A
	506	A		

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example 2	Test Example 3
507	A	В	A
508	A	В	A
509	A	A	A
510	A	В	Ä
511	Α	A	A
512	A	A	A
513	A	Α	A
514	Α	Α	Α
515	Α		С
516	Α	Α	Α
517	Α	Α	Α
518	Α		В
519	A	A	A
520	Α		
521	A	A	Α
522	A	D	A
523	Α	A	A
524	A	A	Α
526	Α	Α	
527	A	Α	A
528	A		A
529	A	D	A
530	A		D
531	A	:	Α

55

Table 4 (Cont'd)

				•
5	No	Test Example	Test Example 2	Test Example 3
	532	A		A
10	533	A	Α	A
	534	A		A
	535	A	A	
15	536			A
	537	A		
20	538	A	A	A
20	539	A		
	540	A		
25	543	A		Α
•	544	A		A
	545	A		Α
30	546	A		A
	547	A	Α	D
	548	A	A	A
35	549	A	Α	D
	550	Α .	C	Α
40	551	A		Α
	552	A		В
	553	A	С	Α
45	554	A		Α
	555	A		В
	557	A	С	В
50	558	A	Α	Α
	1	1		

Table 4 (Cont'd)

	No	Test Example	Test Example 2	Test Example 3
	559	Α		
10	560	A		
	561	Α	С	A
	562	A		A
15	563	Α		A
	564	A		В
	565	Α		A
20	566	Α .		В
·	567	A	D	D
25	568	A	С	A
	569	A	Α	A
	570	A	Α	·
30	571	A	С	
-	573	Α		
	575	A		A
35	576	A		C
-)(-	577	A		A
	579	Α	Α .	A
40	580	A		A
	581	A	В	Α.
45	582	Α		Α
	584	Α	D	
	585	A		Α
50	586	Α		D

Table 4 (Cont'd)

5	
10	
15	
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example
587	Α		
588	Α		
589	A		Α
590	A		
591	A	D	
592	Α		
593	A		
594	A		
595	A	Α	Α
596	D		D
597	В		
598	Α		
599	Α	D	Α
600	Α		
601	Α		
602	Α		Α
603	В		С
604	A		D
605			С
606	Α	D	Α
607	A	A	A .
608	A		
609	A	В	A
610	Α	Α	С

55

Table 4 (Cont'd)

5	No.	Test Example	Test Example 2	Test Example
	611	A		A
10	612	A		D
	613	A		
	614	A		
15	615	A		
	616	A		D
,	617	A	Α	Α
20	618	A	A	A
	619	A	A	Α
25	621	A		
	622	A		
	623	A		Α
30	624	A		
	625	A	D	D
	626	A		
35	628	A	В	Α
	633	A	D	
	634	A		D
40	635	A	D	
	636	A	D	A _.
45	637	A		•
	638	В		
	639	Α .		
50	640	Α		

Table 4 (Cont'd)

5	
10	
15	•
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example
641	D	D	
642	Α		
643	Α		Α
644	Α	A	
645	Α		
646	Α	D.	
647	Α		В
648	A	D	Α
649	A		С
650	A	•	
652	Α		
653	A		
654		D	
656	\mathbf{A}_{\cdot}		Α
657	D		
658	A		
659	A		
660	Α		Α
661	В		D
662	Α		
663	Α	Α	D ,
664	A	Α	
665	Α	Α	В
666	A		D

55

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example 2	Test Example
667	A	A	A
668	A		
669	A	D	Α
670	A		D
671	·A		D
672	A		
673	A	D	D
674	A	D	A
675	A	Α	A
676	A	С	Α
677	Α		
678	Α		
679	Α		Α
680	Α		D
681	Α	Α	A
682	Α		Α
683	A	Α	Α
684	Α	Α	Α
686	Α	Α	Α
687	-A	D	D
688	Α	 	Α
689	Α	D	Α
690	Α		Α
691	Α	D	С

55

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	692	A	D	
10	693	A	A	
	694	A		A
	695	A	A	A
15	696	A	A	Α .
	697	A		A
	698	A	В	A
20	699	A	Α	D
	700	A	Α	Α
25	701	A	Α	A
	703	A	Α	A
	704	A		Α
30	705	A	D	A
	706	A	Α	
	708	D		
35	709	A	A	
	710	A	С	A
. <u>.</u> 40	711	A	C	A
40	712	A	A	Α
	713	A	В	D
45	714	A	Α	A .
	715	A	Α	Α
	716	A	A	Α
50	717	A		Α
	[

Table 4 (Cont'd)

5				
	No	Test Example	Test Example 2	Test Example 3
40	718	A		A
10	719	A	D	
	720	Α		
15	721	A _.		
	722	Α .		Α
	723	D		D
20	724	A		В
	725	Α	Α	
25	727	A	В	Α
	728	A		A
	729	A		A
30	732	A		
	733	A		
35	735			D
	737	A		
	738	D		
40	740	A		A
	741	A	A	A
45	742	A		
	743	D		
	744	С		
50	745	D		

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	749	Α		
10	750	Α		A
	751	Α		Α
	752	Α		
15	753	Α	Α	D
	755	Α		Α
20	758			A
	759			D
	765	•		Α
25	766	Α		
	767	Α	С	A
30	768	Α	В	A
	769	A		D
	770	A	A	A
35	771	Α		С
	772	A		Α
	773	A		A
40	774	A		A
	776	В		D
45	777	A		D
	778	Α		A
	780	Α	A	A
50	781	A	A	A
·				

55

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example 2	Test Example
782	Α		Α
783	Α	A	A
785		A	
788	С		С
790			A
791	Α		Α
793	A		
795	A	В	A
796	A		
797	A		С
798			A
799	A		A
800			С
801	A	A	· D
802	D		
803	A		A
808	A		
819	A	В	A
821	Α		A
822	D		D
824	A		
825	A		

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example 2	Test Example
826	A		A
827	Α		
830	С		•
831	D	D	
832	A		
833	Α		D
835	A		
836	Α		Α
837	Α	·X·	
838	Α	C	Α
839	A		С
840	. A		D
841	Α .	D	
842	Α	A	D
845	Α		
846	Α		
847		D	
848	A		
849	A	В	Α
850	Α		Α
851	A	D	Α
852	Α		D
854	Α		
855	A		

Table 4 (Cont'd)

5	
10	
15	
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example
856	Α		D
858	С	A	
859	D		
860	A		
861	Α		
862	A	D	D
863	A		В
864	A		
865	A		
866	D		
867	A		С
869	A	D	
870	A		
871	A		
872	A		С
874	A	С	A
875	A		
878	С		
879	A		\mathbf{A}_{\cdot}
880		D	
881	A	D	
888	D		

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	889	Α	i	A
10	890	Α	A	A
	891	A	A	A
	892	A		Α ·
15	893	Α	A	Α
	894	A	A	Α
20	895	Α	A	Α
20	901	A	D	A
	902	A		
25	903	A	• A	A
	904	A		
	905	A	. А	Α
30	906	A	D	A
	907	A	A	Α
	908	A	D	Α
35	909	A	A	Α
	910		,	Α
40	911	A		D
	912	Α		
	913	Α		
45	914	A		
	915	A	A	Α
	916	A		
50	917	A	Α	Α

Table 4 (Cont'd)

5	No
	918
10	919
	920
	924
15	925
	927
20	928
20	929
	930
25	931
	932
	933
30	934
	935
	936
35	937
	938
40	939
	940
	941
45	942
	943
	944
50	945

No	Test Example	Test Example	Test Example 3
010	1	2	3
918	A		
919	A		
920	Α		
924	A		
925	Α		A
927	Α	A	A
928	Α		A
929	Α	Α	A
930	A	A	A
931	Α	A	A
932	Α	A	A
933	A	:	A
934	A	A	A
935	A	A	A
936	A	A	A
937	A	A	A
938	A	A	
939	A	A	A
940	A	A	A
941	A	A	A
942	A	Α	A
943	Α	С	A
944	A	Α	Α
945	A	A	A

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example	Test Example
<u> </u>	1	2	3
946	Α		Α
947	Α	A	Α
949	Α	Α	Α
950			С
951	Α	A	Α
952	A		Α
953	A	A	Α
954	Α		Α
955	Α	Α	Α
956	Α		Α
957	A	A	Α
958	A		Α
959	A		Α
965	Α	C	
966	A		В
971	A	Α	Α
972		Α	
973	Α	Α	Α
974	A	Α	Α
975	Α		Α
976	A	Α	Α .
977	A	Α	Α
978	Α	С	Α
979	Α	Α	Α

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	980	A	Α	Α
10	981	Α	Α	A
,	982	Α		С
	983	A	Α	Α
15	984	A		
	985	A	Α	Α
_	986	A		Α .
20	987	A	Α	Α
	988	A		
25	989	A	A	A
	990	A		A
	991	A	A	A
30	992	A		
	993	A		A
	995	Α		A
35	996	Α	A	A
	997	A	Α	A
	998	A	A	A
40	999	A		A
	1000	A	D	A
45	1001	A	Α	A
	1002	Α	Α	A
	1003	A	A	A
50	1004	Α	A `	Α

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example	Test Example
	1	2	3
1005	A	A	A
1006	Α		
1007	Α	Α	A
1008	A		Α
1009	A	A	Α
1010	A		A
1011	Α	A	Α
1013	Α	A	Α
1014	A		Α
1015	Α	D	Α
1016			Α
1017	Α	A	Α
1018	Α	D	Α
1019	Α	A	Α
1020	Α	Α	Α
1021	Α		Α
1022	'A	A	Α
1023	Α	A	Α
1024	A	A	Α
1025	A		
1026	Α		Α .
1027			Α
1028	Α	A	Α
1031	Α	Α	Α

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	1032	A	2	A
10	1033	A	A	A
	1034	A		••
	1035	A	A	Α
15	1036	A	71	A
	1037	A	A	A
	1038	A	A	A
20	1039		\ /	
		A	A	A
	1040	A	Α	Α
25	1041	Α	Α	Α
	1042	Α	Α	Α
	1043	Α		Α
30	1044	A	Α	Α
	1045	A	Α	Α
	1046	A	Α	Α
35	1047	A		Α
	1048	A		Α
	1049	A	Α .	Α .
40	1050	A	Α	Α
	1051	A	Α	Α
45	1052	Α	Α	
45	1053	A	Α	Α
	1054	A	A	A
50	1055	A	A	A
	1000	A	A	А

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example 3
	1056	Α	A	A
10	1057	A		A
	1058	A	Α	A
	1059	Α		
15	1060	A	Α	A
	1061	A	A	A
	1062	A	Α	A
20	1063			A
	1064	A		A
25	1065	A		
	1066	A	С	A
	1067	A	Α	A
30	1068	Α	Α	Α
	1069	A	Α	Α
	1070	Α	Α	Α
35	1071	Α '	С	Α
	1072			A
40	1073	A	Α	A
***	1074	A	A	Α
	1075	A	Α	Α
45	1076	Α		A 25
	1077	Α		A
	1078	Α	Α	Α
50	1079	Α	Α	Α

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
		1	2	3
	1080	A	A	Α
10	1081	A	A	A
	1082	A		i
	1083	A		
15	1086	Α		A
	1087	A	Α	Α
22	1088	Α		Α
20	1089	A		Α
	1099	A		Α
25	1100	Α	С	Α
	1101	A	С	A
	1102	Α .	Α	A
30	1103	A		Α
	1104	A	Α	Α
	1105	A	Α	Α
35	1106	A		Α
	1107	A		Α
40	1108	A		Α
40	1109	A		Α
	1110	A.	C	Α
45	1111	A		Α
	1112	A	Α	Α
1	1113	A	Α	Α
50	1114	A	Α	Α

Table 4 (Cont'd)

5	
10	
15	
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example
1115	A	A	A
1116	A	A	A
1117	A	A	A
1118	A	A	Α
1119	A	A	Α
1120	A	A	Α
1121	Α	A	Α
1122	Α	Α	Α
1123	Α	Α	A
1124	Α	Α	A
1125	Α	A	Α
1126	Α	Α	Α
1127	A	Α	Α
1128	Α	Α	Α
1129	Α	Α	Α
1130	Α	Α	Α
1131	A	Α	Α
1132	Α		Α
1133	Α .		Α
1134	Α		Α
1135	Α		Α ,,
1136	Α		Α
1137	Α		
1138			Α

55

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	1139	Α		
10	1140			Α
	1141	A	A	A
	1142	A	Α	Α
15	1143	A	С	Α
	1144	Α	Α	A
	1145	, A		Α
20	1146	A	С	Α
	1147	Α	A	Α
25	1148	A	Α	A
	1149	A		Α
	1150	Α	Α	Α
30	1151	A	Α	Α
	1152	A	Α	A
	1153	A	A	Α
35	1154	Α	С	Α
	1155	A		A
	1156	Α	Α	Α
40	1157	Α		Α
	1158	· A	D	A
45	1159	Α	A	Α
	1160	Α	A	Α
•	1161	A	D	Α .
50	1162	Α	Α	Α

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	1163	A	Α	A
10	1164	Α	Α	Α
	1165	A	Α	A
	1166	A -	Α	A
15	1167	Α	Α	Α
•	1168	A	Α	Α
0.	1169	A	Α	A
20	1173	Α	Α	Α
	1174	A	Α	A
<i>25</i>	1175	A		
	1178	A		
	1179	Α	Α	<u>-</u>
30	1180	A		_
	1181	A		_
	1182	Α		_
35	1183	- A	Α	_
	1184	A	Α	_
	1185	A	С	_
40	1186	Α	A	_
	1187	A	Α	_
45	1188	A	A	
	1189	A	A	
	1190	A	A	A
50	1191	A	A	A

Table 4 (Cont'd)

5	No.	Test Example	Test Example 2	Test Example
	1192	A	A	A
10	1193	A	A	A
	1202	A	A	A.
	1203	A	D	A
15	1204	A	A	A
	1205	A	A	A
	1205	A	A	A
20	1207	A	A	A
			A	
	1208	A	70	A
25	1209	A	D	A
	1210	Α	Α	A
	1211	A	A	A
30	1212	A	Α	A
	1221	A	A	A
	1222	A	A	A
35	1223	A	A	A
	1224			В
40	1225			A
<i>40</i>	1226			A
	1227	Α .	A	A
45	1228	A	A	Α .
-	1229	A	A	A
	1230	A	A	A
50	1231	A	A	A
				Ī

Table 4 (Cont'd)

5	No	Test Example		Test Example
		1	2	3
	1232	A	Α	Α
10	1233	A	Α	-
	1234	Α	Α	-
	1235	A	D	_
15	1236	A	Α	_
	1237	A	A	_
20	1238	A		_
	1245	A	Α	-
	1246	A	· A	. A
25	1247	С		
	1248	A	A	A
	1249	Α		
30	1250	A	A	A
	1251	A	A	A
	1256	A	A	_
35	1257	A	A	_
	1258	A	A	_
40	1259	A		_
7	1260	A		
	1261	A	A	_
45	1262	A	A	Α .
	1263	A	A	A
	1264	A	A	A
50	1266	A		A
	I	I	ŀ	1

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	1277	A	A	A
10	1278	A	A	A
	1280	A	A	A
	1281	A		A
15	1283	A	_	_
	1284	A	_	Α
	1285	A	_	A
20	1287	A	Α	A
	1288	A		
25	1291	A		
-	1293			A
	1294	A	A	A
30	1295	A	A	A
	1296	A	A	A
	1297	A	A	A
35	1298	A	A	A
	1299	A	A	A
	1300	A	A	A
40	1301	A		A
	1303	A	A	A
45	1304	A		A
	1305	A	A	A
	1306			A
50	1307	A	Α	A

Table 4 (Cont'd)

5		m (P)	m 4 D- 1	6 (2)
v	No	Test Example	lest Example 2	Test Example 3
	1308			C
10	1309	A	A	A
	1310			В
	1311	A	A	A
15	1312	A		A
	1313	A	A	A
	1314	A	A	A
20	1315	A	A	A
	1316	A	A	Α
25	1317	Α	С	Α
25	1318	Α		Α
	1319	A	_	-
30	1321	Α	_	A
	1322	Α		
	1323	A		Α
35	1325	A		A
	1327	Α		
	1328	A		
40	1330	A	A	A
	1331	Α		Α
45	1332	Α		
	1333	A		Α
	1335	A	С	A
50	1337	Α		A
	l	I	l .	I

Table 4 (Cont'd)

5	No	Test Example
•	1338	A
10	1339	Α
	1340	Α
	1341	A
15	1342	Α
	1343	A
20	1345	A
20	1346	A
·	1347	A
25	1348	A
	1349	A
	1350	
30	1351	A
	1352	A
	1353	A
35	1355	A
	1356	A
40	1358	A
40	1360	A
	1361	
45	1362	A
	1363	A
	1364	A
50	1365	A

No	Test Example	Test Example 2	Test Example 3
1338	A		
1339	Α		Α
1340	A		Α
1341	A		
1342	A		
1343	A		Α
1345	A		
1346	A		•
1347	A		
1348	A		
1349	Α		
1350			A
1351	A		A
1352	A		A
1353	A	A	A
1355	A	A	A
1356	A		
1358	A		C ·
1360	A		С
1361			Α
1362	A	A	A
1363	A		
1364	A	A	A
1365	A		

Table 4 (Cont'd)

5	No	Tes
	1366	
10	1367	
	1368	
	1370	
15	1372	
	1373	
20	1374	
20	1376	
	1379	
25	1381	
	1382	
	1383	
30	1384	
	1385	
	1386	
35	1387	
	1388	
40	1389	
40	1390	
	1392	
45	1393	
	1394	
	1395	
50	1398	

No	Test Example	Test Example 2	Test Example 3
1366	A	A	A
1367	Α		
1368	A		
1370	A		Α
1372	A		
1373	A		
1374	A		
1376	A		
1379	A		,
1381	A		С
1382	A	Α	Α
1383	A	Α	Α
1384	A	Α	A
1385	Α	Α	A
1386	Α	Α	Α
1387	Α	Α	A
1388	A	D	A
1389	A	Α	A
1390	A		Α
1392	A		A
1393	A		A .
1394	Α		Α
1395	Α		
1398	A		
			•

Table 4 (Cont'd)

5	
10	
15	
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example
1399	Α	A	A
1400	A	Α	A
1401	A		
1402	A	A	A
1404	Α	С	Α
1406	A	A	Α
1409	A		Α
1410	A		Α
1411	A		Α
1412	A		
1414	A	A	Α
1415	A		Α
1416			Α
1417	A	A	Α
1418	A	Α	Α
1419	A		Α
1420	A		Α
1421	A		
1423	A		
1424		A	Α
1427	A	Α	Α .
1428	A	Α	Α
1429	A	Α	Α
1430	A	D	A

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	1431	A		
10	1432			Α
	1434	A		
15	1436	A	Α	Α
	1438	A	Α	Α
	1439	A ·		С
20	1440	A		
	1441	A		A
	1442	Α		
25	1443	A		·
	1445	Α	A	A
30	1446	Α		
30	1447	Α	Α	Α
	1448	Α	Α	Α
35	1449	Α	Α	Α
	1450	Α	Α	A
	1451	Α		
40	1452	Α	A	A
	1453	A		
45	1454	Α	Α	A
	1456	Α		Α .
	1457	Α		Α
50	1458	Α	Α	Α
		1		

Table 4 (Cont'd)

5	
10	
15	
20	
25	
30	
35	
40	
45	

No	Test Example	Test Example 2	Test Example 3
1459	A	A	
1460	Α	C	Α
1461	A		Α
1464	A	A	A
1465	A		A
1466	A	A	Α
1467	A	A	Α
1468	A	С	A
1469	A		
1470	· A		A
1472	A	A	A
1473	A	A	A
1474	A	A	A
1475	A	A	A
1476	A		
1478	A	A	A
1479	A	A	A
1480			A
1481	A	A	A
1482	A	A	A
1484	A	A	A
1485	A	A	A

55

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	1486	A	A	
10	1487	A	Α	
	1488	A		A
	1489	A	A	A
15	1490	A	,	A
	1491	A	A	· A
	1492	A		A
20	1493	A		Α
	1494	A	A	Α
25	1495	A	A	Α
	1496	A	A	Α
•	1497	A	A	A
30	1498	A	A	A
	1499	A	A	A
	1500	A	A	A
35	1501	A	A	A
	1502	A	A	A
	1503	A	С	A
40	1504	A	A	A
	1505	A	A	A
45	1506	A	A	Α .
	1507	A	A	A
	1508	A	С	A
50	1509	A	С	A A
	•			

Table 4 (Cont'd)

5	No	Test Example		Test Example
	17.1	1	2	3
	1510	Α		Α
10	1511	Α		
	1512	A	Α	A
	1513	A		
15	1514	A		A
	1515	A		
	1516	A	A	A
20	1517	A		A
	1518	Α		·
25	1519	A		A
	1520			Α
	1521	A		
30	1522	A	Α	A
	1523	A		A
	1524	Α .	Α	Α
35	1525	A	Α	A
	1526	A	A	A
	1527	A		Α
40	1528	A		Α
	1529	A		Α
45	1530	A		Α
	1531	A	A	Α
	1532	A	Α	A
50	1533	A	Α	A

Table 4. (Cont'd)

5	No	Test Example	Test Example 2	Test Example
	1534	A	A	A
10	1535	Α	A	Α
	1536	A	A	A
	1537	A	A	A
15	1538	A	A	A
	1539	A	A	A
	1540	A	A	A
20	1541	A	A	A
	1542	A	D	A
25	1543	Α	Α	A
	1544	A	Α	A
	1545	Α	D	A
30	1546	A	С	Α
	1547	Α		A
	1548	A		A
35	1549	Α	D	Α
	1550	Α	A	A
	1551	Α	D	A
40	1552	A	A	A
	1553	Α		Α
45	1554	Α	A.	
	1555	A		Α
	1556	· A	A	A
50	1557	A	A	Α

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example	Test Example
	11	22	3
1558	Α	С	Α
1559	Α	Α	Α
1560	Α	Α	A
1561	Α .	Α	A
1562	Α	. A	Α
1563	Α	A	Α
1565	· A	A	A
1566	Α	A	A
1567	A	A	A
1568	A	A	A
1569	Α	A	A
1570			A
1571	A	A	A
1572	Α	A	A
1573	Α	A	A
1574	A	A	A
1575	A		A
1576	A	D	A
1577	A		
1578	A		A
1579	A		
1580	A	A	A
1581	A	A	Α
1582	Α		Α
			<u></u>

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
		1	2	3
	1583	A		A
10	1584	A		A
	1585	A	Α	A
	1586	A	С	A
15	1587	A	A	A
	1588	A		A
20	1589	Α		Α
	1590	Α	Α	Α
	1594	Α	A	-
25	1595	Α	Α	_
	1596	A	A	-
	1597	A	A	_
30	1598	A	Α	-
	1599	Α	Α	-
	1600	A	Α	_
35	1601	A	Α	
	1602	Α	Α	Α
40	1603	Α	Α	Α
	1604	Α	Α	Α
	1605	Α	С	-
45	1606	Α	Α	
	1607	A	Α	_
	1608	A	Α	_
50	1609	A	Α	-
				l

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
		1	2	3
	1610	A	Α	_
10	1611	A		_
	1612	Α	Α	-
	1613	A	Α	_
15	1614	Α	С	A
	1615	Α	Α	A
20	1617	Α		Α
	1618	A	С	A
	1619	A	A	_
25	1620	A		A
	1622	A	A	A
	1623	A	С	A
30	1624	A	D	A
	1625	A	A	A
	1626	A		A
35	1627	A	A	A
	1628	A	A	A
40	1629	A	A	A
40	1632	Α .		A
	1633	A	A	A
45	1634	A	A	A
	1635	A	A	Α
	1636	A	D	A
50	1637	A	A	A

Table 4 (Cont'd)

5	No	Test Example	Test Example	Test Example
		1	2	3
	1638	A	Α	A
10	1639	A	A	A
	1640	A	A	A
	1641	A		A
15	1642	A	A	A
	1643	A		A
20	1644	A	A	A
20	1645	A	A	A
•	1646	Α	A	A
25	1647	A	A	A
	1648	A	A	A
	1649	A	A	Α.
30	1650	A	A	A
	1651	A	A	A
	1652	A	A	A
35	1653	A	A	A
	1654	A	A	A
40	1655	A	A	A
40	1656	A	A	A
·	1658	A		
45	1659	A		A
	1660	A	A	_
	1661	A		_
50	1662	Α	A	_

Table 4 (Cont'd)

5	No	Test Example	Test Example	
		1	2	3
	1663	Α	Α	_
10	1664	A	A	_
	1665	A	A	_
	1666	A	A	_
15	1667	Α	A	_
	1668	Α	A	_
	1669	A	A	-
20	1670	A	A	_
	1671	A	A	_
25	1672	A	· A	_
	1673	A		_
	1674	A	A	-
30	1679	A	A	A
	1680	A	A	A
	1681	A	A	A
.35	1682	A	A	_
	1683	A		
	1684	A	A	_
40	1685	A	A	_
	1686	A	• А	
45	1689	A	A	–
	1690	A	A	_
	1691	A	A	–
50	1692	A		_

Table 4 (Cont'd)

5		
10		
15		
20		
25		
30		
35		
40		
45		

No	Test Example	Test Example 2	Test Example
1693	A	A	_
1694	A		_
1695	Α	Α	_
1696	Α	A	_
1697	Α	Α	_
1698	Α		_
1699	A	A	_
1700	A	D	_
1714	A	Α	Α
1715	Α	A	Α
1716	A	A	Α
1717	A	A	A
1722	Α	A	Α
1723	Α	A	A
1726	A		Α
1727	Α		Α
1732	Α	Α	Α
1733	Α	Α	A
1737	Α	С	Α
1742	Α		Α
1743	Α		A
1747	Α	С	Α
1748	Α		Α
1750	Α		Α

55

Table 4 (Cont'd)

5	No	Test Example	Test Example 2	Test Example 3
	1752	Α		Α
10	1763	Α		. A
	1764	Α	A	Α
15	1765	Α	Α	Α
	1766	Α	Α	A
	1767	Α		
20	1768	Α		A
	1770	Α	-	A
25	1772	A	-	Α
	1773	Α	Α	A
	1774	Α	-	\mathbf{A}°
30	1775	Α		Α
	1776	Α	_	A
	1777	Α	·A	A
35	1778	Α		Α
	1799	Α	A	A
40	1800	Α	Α	A
	1801	Α	Α	A
	1802	Α		
45	1803	Α	Α .	A
	1804	Α		
50	1805	A		,

⁵⁵ In Table 4, "-" means that test is not conducted.

Claims

5

10

15

20

25

30

35

40

45

50

55

A phthalic acid diamide derivative represented by the general formula (I).

$$\begin{array}{c}
X_{1} \\
\downarrow \\
C-N(\mathbb{R}^{1}) \mathbb{R}^{2}
\end{array}$$

$$\begin{array}{c}
Y_{m} \\
\downarrow \\
Z^{2}
\end{array}$$

wherein R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a cyano group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group or a group of the formula - A^1 - Q_ℓ (wherein A^1 is -O-, -S-, -SO₂-, -C(=O)-, a group of the formula - $N(R^4)$ - (wherein R^4 is a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenylcarbonyl group, or a substituted phenylcarbonyl group having at least one substituent which may be the same or different, and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a cyano group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different), a C_1 - C_6 alkylene group, a C_3 - C_6 alkenylene group or a C_3 - C_6 alkynylene group;

(1) when A^1 is -O- or a group of the formula -N(R^4)-(wherein R^4 is the same as defined above), then Q is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkynyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group and a di- C_1 - C_6 alkylamino group which may be the same or different, a phenyl- C_1 - C_4 alkyl group or a substituted phenyl- C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6

(2) when A1 is -S-, -SO2- or -C(=O)-, then Q is a C₁-C6 alkyl group, a halo-C₁-C6 alkyl group, a C3-C6 alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a C₁-C₆ alkoxy group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a C₁-C₆ alkoxycarbonylamino group, a C1-C6 alkoxycarbonyl-C1-C6 alkylamino group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1- C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a phenylamino group, a substituted phenylamino group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may

be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1- C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or a pyrazolyl group), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁- C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different,

10

15

20

25

30

35

40

45

50

55

(3) when A¹ is a C₁-C₈ alkylene group, a C₃-C₆ alkenylene group or a C₃-C₆ alkynylene group, then Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C1-C6 alkyl group, a C3-C6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a di- C_1 - C_6 alkoxyphosphoryl group which may be the same or different, a di-C₁-C₆ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-Ce alkylsulfinyl group, a C1-Ce alkylsulfonyl group, a halo-C1-Ce alkylsulfonyl group, a mono-C1-Ce alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkythio group, a halo-C1-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -Z3-R5 (wherein Z3 is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R6)-(wherein R6 is a hydrogen atom, a C1-C6 alkylcarbonyl group, a halo-C1-C6 alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a phenyl C1-C4 alkoxycarbonyl group, or a substituted phenyl C1-C4 alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different); and

R⁵ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆

alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₁-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkylcarbonyl group, a halo C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 -C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl C1-C4 alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkytthio group, a C_1 - C_6 alkytsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkył group, a halo-C₁-C₆ alkył group, a C₂-C₆ alkenył group, a halo-C₂-C₆ alkenył group, a C₂-C₆ alkynył group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different);

ℓ is an integer of 1 to 4); further,

10

15

20

25

30

35

40

45

50

55

 R^1 and R^2 may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X may be the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, or a group of the formula -A2-R7 (wherein A2 is -O-, -S-, -SO-, -SO2-, -C(=O)-, -C(=NOR8)- (wherein R^8 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6

10

15

20

25

30

35

40

45

50

55

alkenyl group, a C_3 - C_6 alkynyl group, a C_3 - C_6 cycloalkyl group, a phenyl- C_1 - C_4 alkyl group, or a substituted phenyl- C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkynyl group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a halo- C_2 - C_6 alkenylene group, a halo- C_3 - C_6 alkynylene group;

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo C_1 - C_6 alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_3 - C_6 cycloalkenyl group, a halo-C₃-C₆ cycloalkenyl group, a di-C₁-C₆ alkoxyphosphoryl group which may be the same or different, a di- C_1 - C_6 alkoxythiophosphoryl group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, or a group of the formula -A3-R9 (wherein A3 is -C(=O)-, -SO2-, a C1-C6 alkylene group, a halo-C1-C6 alkylene group, a C2-C6 alkenylene group, a halo-C2-C6 alkenylene group, a C₃-C₆ alkynylene group, or a halo-C₃-C₆ alkynylene group,

(i) when A3 is -C(=O)- or -SO2-, then R9 is a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a sub-

5

10

15

20

25

30

35

40

45

50

55

stituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkenyl group, a C_2 - C_6 alkenyl group, a C_2 - C_6 alkenyl group, a C_2 - C_6 alkenyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylyl group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different,

(ii) when A3 is a C1-C6 alkylene group, a halo-C1-C6 alkylene group, a C2-C6 alkenylene group, a halo-C2-C6 alkenylene group, a C3-C6 alkynylene group or a halo-C3-C6 alkynylene group, then R⁹ is a hydrogen atom, a halogen atom, a cyano group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group, a sulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A⁴-R¹⁰ (wherein A^4 is -O-, -S-, -SO-, -SO₂-, -C(=O)-, or a group of the formula -N(R^{11})- (wherein R^{11} is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 -C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkythio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group, a sulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different); and

 R^{10} is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a C_3 - C_6 alkynyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group, a bennyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkynyl group, a C_2 - C_6 alkynyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_2 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_2 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_2 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_2 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_2 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_1 - C_2 - C_1 - C_2 - C_1 - C_2 - C_2 - C_3 - C_1 - C_2 - C_3 -

5

10

15

20

25

30

35

40

45

50

55

halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylsulfining group, a halo- C_1 - C_6 alkylsulfining group, a halo- C_1 - C_6 alkylsulfining group, a halo- C_1 - C_6 alkylsulfining group, a halo- C_1 - C_6 alkylsulfining group, a halo- C_1 - C_6 alkylsulfining group, a halo- C_1 - C_6 alkylsulfining group which may be the same or different, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylsulfinyl

(2) when A2 is -C(=O)- or a group of the formula -C(=NOR8)-(wherein R8 is the same as defined above), then R7 is a C1-C6 alkyl group, a halo C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a mono- C_1 - C_6 alkylamino group, a di- C_1 - C_6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁- C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having one or more substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- $\textbf{C}_1\textbf{-}\textbf{C}_6 \text{ alkyl group, a } \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkenyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkenyl group, a } \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_6 \text{ alkynyl group, a halo-} \textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}\textbf{C}_2\textbf{-}$ C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁- C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 -C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different,

(3) when A^2 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group, then R^7 is a hydrogen atom, a halogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_1 - C_6 alkylyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group,

5

10

20

25

30

35

40

45

50

55

a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R¹³)-(wherein R¹³ is a hydrogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a C2-C6 alkynyl group, a C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different); and R¹² is a hydrogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-

 C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is -C(=O)-, -SO₂-, a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C₂-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group;

(i) when A⁶ is -C(=O)- or -SO₂-, then R¹⁴ is a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a phenyl

5

10

15

group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocydic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a

 C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group

20

25

30

35

40

45

50

55

which may be the same or different; (ii) when A^6 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo-C2-C6 alkenylene group, a C2-C6 alkynylene group or a halo-C3-C6 alkynylene group, then R¹⁴ is a hydrogen atom, a halogen atom, a cyano group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a C₁-C₆ alkylcarbonyl group, a halo-C₁-C₆ alkylcarbonyl group, a C1-C6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a $\hbox{halo-C$_1$-C$_6$ alkyl group, a C_2-C$_6$ alkenyl group, a halo-C$_2$-C$_6$ alkenyl group, a C_2-C$_6$ alkynyl group, a C_2 a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsutfonyl group, a halo-C₁-C₆ alkylsutfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆- alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or dif-

ferent and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a $C_1\text{-}C_6$ alkyl group, a halo- $C_1\text{-}C_6$ alkyl group, a $C_2\text{-}C_6$ alkynyl group, a $C_2\text{-}C_6$ alkynyl group, a $C_1\text{-}C_6$ alkynyl group, a $C_1\text{-}C_6$ alkynyl group, a $C_1\text{-}C_6$ alkylthio group, a $C_1\text{-}C_6$ alkylthio group, a $C_1\text{-}C_6$ alkylsulfinyl group, a halo- $C_1\text{-}C_6$ alkylsulfinyl group, a halo- $C_1\text{-}C_6$ alkylsulfinyl group, a halo- $C_1\text{-}C_6$ alkylsulfinyl group, a mono- $C_1\text{-}C_6$ alkylsulfinyl group and a di- $C_1\text{-}C_6$ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a $C_1\text{-}C_6$ alkyl group, a halo- $C_1\text{-}C_6$ alkynyl group, a $C_2\text{-}C_6$ alkenyl group, a halo- $C_2\text{-}C_6$ alkenyl group, a $C_1\text{-}C_6$ alkynyl group, a halo- $C_2\text{-}C_6$ alkoyy group, a $C_1\text{-}C_6$ alkylsulfinyl group, a halo- $C_1\text{-}C_6$ alky

n is an integer of 1 to 4;

5

10

15

20

25

30

35

40

45

50

55

further, X may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole), by combining together with the adjacent carbon atoms in the phenyl ring, and said condensed ring may have at least one substituents, which may be the same or different, and selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylaulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different:

Y is the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a di- C_1 - C_6 alkoxyphosphoryl group which may be the same or different, a di- C_1 - C_6 alkoxythiophosphoryl group which may be the same or different, a benyl group, a substituted phenyl group having at least one substitutent which may be the same or different and is selected from the group consisting of a hal-

ogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, or a group of the formula -A²-R⁷ (wherein A² and R⁷ are the same as defined above);

m is an integer of 1 to 5;

5

10

15

20

25

30

35

40

45

50

55

further, Y may form a condensed ring (the condensed ring is the same as defined above), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituents, which may be the same or different, and selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein said heterocyclic group is the same as defined above) having at lease one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different;

Z¹ and Z² are each represents an oxygen atom or a sulfur atom; provided that,

5

10

15

20

25

30

35

40

45

50

55

- (1) when X, R^1 and R^3 are hydrogen atoms at the same time; \underline{m} is an integer of 2; Y at 2-position is a fluorine atom and Y at 3-position is a chlorine atom; then R^2 is not ethyl group, isopropyl group, cyclohexyl group, 2-propenyl group, methylthiopropyl group and α -methylbenzyl group,
- (2) when X and R^3 are hydrogen atoms at the same time; \underline{m} is an integer of 2; Y at 2-position is a fluorine atom and Y at 3-position is a chlorine atom; then the 4 to 7 membered ring by combining R^1 and R^2 to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom is not morpholino group.
- (3) when X, R¹ and R³ are hydrogen atoms at the same time; and R² is 1,2,2-trimethyl-propyl group; then Y is not a hydrogen atom,
- (4) when X, R^1 and R^3 are hydrogen atoms at the same time; R^2 is 2,2-dimethylpropyl group; and \underline{m} is an integer of 1; then Y is not 2-ethoxy group, and
- (5) when X, R^1 and R^3 are hydrogen atoms at the same time; and R^2 is <u>tert</u>-butyl group group; and \underline{m} is an integer of 1; then Y is not 4-chlorine atom, 2-nitro group, 4-nitro group, 3-methoxy group, 4-methoxy group and 2,6-dimethyl groups.
- The phthalic acid diamide derivative according to Claim 1, wherein R1, R2 and R3 may be the same or different, and are each a hydrogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, or a group of the formula -A¹- Q_{ℓ} (wherein A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C_3 - C_6 alkynylene group; and Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C1-C6 alkyl group, a C3-C6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a di- C_1 - C_6 alkoxyphosphoryl group which may be the same or different, a di-C₁-C₆ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula -Z3-R5 (wherein Z3 is -O-, -S-, -SO2- or a group of the formula -N(R6)- (wherein R6 is a hydrogen atom, a C₁-C₆ alkylcarbonyl group, a halo-C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenyl C₁-C₄ alkoxycarbonyl group, or a substituted phenyl C₁-C₄ alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group); and

 $m R^5$ is a hydrogen atom, a $m C_1$ - $m C_6$ alkyl group, a halo $m C_1$ - $m C_6$ alkyl group, a $m C_3$ - $m C_6$ alkynyl group, a halo- $m C_3$ - $m C_6$ alkynyl group, a halo- $m C_3$ - $m C_6$ alkynyl group, a halo- $m C_3$ - $m C_6$ alkynyl group, a halo- $m C_3$ - $m C_6$ alkylcarbonyl group, a halo- $m C_3$ - $m C_6$ alkylcarbonyl group, a halo- $m C_1$ - $m C_6$ alkylcarbonyl group, a halo- $m C_1$ - $m C_6$ alkylcarbonyl group, a halo- $m C_1$ - $m C_6$ alkylcarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a $m C_1$ - $m C_6$ alkylthio group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a $m C_1$ - $m C_6$ alkylsulfinyl group, a $m C_1$ - $m C_6$ alkylsulfinyl group, a $m C_1$ - $m C_6$ alkylsulfinyl group, a $m C_1$ - $m C_6$ alkylsulfonyl group and a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a phenyl $m C_1$ - $m C_4$ alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfinyl group, a halo- $m C_1$ - $m C_6$ alkylsulfin

the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group); and $\underline{\ell}$ is an integer of 1 to 4);

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X may be the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C₃-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁- C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula $-A^2-R^7$ (wherein A^2 is -O-, -S-, -SO-, -SO₂-, -C(=O)-, -C(=NOR⁸)-(wherein R⁸ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a C3-C6 cycloalkyl group, a phenyl-C1-C4 alkyl group, or a substituted phenyl-C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkytthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group), a C_1 - C_6 alkylene group, a halo-C₁-C₆ alkylene group, a C₂-C₆ alkenylene group, a halo-C₂-C₆ alkenylene group, a C₂-C₆ alkynylene group or a halo-C3-C6 alkynylene group;

30

35

40

45

50

55

5

10

15

20

25

(1) when A² is -O-, -S-, -SO- or -SO₂-, then R⁷ is a halo-C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a $C_1 - C_6 \text{ alkylthio group, a halo-} \\ C_1 - C_6 \text{ alkylthio group, a } C_1 - C_6 \text{ alkylsulfinyl group, a halo-} \\ C_1 - C_6 \text{ alkylsulfinyl group,$ group, a C₁-C₆ alkylsulfonyl group and a halo-C₂-C₆ alkylsulfonyl group, or a group of the formula -A³-R⁹ (wherein A^3 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_3 - C_6 alkenylene group, a halo- C_3 - C_6 alkenylene group, a C_3 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group; R^9 is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^4 - R^{10} (wherein A^4 is -O-, -S-, -SO-, -SO₂- or -C(=O)-, and $\rm R^{10}$ is a $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkyl group, a $\rm C_3$ - $\rm C_6$ alkenyl group, a halo- $\rm C_3$ - $\rm C_6$ alkenyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl

5

10

15

20

25

30

35

40

45

55

group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group));

(2) when A² is -C(=O)- or a group of the formula -C(=NOR⁸)-(wherein R⁸ is the same as defined the above), then \mathbb{R}^7 is a \mathbb{C}_1 - \mathbb{C}_6 alkyl group, a halo \mathbb{C}_1 - \mathbb{C}_6 alkyl group, a \mathbb{C}_2 - \mathbb{C}_6 alkenyl group, a halo- \mathbb{C}_2 - \mathbb{C}_6 alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkylthio group, a mono-C1-C6 alkylamino group, a di-C1-C6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group,

(3) when A^2 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, C_2 - C_6 alkenylene group, a halo- C_2 -C₆ alkenylene group, a C₂-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group, then R⁷ is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula -A5-R12 (wherein A5 is -O-, -S-, -SO- or -SO2-; and R12 is a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C₂-C₆ alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group, or a halo- C_3 - C_6 alkynylene group; and R¹⁴ is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁- C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 -C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfinyl group group, a halo-C₁-C₆ alkylsulfinyl group group group group group group group group gr tonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a

 C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group)));

n is an integer of 1 to 4;

5

10

15

20

25

30

35

40

45

50

55

further, X may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituent, which may be the same or different, and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a phenyl group, a substituted phenyl group, a subs nyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; Y is the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula $-A^2$ - R^7 (wherein A^2 and R^7 are the same as defined above);

m is an integer of 1 to 5;

further, Y may form a condensed ring (the same as defined above), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituent, which may be the same or different, and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a chalo- C_1 - C_6 alkylsulfinyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkyls

 C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group; and Z^1 and Z^2 are each represents an oxygen atom or a sulfur atom.

The phthalic acid diamide derivative according to Claim 2, represented by the general formula (I-1),

$$\begin{array}{c|c}
X & Z^1 \\
\downarrow \\
C-N(R^1) & R^2 \\
\downarrow \\
Z^2
\end{array}$$

$$\begin{array}{c}
Y^3 \\
Y^2
\end{array}$$
(I-1)

15

20

25

30

35

40

45

50

55

10

(wherein, R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group or a group of the formula - A^1 - Q_ℓ (wherein, A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C_3 - C_6 alkynylene group; Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₁-C₆ alkyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a $di-C_1-C_6$ alkoxyphosphoryl group which may be the same or different, a $di-C_1-C_6$ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - Z^3 - R^5 (wherein Z^3 is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(\mathbb{R}^5)- (wherein \mathbb{R}^6 is a hydrogen atom, a \mathbb{C}_1 - \mathbb{C}_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenyl C₁-C₄ alkoxycarbonyl group, or a substituted phenyl C₁-C₄ alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfinyl group. fonyl group); and R^5 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 alkynyl group, a halo- C_3 - C_6 alkynyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkylcarbonyl group, a halo C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-

 C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group); and ℓ is an integer of 1 to 4); further,

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom:

X is a hydrogen atom or a nitro group;

5

10

15

20

25

30

35

40

45

50

55

 $\rm Y^1$ and $\rm Y^3$ may be the same or different and are each a hydrogen atom, a halogen atom, a $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkyl group, a $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylthio group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkyl group, a $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylthio group, a $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfonyl group, a pyridyloxy group, a substituted pyridyloxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group, a halo- $\rm C_1$ - $\rm C_6$ alkylsulfinyl group;

 Y^2 is a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfinyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group, and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^2 - R^7 (wherein A^2 is -O-, -S-, -SO-, -SO₂-, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_3 - C_6 alkynylene group and,

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a halo- C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁- C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 tonyl group, or a group of the formula -A3-R9 (wherein A3 is a halo-C1-C6 alkylene group, a halo-C3-C6 alkenylene group, a C₃-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group; and R⁹ is a hydrogen atom, a halogen atom, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group or a group of the formula -A⁴-R¹⁰ (wherein A⁴ is -O-, -S- or -SO₂-; and R^{10} is a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as

group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group)); (2) when A^2 is a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group or a halo- C_3 - C_6 alkynylene group, then R^7 is a hydrogen atom, a halogen atom, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group, and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^5 - R^{12} (wherein A^5 is -O-, -S-, -SO-

defined above), a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group

consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio

5

10

15

20

25

30

35

40

45

50

or -SO $_2$ -; and R 12 is a C $_3$ -C $_6$ cycloalkyl group, a halo-C $_3$ -C $_6$ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group; or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C2-C6 alkenylene group, a halo-C2-C6 alkenylene group, a C2-C6 alkynylene group or a halo-C3-C6 alkynylene group; and R^{14} is a hydrogen atom, a halogen atom, a halo- C_3 - C_6 cycloalkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkytthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo-C1-C6-alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group))); further,

Y and Y may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole) by combining to each other together with the adjacent Y3, said condensed ring may have at least one substituent, which is the same or different, selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 sulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; and Z^1 and Z^2 are each an oxygen atom or a sulfur atom).

The phthalic acid diamide derivative according to Claim 2, represented by the general formula (I-2),

50

$$X^{2}$$
 $C-N(R^{3})$
 Ym

(1-2)

5

10

15

20

25

30

35

40

45

50

55

(wherein, R1, R2 and R3 may be the same or different, and are each a hydrogen atom, a C3-C6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group or a group of the formula $-A^1$ - Q_ℓ (wherein, A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C₃-C₆ alkynylene group; Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₁-C₆ alkyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a $di-C_1-C_6$ alkoxyphosphoryl group which may be the same or different, a $di-C_1-C_6$ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo- C_1 - C_6 alkoyty group, a C_1 - C_6 alkythio group, a halo- C_1 - C_6 alkythio group, a C_1 - C_6 alkytsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -Z³-R⁵ (wherein Z³ is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R6)- (wherein R6 is a hydrogen atom, a C1-C6 alkylcarbonyl group a halo-C1-C6 alkylcarbonyl group, a C1-C6 alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 - C_1 - C_6 - C_1 - C_6 - C_1 - C_6 - C_1 - C_6 - C_1 - C_6 - C_1 - C_1 - C_6 - C_1 a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkoxycarbonyl group, or a substituted phenyl C₁-C₄ alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsul-

 R^5 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group and a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group); and £ is an integer of 1 to 4); further,

 R^1 and R^2 may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

 \dot{X}^1 and \dot{X}^2 may be the same or different and are each a halogen atom, a cyano group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfing group, a C_1 - C_6 alkylsulfing group, a halo- C_1 - C_6 alkylsulfing group, a C_1 - C_6 alkylsulfing group, a halo- C_1 - C_6 alkylsulfing group, a C_1 - C_6 alkylsulfong group; further, \dot{X}^1 and \dot{X}^2 may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole) by combining to each other, and said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a hal-

ogen atom, a C_1 - C_6 alkyi group, a halo- C_1 - C_6 alkyi group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkyl sulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; Y is the same or different, and are each a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of -A²-R⁷ (wherein A² is -O-, -S-, -SO-, -SO₂-, -C(=O)-, -C(=NOR 8)- (wherein R 8 is a hydrogen atom, a C $_1$ -C $_6$ alkyl group, a halo-C $_1$ -C $_6$ alkyl group, a C $_3$ -C $_6$ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a phenyl-C₁-C₄ alkyl

group, or a substituted phenyl- C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group), a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene

group, a halo-C2-C6 alkenylene group, a C2-C6 alkynylene group or a halo-C3-C6 alkynylene group;

10

15

20

25

30

35

40

50

55

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a halo-C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A³-R⁹ (wherein A^3 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_3 - C_6 alkenylene group, a halo- C_3 - C_6 alkenylene group, a C_3 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group; R^9 is a hydrogen atom, a halogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A⁴-R¹⁰ (wherein A⁴ is -O-, -S-, -SO-, -SO₂- or -C(=O)-, and R¹⁰ is a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the hetero-

10

15

20

25

30

35

40

45

50

55

cyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group);

(2) when A² is -C(=O)- or a group of the formula -C(=NOR⁸)-(wherein R⁸ is the same as defined the above), then R7 is a C1-C6 alkyl group, a halo C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxy group, a $C_$ group, a mono-C1-C6 alkylamino group, a di-C1-C6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group,

(3) when A² is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, C₂-C₆ alkenylene group, a halo-C₂-C₆ alkenylene group, a C₂-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group, then R⁷ is a hydrogen atom, a halogen atom, a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a C1-C6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula -A5-R12 (wherein A5 is -O-, -S-, -SO- or -SO2-; and R12 is a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula -A6-R14 (wherein A6 is a C1-C6 alkylene group, a halo-C1-C6 alkylene group, a C2-C6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group, or a halo- C_3 - C_6 alkynylene group; and R¹⁴ is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one

5

10

15

20

25

30

35

40

45

50

55

substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group

further, Y may form a condensed ring (which is the same as defined above) by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; Z^1 and Z^2 are each an oxygen atom or a sulfur atom).

5. The phthalic acid diamide derivative according to Claim 4, represented by the general formula (I-3),

$$X^{2} \xrightarrow{X^{1}} C-N(R^{1}) R^{2}$$

$$C-N(R^{3}) \xrightarrow{Y^{2}} Y^{2}$$

$$Z^{2} \xrightarrow{Y^{1}} Y^{2}$$

$$(I-3)$$

{wherein, R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group or a group of the formula $-A^1$ - Q_ℓ (wherein, A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C_3 - C_6 alkynylene group; Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxyphosphoryl group which may be the same or different, a di- C_1 - C_6 alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thianyl group, tetrahydropyranyl group, tetrahydrofuryl group, isoxazolyl group, oxazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl

5

10

15

25

30

35

40

45

50

55

group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -Z³-R⁵ (wherein Z³ is -O-, -S-, -SO-, -SO2- or a group of the formula -N(R6)- (wherein R6 is a hydrogen atom, a C1-C6 alkylcarbonyl group a halo-C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkoxycarbonyl group, or a substituted phenyl C₁-C₄ alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group. fonyl group,); and

R⁵ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkylcarbonyl group, a halo-C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfinyl group. fonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group); and £ is an integer of 1 to 4); further,

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X1 and X2 may be the same or different and are each a halogen atom, a cyano group, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁- C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; further, X¹ and X² may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole) by combining to each other, and said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a $C_1 - C_6 \text{ alkylthio group, a halo-} \\ C_1 - C_6 \text{ alkylthio group, a } C_1 - C_6 \text{ alkylsulfinyl group, a halo-} \\ C_1 - C_6 \text{ alkylsulfinyl group,$ group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group; Y^1 and Y^3 may be the same or different, and are each a hydrogen atom, a halogen atom, a C_1 - C_6 alkyl group,

a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a pyridyloxy group, or a substituted pyridyloxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group, and a halo- C_1 - C_6 alkylsulfonyl group, group, group, group, and a halo- C_1 - C_6 alkylsulfonyl group,

 Y^2 is a hydrogen atom, a halogen atom, a halo- C_3 - C_6 cycloalkyl group or a group of the formula - A^2 - R^7 (wherein A^2 -O-, -S-, -SO-, -SO₂-, a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group, and

10

15

20

25

30

35

40

45

50

55

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a halo- C_3 - C_6 cycloalkyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a substituted pyridyloxy group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A³-R⁹ (wherein A³ is a halo- C_1 - C_6 alkylene group, or a halo- C_3 - C_6 alkenylene group; and R^9 is a hydrogen atom, a halogen atom, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C₁-C₆ alkytsulfonyl group, or a group of the formula -A⁴-R¹⁰ (wherein A⁴ is -O-, -S-, -SO- or -SO₂-; R¹⁰ is a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, or a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group)),

(2) when A^2 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group, a halo- C_3 - C_6 alkynylene group, then R^7 is a hydrogen atom, a halogen atom, a halo- $\mathrm{C_3\text{-}C_6}$ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁ $m C_6$ alkylsulfinyl group and a halo- $m C_1$ - $m C_6$ alkylsulfonyl group, or a group of the formula - $m A^5$ - $m R^{12}$ (wherein $m A^5$ is -O-, -S-, -SO- or -SO $_2$ -; and R^{12} is a C_3 -C $_6$ cycloalkyl group, a halo-C $_3$ -C $_6$ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a $m C_2$ - $m C_6$ alkenylene group, a halo- $m C_2$ - $m C_6$ alkenylene group; and $m R^{14}$ is a hydrogen atom, a halogen atom, a halo-C₃-C₆ cycloalkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a phenylthio group, or a substituted phenylthio group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group));

further, Y^1 and Y^2 may form a condensed ring (the condensed ring is the same as defined above) by combining to each other together with Y^3 , and said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoy group, a halo- C_1 - C_6

a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a phenyl group, and a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group;

 Z^1 and Z^2 are each an oxygen atom or a sulfur atom}.

6. An agricultural and horticultural insecticides, which is characterized by containing, as the effective ingredient, a phthalic acid diamide derivative represented by the general formula (I),

$$Z^{1}$$

$$C-N(R^{1}) R^{2}$$

$$Ym$$

$$Z^{2}$$

$$Z^{2}$$

$$(I)$$

wherein R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a cyano group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group or a group of the formula - A^1 - Q_ℓ (wherein A^1 is -O-, -S-, -SO₂-, -C(=O)-, a group of the formula - $N(R^4)$ - (wherein R^4 is a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenylcarbonyl group, or a substituted phenylcarbonyl group having at least one substituent which may be the same or different, and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkoxy group, a halo- C_2 - C_6 alkoxy group, a C_1 - C_6 alkylyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different), a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C_3 - C_6 alkynylene group;

(1) when A^1 is -O- or a group of the formula -N(R^4)-(wherein R^4 is the same as defined above), then Q is a hydrogen atom, a C_1 -C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_3 -C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 -C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_2 -C₆ alkenyl group, a halo-C₂-C₆ alkynyl group, a C_1 -C₆ alkynyl group, a halo-C₁-C₆ alkynyl group, a C_1 -C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 -C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 -C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl-C₁-C₄ alkyl group or a substituted phenyl-C₁-C₄ alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 -C₆ alkyl group, a halo-C₂-C₆ alkenyl group, a C_2 -C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_1 -C₆ alkyl group, a halo-C₁-C₆ alkyl group, a halo-C₁-C

(2) when A^1 is -S-, -SO₂- or -C(=O)-, then Q is a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a C_3 - C_6 alkenyl group, a C_1 - C_6 alkoxy group, a mono- C_1 - C_6 alkylamino group, a di- C_1 - C_6 alkylamino group which may be the same or different, a C_1 - C_6 alkoxycarbonyl-amino group, a C_1 - C_6 alkoxycarbonyl- C_1 - C_6 alkylamino group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkenyl group, a C_1 - C_6 alkynyl group, a C_1 - C_6 alkynyl group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1

C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a phenylamino group, a substituted phenylamino group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 -C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or a pyrazolyl group), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkenyl group, a halo-C1-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁- C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different,

10

15

20

25

30

35

40

45

50

55

(3) when A¹ is a C₁-C₈ alkylene group, a C₃-C₆ alkenylene group or a C₃-C₆ alkynylene group, then Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C1-C6 alkyl group, a C3-C6 cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a di-C₁-C₆ alkoxyphosphoryl group which may be the same or different, a di-C1-C6 alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁- C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkytthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 -C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -Z3-R5 (wherein Z3 is -O-, -S-, -SO-, -SO2- or a group of the formula -N(R^6)-(wherein R^6 is a hydrogen atom, a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C1-C6 alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁- C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl C₁-C₄ alkoxycarbonyl

group, or a substituted phenyl C_1 - C_4 alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkynyl group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylsulfinyl group and a di- C_1 - C_6 alkylsulfinyl group which may be the same or different); and

R⁵ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C_1 - C_6 alkylcarbonyl group, a halo C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a phenyl C1-C4 alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkytthio group, a halo-C1-C6 alkytthio group, a C1-C6 alkylsulfinyl group, a hato-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 $a\ C_1-C_6\ alkylsulfinyl\ group,\ a\ halo-C_1-C_6\ alkylsulfinyl\ group,\ a\ C_1-C_6\ alkylsulfonyl\ group,\ a\ halo-C_1-C_6\ onyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different);

Lis an integer of 1 to 4); further,

10

20

25

30

35

40

45

50

55

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom:

X may be the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above)

5

10

20

25

35

40

45

50

55

having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2- C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, or a group of the formula -A2-R7 (wherein A2 is -O-, -S-, -SO-, -SO2-, -C(=O)-, -C(=NOR8)- (wherein R^8 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a phenyl-C₁-C₄ alkyl group, or a substituted phenyl- C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylsulfonyl group and a di- C_1 - C_6 alkylsulfonyl group. C_6 alkylamino group which may be the same or different), a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C2-C6 alkenylene group, a halo-C2-C6 alkenylene group, a C2-C6 alkynylene group or a halo-C3-C₆ alkynylene group;

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo C_1 - C_6 alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₃-C₆ cycloalkenyl group, a halo-C₃-C₆ cycloalkenyl group, a di-C₁-C₆ alkoxyphosphoryl group which may be the same or different, a di-C₁-C₆ alkoxythiophosphoryl group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, or a group of the formula -A3-R9 (wherein A3 is -C(=O)-, -SO2-, a C1-C6 alkylene group, a halo-C1-C6 alkylene group, a C2-C6 alkenylene group, a halo-C2-C6 alkenylene group, a C₃-C₆ alkynylene group, or a halo-C₃-C₆ alkynylene group,

(i) when A^3 is -C(=O)- or $-SO_2$ -, then R^9 is a C_1-C_6 alkyl group, a halo- C_1-C_6 alkyl group, a C_1-C_6 alkylamino group, a di- C_1-C_6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1-C_6 alkyl group, a halo- C_1-C_6 alkyl group, a C_2-C_6 alkenyl group, a halo- C_2-C_6 alkenyl group, a C_1-C_6 alkynyl group, a halo- C_1-C_6 alkynyl group, a C_1-C_6 alkyl group, a halo- C_1-C_6 alkyl group, a C_1-C_6 alkyl group, a halo- C_1-C_6 alkyl group, a h

5

10

15

20

25

30

35

40

45

50

55

be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different,

(ii) when A^3 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo-C2-C6 alkenylene group, a C3-C6 alkynylene group or a halo-C3-C6 alkynylene group, then R⁹ is a hydrogen atom, a halogen atom, a cyano group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C1-C6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A4-R10 (wherein A⁴ is -O-, -S-, -SO-, -SO₂-, -C(=O)-, or a group of the formula -N(R¹¹)- (wherein R¹¹ is a hydrogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C3-C6 alkenyl group, a C3-C6 alkynyl group, a C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_2 - C_1 - C_2 - C_3 - C_4 - C_6 sulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different); and

 R^{10} is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a C_3 - C_6 alkynyl group, a halo- C_3 - C_6 alkynyl group, a C_3 - C_6 alkynyl group, a halo- C_3 - C_6 cycloalkyl group, a C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6

5

10

15

20

25

30

35

45

50

55

alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylaulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different));

(2) when A2 is -C(=O)- or a group of the formula -C(=NOR8)-(wherein R8 is the same as defined above), then R7 is a C1-C6 alkyl group, a halo C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a mono- C_1 - C_6 alkylamino group, a di- C_1 - C_6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having one or more substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group- a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group- a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different,

(3) when A^2 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group, then R^7 is a hydrogen atom, a halogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent

5

10

15

20

25

30

35

40

45

50

55

which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyi group, a halo- C_2 - C_6 alkynyi group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R¹³)-(wherein R¹³ is a hydrogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different); and R¹² is a hydrogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkynyl group, a group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl

5

10

15

20

25

30

35

40

45

50

55

group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_1 - C_6 alkynyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a mono- C_1 - C_6 alkylsulfinyl group and a di- C_1 - C_6 alkylamino group which may be the same or different, or a group of the formula - A^6 - A^{14} (wherein A^6 is -C(=O)-, - SO_2 -, a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a halo- C_2 - C_6 alkylene group, a halo- C_3 - C_6 alkynylene group;

(i) when A^6 is -C(=0)- or $-SO_2$ -, then R^{14} is a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different;

(ii) when A^6 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo-C₂-C₆ alkenylene group, a C₂-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group, then R14 is a hydrogen atom, a halogen atom, a cyano group, a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and are selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkytthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a phenylthio group, a sub-

stituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6- alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different)));

 \underline{n} is an integer of 1 to 4;

5

10

15

20

25

30

35

40

45

50

55

further, X may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole), by combining together with the adjacent carbon atoms in the phenyl ring, and said condensed ring may have at least one substituents, which may be the same or different, and selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkytthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_2 sulfonyl group, a mono-C1-C6 alkylamino group, a di-C1-C6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 -C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo-C1-C6 alkylsulfinyl group, a halo-C1-C6 alkyl sulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C2-C6 alkenyl group, a C2-C6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkytthio group, a C_1 - C_6 alkytsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different;

Y is the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a di- C_1 - C_6 alkoxyphosphoryl group which may be the

same or different, a di-C1-C6 alkoxythiophosphoryl group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1- C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a haio-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a haio-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkytthio group, a C_1 - C_6 alkytsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁-C₆ alkylamino group and a di-C₁-C₆ alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a C_2 - C_6 a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_2 - C_1 - C_2 - C_3 - C_4 - C_6 fonyl group, a halo-C1-C6 alkylsulfonyl group, a mono-C1-C6 alkylamino group and a di-C1-C6 alkylamino group which may be the same or different, or a group of the formula -A2-R7 (wherein A2 and R7 are the same as defined above);

m is an integer of 1 to 5;

10

15

20

25

30

35

40

45

50

55

further, Y may form a condensed ring (the condensed ring is the same as defined above), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituents, which may be the same or different, and selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C2-C6 alkenyl group, a halo-C₂-C₆ alkenyl group, a C₂-C₆ alkynyl group, a halo-C₂-C₆ alkynyl group, a C₁-C₆ alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkytthio group, a C_1 - C_6 alkytsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a mono-C₁- C_6 alkylamino group, a di- C_1 - C_6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a $m C_{1} ext{-}C_{6}$ alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- $\mathrm{C_1\text{-}C_6}$ alkylsuifonyl group, a mono- $\mathrm{C_1\text{-}C_6}$ alkylamino group and a di- $\mathrm{C_1\text{-}C_6}$ alkylamino group which may be the same or different, a naphthyl group, a substituted naphthyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo- C_2 - C_6 alkynyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkytthio group, a halo- C_1 - C_6 alkytthio group, a C_1 - C_6 alkytsuffinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein said heterocyclic group is the same as defined above) having at lease one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a cyano group, a nitro group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_2 - C_6 alkenyl group, a halo- C_2 - C_6 alkenyl group, a C_2 - C_6 alkynyl group, a halo-C2-C6 alkynyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_2 - C_1 - C_2 - C_3 - C_4 - C_6 -Cfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a mono- C_1 - C_6 alkylamino group and a di- C_1 - C_6 alkylamino group which may be the same or different;

 Z^1 and Z^2 are each represents an oxygen atom or a sulfur atom.

7. The agricultural and horticultural insecticides according to Claim 6,

wherein R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, or a group of the formula -A¹-Q₂ (wherein A¹ is a C_1 - C_8 alkylene group, a C_3 -

 C_6 alkenylene group or a C_3 - C_6 alkynylene group; and Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C1-C6 alkyl group, a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a C1-C6 alkoxycarbonyl group, a di- C_1 - C_6 alkoxyphosphoryl group which may be the same or different, a di- C_1 - C_6 alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolył group, oxadiazolył group, thiazolył group, isothiazolył group, thiadiazolył group, imidazolył group, triazolył group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula $-Z^3$ - R^5 (wherein Z^3 is -O-, -S-, -SO₂- or a group of the formula -N(\mathbb{R}^6)- (wherein \mathbb{R}^6 is a hydrogen atom, a \mathbb{C}_1 - \mathbb{C}_6 alkylcarbonyl group, a halo-C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenyl C₁-C₄ alkoxycarbonyl group, or a substituted phenyl C₁-C₄ alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group,); and

R⁵ is a hydrogen atom, a C₁-C₆ alkyl group, a halo C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C_1 - C_6 alkylcarbonyl group, a halo C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkytthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group); and £ is an integer of 1 to 4);

45

50

55

5

10

20

25

30

35

40

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X may be the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituents which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkysulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a halo- $C_$

10

15

20

25

30

35

40

45

50

55

alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula -A²-R² (wherein A² is -O-, -S-, -SO-, -SO₂-, -C(=O)-, -C(=NOR³)-(wherein R³ is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 alkyl group, a phenyl- C_1 - C_4 alkyl group, or a substituted phenyl- C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group), a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_1 - C_6 alkylene group, a C_2 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group;

(1) when A² is -O-, -S-, -SO- or -SO₂-, then R⁷ is a halo-C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a $\mathsf{C}_1 ext{-}\mathsf{C}_6$ alkylsulfonyl group and a halo- $\mathsf{C}_1 ext{-}\mathsf{C}_6$ alkylsulfonyl group, or a group of the formula - $\mathsf{A}^3 ext{-}\mathsf{R}^9$ (wherein A^3 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_3 - C_6 alkenylene group, a halo- C_3 - C_6 alkenylene group, a C_3 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group; R^9 is a hydrogen atom, a halogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^4 - R^{10} (wherein A^4 is -O-, -S-, -SO-, -SO₂- or -C(=O)-, and R^{10} is a $\mathsf{C}_1\text{-}\mathsf{C}_6$ alkyl group, a halo- $\mathsf{C}_1\text{-}\mathsf{C}_6$ alkyl group, a $\mathsf{C}_3\text{-}\mathsf{C}_6$ alkenyl group, a halo- $\mathsf{C}_3\text{-}\mathsf{C}_6$ alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group));

(2) when A^2 is -C(=O)- or a group of the formula -C(=NOR⁸)-(wherein R⁸ is the same as defined the above), then R^7 is a $\mathsf{C}_1 ext{-}\mathsf{C}_6$ alkyl group, a halo $\mathsf{C}_1 ext{-}\mathsf{C}_6$ alkyl group, a $\mathsf{C}_2 ext{-}\mathsf{C}_6$ alkenyl group, a halo- $\mathsf{C}_2 ext{-}\mathsf{C}_6$ alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a mono- C_1 - C_6 alkylamino group, a di- C_1 - C_6 alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkytthio group, a halo-C1-C6 alkytthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy

group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group,

(3) when A² is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, C₂-C₆ alkenylene group, a halo-C₂-C₆ alkenylene group, a C₂-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group, then R⁷ is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkythio group, a halo- C_1 - C_6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SO- or -SO₂-; and R¹² is a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above). a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkytthio group, a halo-C1-C6 alkytthio group, a C1-C6 alkytsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula $-A^6-R^{14}$ (wherein A^6 is a C_1-C_6 alkylene group, a halo- C_1-C_6 alkylene group, a C_2-C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group, or a halo- C_3 - C_6 alkynylene group; and $\rm R^{14}$ is a hydrogen atom, a halogen atom, a $\rm C_3\text{-}C_6$ cycloalkyl group, a halo- $\rm C_3\text{-}C_6$ cycloalkyl group, a $\rm C_1\text{-}$ C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁- C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alky fonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_2 - C_1 - C_2 - C_3 - C_4 - C_5 - C_6 - C_6 - C_7 - C_8 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkył group, a hało-C₁-C₆ alkył group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆- alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group)));

n is an integer of 1 to 4;

5

10

15

20

25

30

35

40

45

50

55

further, X may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituent, which may be the same or different, and is selected from the group consisting of

a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkyls

Y is the same or different, and is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^2 - A^7 (wherein A^2 and A^7 are the same as defined above);

m is an integer of 1 to 5;

5

10

15

20

25

30

35

40

45

50

55

further, Y may form a condensed ring (the same as defined above), by combining together with the adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituent, which may be the same or different, and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), and a substituted heterocyclic group (wherein said heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; and Z^1 and Z^2 are each represents an oxygen atom or a sulfur atom.

8. The agricultural and horticultural insecticides according to Claim 7, containing as the effective ingredient, a phthalic acid diamide derivative represented by the general formula (I-1),

$$\begin{array}{c|c}
X & Z^1 \\
C-N(R^1) & R^2 \\
Z^2 & Y^1
\end{array}$$
(I-1)

{wherein, R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group or a group of the formula - A^1 - Q_ℓ (wherein, A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alke-

nylene group or a C₃-C₆ alkynylene group; Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₁-C₆ alkył group, a C₃-C₆ cycloalkył group, a halo-C₃-C₆ cycloalkył group, a C₁-C₆ alkoxycarbonyl group, a $di-C_1-C_6$ alkoxyphosphoryl group which may be the same or different, a $di-C_1-C_6$ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkytthio group, a halo-C1-C6 alkytthio group, a C1-C6 alkytsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -Z³-R⁵ (wherein Z³ is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R^6)- (wherein R^6 is a hydrogen atom, a C_1 - C_6 alkylcarbonyl group, a halo-C₁-C₆ alkylcarbonyl group, a C₁-C₆ alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C6 alkylsulfonyl group and a halo-C₁-C6 alkylsulfonyl group, a phenyl C₁-C₄ alkoxycarbonyl group, or a substituted phenyl C1-C4 alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkytthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfinyl group. fonyl group); and R⁵ is a hydrogen atom, a C₁-C₆ alkyl group, a halo C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ alkynyl group, a halo- C_6 cycloalkyl group, a C_1 - C_6 alkylcarbonyl group, a halo C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group); and $\underline{\ell}$ is an integer of 1 to 4); further,

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X is a hydrogen atom or a nitro group;

5

10

15

20

25

30

35

40

45

50

55

 Y^1 and Y^3 may be the same or different and are each a hydrogen atom, a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a balo- C_1 - C_6 alkylsulfinyl group, a balo- C_1 - C_6 alkylsulfonyl group, a pyridyloxy group, a substituted pyridyloxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6

10

15

20

25

30

35

40

45

50

55

 Y^2 is a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfinyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^2 - R^7 (wherein A^2 is -O-, -S-, -SO-, -SO₂-, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_3 - C_6 alkynylene group and,

(1) when A² is -O-, -S-, -SO- or -SO₂-, then R⁷ is a halo-C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₅ alkyl group, a halo-C₁-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsultonyl group, or a group of the formula -A3-R9 (wherein A3 is a halo-C1-C6 alkylene group, a halo-C3-C6 alkenylene group, a C₃-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group; and R⁹ is a hydrogen atom, a halogen atom, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group or a group of the formula -A⁴-R¹⁰ (wherein A⁴ is -O-, -S- or -SO₂-; and R¹⁰ is a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₃-C₆ alkenyl group, a halo-C₃-C₆ alkenyl group, a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group));

(2) when A² is a halo-C₁-C₆ alkylene group, a C₂-C₆ alkenylene group, a halo-C₂-C₆ alkenylene group, a C_2 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group, then R^7 is a hydrogen atom, a halogen atom, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SOor -SO₂-; and R¹² is a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group; or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C2-C6 alkenylene group, a halo-C2-C6 alkenylene group, a C2-C6 alkynylene group or a halo-C3-C6 alkynylene group; and R14 is a hydrogen atom, a halogen atom, a halo-C3-C6 cycloalkyl group, a halo-C1-C6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group and a

halo- C_1 - C_6 alkylsulfonyl group, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 -alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group, and a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfinyl group))); further,

 Y^1 and Y^2 may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole) by combining to each other together with the adjacent Y3, said condensed ring may have at least one substituent, which is the same or different, selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_2 - C_1 - C_2 - C_3 - C_4 - C_6 sulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group; and Z^1 and Z^2 are each an oxygen atom or a sulfur atom).

9. The agricultural and horticultural insecticides according to Claim 7, containing as the active ingredient, a phthalic acid diamide derivative represented by the general formula (I-2),

$$X^{2} \xrightarrow{X^{1}} \begin{bmatrix} Z^{1} \\ \vdots \\ C-N(R^{1}) R^{2} \end{bmatrix}$$

$$Ym \qquad (I-2)$$

40

45

50

55

10

15

20

25

30

35

{wherein, R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group or a group of the formula - A^1 - Q_ℓ (wherein, A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C₃-C₆ alkynylene group; Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₁-C₆ alkyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a $di-C_1-C_6$ alkoxyphosphoryl group which may be the same or different, a $di-C_1-C_6$ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6

alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - Z^3 - Z^5 - Z^5 (wherein Z^3 is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(Z^5 -

5

10

15

20

25

30

35

45

50

55

 R^5 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C1-C6 alkylcarbonyl group, a halo-C1-C6 alkylcarbonyl group and a C1-C6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C_1 - C_4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfinyl group. fonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkytthio group, a halo-C₁-C₆ alkytthio group, a C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group); and £ is an integer of 1 to 4); further,

R¹ and R² may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom:

X1 and X2 may be the same or different and are each a halogen atom, a cyano group, a C1-C6 alkyl group, a halo-C₁-C₆ alkyi group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkyithio group, a halo-C₁- C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group; further, X¹ and X² may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole) by combining to each other, and said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a $C_1 - C_6 \text{ alkylthio group, a halo-} \\ C_1 - C_6 \text{ alkylthio group, a } C_1 - C_6 \text{ alkylsulfinyl group, a halo-} \\ C_1 - C_6 \text{ alkylsulfinyl group,$ group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; Y is the same or different, and are each a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-

5

10

15

20

25

30

35

40

45

50

55

 C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, or a group of - C_1 - C_2 alkylsulfinyl group, a C_1 - C_2 alkylsulfonyl group, a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halogen atom, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_2 alkylsulfonyl group, a case of a halo- C_1 - C_3 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsulfonyl group, a case of a halo- C_1 - C_3 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsulfonyl group, a case of a halo- C_1 - C_4 alkylsu

(1) when A² is -O-, -S-, -SO- or -SO₂-, then R⁷ is a halo-C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkenyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkythio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A³-R⁹ (wherein A^3 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_3 - C_6 alkenylene group, a halo- C_3 -C₆ alkenylene group, a C₃-C₆ alkynylene group or a halo-C₃-C₆ alkynylene group; R⁹ is a hydrogen atom, a halogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A⁴-R¹⁰ (wherein A⁴ is -O-, -S-, -SO-, -SO₂- or -C(=O)-, and R^{10} is a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group));

(2) when A^2 is -C(=O)- or a group of the formula -C(=NOR⁸)-(wherein R⁸ is the same as defined the above), then R⁷ is a C₁-C₆ alkyl group, a halo C₁-C₆ alkyl group, a C₂-C₆ alkenyl group, a halo-C₂-C₆ alkyl group, a C₁-C₆ alkyl group, a C₁-C₆ alkylamino group, a C₁-C₆ alkylamino group, a di-C₁-C₆ alkylamino group which may be the same or different, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenylamino group, a substituted phenylamino group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoy group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoy group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoy group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoy group, a halo-C₁-C₆ alkoy group, a C₁-C₆ alkoy group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoy group gr

5

10

15

20

25

30

35

40

45

50

55

alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined the above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - $C_$

(3) when A^2 is a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group or a halo- C_3 - C_6 alkynylene group, then R^7 is a hydrogen atom, a halogen atom, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SO- or -SO₂-; and R¹² is a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C₂-C₆ alkenylene group, a halo- C_2 - C_6 alkenylene group, a C_2 - C_6 alkynylene group, or a halo- C_3 - C_6 alkynylene group; and R^{14} is a hydrogen atom, a halogen atom, a $\mathrm{C}_3\text{-}\mathrm{C}_6$ cycloalkyl group, a halo- $\mathrm{C}_3\text{-}\mathrm{C}_6$ cycloalkyl group, a $\mathrm{C}_1\text{-}$ C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_2 alkylsulfonyl group, a halo- C_2 - C_3 - C_4 - C_5 fonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group, a phenylthio group, a substituted phenylthio group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆- alkylsulfonyl group, a heterocyclic group (which is the same as defined the above), or a substituted heterocyclic group (wherein the heterocyclic ring is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo- C_1 - C_6 alkylsulfonyl group))); and \underline{m} is an integer of 1 to 5;

further, Y may form a condensed ring (which is the same as defined above) by combining together with the

adjacent carbon atoms in the phenyl ring, said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl group, a C_1 - C_6 alkylsulfonyl group, a halo- C_1 - C_6 alkylsulfonyl

10. The agricultural and horticultural insecticides according to Claim 9, containing as the effective ingredient, a phthalic acid diamide derivative represented by the general formula (I-3),

$$X^{2} \xrightarrow{\begin{array}{c} X^{1} & Z^{1} \\ C-N(R^{1}) & R^{2} \end{array}} Y^{3}$$

$$C^{-N(R^{3})} \xrightarrow{\begin{array}{c} Y^{2} \\ Z^{2} \end{array}} Y^{2}$$

$$(I-3)$$

30

35

40

45

50

55

10

15

20

25

{wherein, R^1 , R^2 and R^3 may be the same or different, and are each a hydrogen atom, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group or a group of the formula - A^1 - Q_ℓ (wherein, A^1 is a C_1 - C_8 alkylene group, a C_3 - C_6 alkenylene group or a C₃-C₆ alkynylene group; Q is a hydrogen atom, a halogen atom, a cyano group, a nitro group, a halo-C₁-C₆ alkyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C₁-C₆ alkoxycarbonyl group, a $di-C_1-C_6$ alkoxyphosphoryl group which may be the same or different, a $di-C_1-C_6$ alkoxythiophosphoryl group which may be the same or different, a diphenylphosphino group, a diphenylphosphono group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C1-C6 alkoxy group, a halo-C1-C6 alkythio group, a C1-C6 alkythio group, a halo-C1-C6 alkythio group, a C1-C6 alkytsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which means pyridyl group, pyridine-N-oxide group, pyrimidinyl group, furyl group, tetrahydrofuryl group, thienyl group, tetrahydrothienyl group, tetrahydropyranyl group, tetrahydrothiopyranyl group, oxazolyl group, isoxazolyl group, oxadiazolyl group, thiazolyl group, isothiazolyl group, thiadiazolyl group, imidazolyl group, triazolyl group or pyrazolyl group), a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_2 - C_6 alkoxy group, a C_1 - C_2 - C_2 - C_3 - C_4 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfonyl group and halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - Z^3 - R^5 (wherein Z^3 is -O-, -S-, -SO-, -SO₂- or a group of the formula -N(R⁶)- (wherein R⁶ is a hydrogen atom, a C₁-C₆ alkylcarbonyl group a halo-C1-C6 alkylcarbonyl group, a C1-C6 alkoxycarbonyl group, a phenylcarbonyl group, a substituted phenylcarbonyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 - C_1 - C_6 - C_1 - C_6 - C_1 - C_1 - C_1 - C_1 - C_1 - C_1 - C_1 - C_2 - C_1 - C_2 - C_1 - C_2 - C_1 - C_2 - C_1 - C_2 - C_1 - C_2 - C_1 - C_2 - C_2 - C_2 - C_1 - C_2 - $C_$ a C1-C6 alkylthio group, a halo-C1-C6 alkylthio group, a C1-C6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkoxycarbonyl group, or a substituted phenyl C1-C4 alkoxycarbonyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group.

fonyl group); and

10

15

20

25

30

35

40

45

50

55

 R^5 is a hydrogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_3 - C_6 alkenyl group, a halo- C_3 - C_6 alkenyl group, a C₃-C₆ alkynyl group, a halo-C₃-C₆ alkynyl group, a C₃-C₆ cycloalkyl group, a halo-C₃-C₆ cycloalkyl group, a C_1 - C_6 alkylcarbonyl group, a halo- C_1 - C_6 alkylcarbonyl group, a C_1 - C_6 alkoxycarbonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C1-C6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a phenyl C1-C4 alkyl group, a substituted phenyl C1-C4 alkyl group having at least one substituent, in the phenyl ring, which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group, a heterocyclic group (which is the same as defined above), or a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group); and £ is an integer of 1 to 4); further,

 R^1 and R^2 may form a 4 to 7 membered ring by combining to each other, in which the ring may contain the same or different 1 to 3 hetero atoms selected from the group consisting of oxygen atom, sulfur atom and nitrogen atom;

X1 and X2 may be the same or different and are each a halogen atom, a cyano group, a C1-C6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group; further, X¹ and X² may form a condensed ring (which means naphthalene, tetrahydronaphthalene, indene, indane, quinoline, quinazoline, chroman, isochroman, indole, indoline, benzodioxane, benzodioxole, benzofuran, dihydrobenzofuran, benzothiophene, dihydrobenzothiophene, benzoxazole, benzothiazole, benzimidazole or indazole) by combining to each other, and said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group, a halo-C₁-C₆ alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C_1 - C_6 alkylthio group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C1-C6 alkylsulfonyl group and a halo-C1-C6 alkylsulfonyl group, a heterocyclic group (which is the same as defined above), and a substituted heterocyclic group (wherein the heterocyclic group is the same as defined above) having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfonyl group and a halo-C₁-C₆ alkylsulfonyl group; Y^1 and Y^3 may be the same or different, and are each a hydrogen atom, a halogen atom, a C_1 - C_6 alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ C₆ alkylthio group, a phenoxy group, a substituted phenoxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C₁-C₆ alkyl group, a halo-C₁-C₆ alkyl group, a C₁-C₆ alkoxy group, a halo-C₁-C₆ alkoxy group, a C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylthio group, a C₁-C₆ alkylsulfinyl group, a halo-C₁-C₆ alkylsulfinyl group, a C₁-C₆ alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a pyridyloxy group, or a substituted pyridyloxy group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group,

 Y^2 is a hydrogen atom, a halogen atom, a halo- C_3 - C_6 cycloalkyl group or a group of the formula - A^2 - B^7 (wherein A^2 -O-, -S-, -SO-, -SO₂-, a C_1 - C_6 alkylene group, a halo- C_1 - C_6 alkylene group, a C_2 - C_6 alkenylene group, a halo- C_2 - C_6 alkenylene group, and

(1) when A^2 is -O-, -S-, -SO- or -SO₂-, then R^7 is a halo- C_3 - C_6 cycloalkyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of

a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, a substituted pyridyloxy group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo- C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkylsulfinyl group and a halo- C_1 - C_6 alkylsulfonyl group, or a group of the formula - A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^9 (wherein A^3 is a halo- A^3 - A^3 - A^3 (wherein A^3 is a halo- A^3 - A^3 - A^3 (wherein A^3 is a halo- A^3 - A^3 - A^3 - A^3 (wherein A^3 -

(2) when A² is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C₂-C₆ alkenylene group, a halo-C₂- C_6 alkenylene group, a C_2 - C_6 alkynylene group, a halo- C_3 - C_6 alkynylene group, then R' is a hydrogen atom, a halogen atom, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group, or a group of the formula -A⁵-R¹² (wherein A⁵ is -O-, -S-, -SO- or -SO2-; and R12 is a C3-C6 cycloalkyl group, a halo-C3-C6 cycloalkyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, or a group of the formula -A⁶-R¹⁴ (wherein A⁶ is a C₁-C₆ alkylene group, a halo-C₁-C₆ alkylene group, a C₂-C₆ alkenylene group, a halo-C₂-C₆ alkenylene group; and R¹⁴ is a hydrogen atom, a halogen atom, a halo-C₃-C₆ cycloalkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group, a halo-C1-C6 alkylsulfonyl group, a phenyl group, a substituted phenyl group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C₁-C₆ alkyl group, a halo-C₁-C₆ alkoxy group, a halo-C₁-C₆ alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a phenoxy group, a substituted phenoxy group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C1-C6 alkylsulfinyl group and a halo-C1-C6 alkylsulfonyl group, a phenylthio group, or a substituted phenylthio group having at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a halo-C1-C6 alkyl group, a halo-C1-C6 alkoxy group, a halo-C1-C6 alkylthio group, a halo-C₁-C₆ alkylsulfinyl group and a halo-C₁-C₆ alkylsulfonyl group));

further, Y¹ and Y² may form a condensed ring (the condensed ring is the same as defined above) by combining to each other together with Y³, and said condensed ring may have at least one substituent, which is the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkoxy group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a phenyl group, and a substituted phenyl group having at least one substituent which may be the same or different and is selected from the group consisting of a halogen atom, a C_1 - C_6 alkyl group, a halo- C_1 - C_6 alkyl group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylthio group, a C_1 - C_6 alkylsulfinyl group, a halo- C_1 - C_6 alkylsulfinyl group, a ha

 Z^1 and Z^2 are each an oxygen atom or a sulfur atom).

11. A method for controlling undesirable insect pests for a useful crop, characterized by treating an objective crop with an effective amount of the agricultural and horticultural insecticides as claimed in any one of Claims 6 to 10.

55

5

10

15

20

25

30

35

40

45

50



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 0 919 542 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3: 12.04.2000 Bulletin 2000/15

(43) Date of publication A2: 02.06.1999 Bulletin 1999/22

(21) Application number: 98122107.0

(22) Date of filing: 23.11.1998

(51) Int. CI.⁷: **C07C 233/64**, C07C 235/42, C07C 317/32, C07C 323/42, C07D 295/192, A01N 37/18, A01N 43/00, A01N 43/84, A01N 57/00

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE Designated Extension States: AL LT LV MK RO SI

(30) Priority: 25.11.1997 JP 33939397

(71) Applicant:
NIHON NOHYAKU CO., LTD.
Chuo-ku Tokyo100-0027 (JP)

(72) Inventors:

Tohnishi, Masanori
 Sakai-shi (JP)

Nakao, Hayami
 Kawachinagano-shi (JP)

• Kohno, Eiji Habikino-shi (JP)

 Nishida, Tateki Tondabayashi-shi (JP) Furuya, Takashi Izumisano-shi (JP)

 Shimizu, Toshiaki Kawachinagano-shi (JP)

Seo, Akira
 Hashimoto-shi (JP)

 Sakata, Kazuyuki Kawachinagano-shi (JP)

• Fujioka, Shinsuke Kawachinagano-shi (JP)

 Kanno, Hideo Ibaraki-shi (JP)

(74) Representative:

Grünecker, Kinkeldey, Stockmair & Schwanhäusser Anwaltssozietät Maximilianstrasse 58 80538 München (DE)

- (54) Phthalic acid diamide derivatives, agricultural and horticultural insecticides, and a method for application of the insecticides
- (57) The present invention provides a phthalic acid diamide derivative represented by the general formula (I),

$$\begin{array}{c|c}
X_{n} & Z_{1} \\
C-N(R^{1}) R^{2} \\
C-N(R^{2}) & Y_{m}
\end{array}$$
(1)

{wherein R^1 , R^2 and R^3 may be same or different, and are each a hydrogen atom, a cyano group, a C_3 - C_6 cycloalkyl group, a halo- C_3 - C_6 cycloalkyl group, a C_3 - C_6 cycloalkenyl group, a halo- C_3 - C_6 cycloalkenyl group, or a group of the formula $-A^1$ - Q_ℓ ; X may be the same or different and are each a hydrogen atom, a halogen atom, a cyano group, a nitro group, a phenyl group, a substituted phenyl group, a naphthyl group, a substituted naphthyl group, a heterocyclic group, a substituted heterocyclic group; or a group of the formula $-A^2$ - R^7 ; n is an integer of 1 to 4; Y may be same or different and are each a hydrogen atom, a halogen atom, a cyano group, a nitro group, a phenyl group, a substituted phenyl group, a naphthyl group, a substituted naphthyl group, a heterocyclic group, a substituted heterocyclic group or a group

of the formula $-A^2-R^7$; \underline{m} is an integer of 1 to 5; Z^1 and Z^2 are each an oxygen atom or a sulfur atom}, and an agricultural and horticultural insecticide containing said phthaldiamide derivative, as well as to provide a method for use of said insecticide.

The agricultural and horticultural insecticides of the present invention show excellent activities for controlling injurious insects.



Application Number

which under Rule 45 of the European Patent Convention EP 98 12 2107 shall be considered, for the purposes of subsequent proceedings, as the European search report

		ERED TO BE RELEVANT	1	
Category	Citation of document with in of relevant passe	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.6)	
х	DE 38 02 175 A (HOE 3 August 1989 (1989 * page 2 - page 7;	-08-03)	1-11	C07C233/64 C07C235/42 C07C317/32 C07C323/42
X,D	PATENT ABSTRACTS OF vol. 011, no. 003 (7 January 1987 (198 & JP 61 180753 A (N 13 August 1986 (198 * abstract *	C-395), 7-01-07) IPPON KAYAKU CO LTD),	1	C07D295/192 A01N37/18 A01N43/00 A01N43/84 A01N57/00
X D	EP 0 119 428 A (BAY 26 September 1984 (* page 28; claim 1 & JP 59 163353 A	1984-09-26)	1	
X	PATENT ABSTRACTS OF vol. 015, no. 466 (26 November 1991 (1 & JP 03 198049 A (F 29 August 1991 (199 * abstract *	P-1280), 991-11-26) UJI PHOTO FILM CO LTD),	1	TECHNICAL FIELDS SEARCHED (Inl.Cl.6)
	abstract -	-/		C07C C07D A01N
INCO	MPLETE SEARCH			
not comp be carried Claims se	ly with the EPC to such an extent that d out, or can only be carried out partial earched completely;	application, or one or more of its claims, does a meaningful search into the state of the art or ly, for these claims.	/do annot	
	earched incompletely :			
,,	or the limitation of the search: sheet C			
			·	
	Place of search	Date of completion of the search	<u> </u>	Examiner
	BERLIN	20 January 2000	Rut	et, J
X : par Y : par doo	ATEGORY OF CITED DOCUMENTS ticularly relevant if taken alone ticularly relevant if combined with anciument of the same category	T : theory or principle E : earlier patent doc after the filing dab ther D : document citled for L : document citled for	oument, but publi o n the application	
O : nor	hnological background n-written disclosure ermediate document	& : member of the sa document		

SACO ECON 1500 CO



PARTIAL EUROPEAN SEARCH REPORT Application Number

	DOCUMENTS CONSIDERED TO BE RELEVANT		CLASSIFICATION OF THE APPLICATION (Int.CI.6)	
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim		
х .	CHEMICAL ABSTRACTS, vol. 124, no. 6, 5 February 1996 (1996-02-05) Columbus, Ohio, US; abstract no. 56894r, PERRY ROBERT J. ET AL.: "Polyimide formation through the palladium-mediated carbonylation" page 12; column 1; XP002095523 * abstract * & MACROMOLECULES, vol. 29, no. 3, 1996, pages 1014-1020,	1		
X	CHEMICAL ABSTRACTS, vol. 123, no. 26, 25 December 1995 (1995-12-25) Columbus, Ohio, US; abstract no. 343359e, HALL, NIGEL: page 180; column 2; XP002095524 * abstract * & WO 95 20014 A (ZENECA LTD)	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6)	
	CHEMICAL ABSTRACTS, vol. 120, no. 21, 23 May 1994 (1994-05-23) Columbus, Ohio, US; abstract no. 269853f, BEELY, NIGEL ET AL.: page 1012; column 1; XP002095525 * abstract * & WO 93 25517 A (CELLTECH LTD)			
			*	



Application Number

	DOCUMENTS CONSIDERED TO BE RELEVANT		CLASSIFICATION OF THE APPLICATION (Int.CI.6)	
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim		
x	CHEMICAL ABSTRACTS, vol. 113, no. 19, 5 November 1990 (1990-11-05) Columbus, Ohio, US; abstract no. 171822a, ISMAIL, M. FEKRY ET AL.: "Reaction of N-arylphthalisoimidium perchlorates with amines" page 686; column 2; XP002095526 * abstract * & ZEITSCHRIFT FUR NATURFORSCHUNG, TEIL B: CHEM. SCI., vol. 45, no. 5, 1990, pages 707-710,	1		
х	CHEMICAL ABSTRACTS, vol. 109, no. 5, 1 August 1988 (1988-08-01) Columbus, Ohio, US; abstract no. 37697c, GANIN, E. V. ET AL.: "N-substituted isophthalimide formation." page 604; column 1; XP002095527 * abstract * & UKR. KHIM. ZH., vol. 53, no. 9, 1987, pages 964-967,	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6)	
X	CHEMICAL ABSTRACTS, vol. 91, no. 9, 27 August 1979 (1979-08-27) Columbus, Ohio, US; abstract no. 74313e, DE SILVA, S. OSMUND ET AL.: "Directed lithiation" page 569; column 1; XP002095528 * abstract * & TETRAHEDRON LETTERS., vol. 51, 1978, pages 5099-5102, OXFORD GB	1		
			·	



Application Number

	DOCUMENTS CONSIDERED TO BE RELEVANT	CLASSIFICATION OF THE APPLICATION (InlCI.6)	
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	CHEMICAL ABSTRACTS, vol. 68, no. 3, 15 January 1968 (1968-01-15) Columbus, Ohio, US; abstract no. 12654h, EDITH G. DIAZ DE TORANZO ET AL.: "Syntheses of unsymmetric o-phthalic diamides." page 1198; column 2; XP002095529 * abstract * & JOURNAL OF MEDICINAL CHEMISTRY., vol. 10, no. 5, 1967, pages 982-983, WASHINGTON US	1	
X	CHEMICAL ABSTRACTS, vol. 70, no. 15, 14 April 1969 (1969-04-14) Columbus, Ohio, US; abstract no. 68102k, RESPLANDY, ALBERT ET AL.: "Synthesis of 6-phenanthridone-7-carboxylic acids from N-arylphthalamic acids." page 353; column 1; XP002095530 * abstract * & BULLETIN DE LA SOCIETE CHIMIQUE DE FRANCE., vol. 12, 1968, pages 4947-4953, PARIS FR	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
x	US 4 732 845 A (KEIJI MIHAYASHI ET AL) 22 March 1988 (1988-03-22) * column 10-16; claim 1 *	1	
X	US 4 694 016 A (LU JING-JONG ET AL) 15 September 1987 (1987-09-15) * the whole document */	1	



Application Number

	DOCUMENTS CONSIDERED TO BE RELEVANT	CLASSIFICATION OF THE APPLICATION (Int.CI.6)	
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	US 3 502 685 A (GEVIRTZ ARTHUR H ET AL) 24 March 1970 (1970-03-24) * the whole document *	1	
X	CHEMICAL ABSTRACTS, vol. 126, no. 21, 26 May 1997 (1997-05-26) Columbus, Ohio, US; abstract no. 277494d, page 673; column 1; XP002128153 * abstract * & JP 09 059236 A (DAIICHI SEIYAKU)	1	·
X	CHEMICAL ABSTRACTS, vol. 124, no. 3, 15 January 1996 (1996-01-15) Columbus, Ohio, US; abstract no. 29077v, ISMAIL, M. FEKRY ET AL.: "Studies on the mode of ring-opening of N-aryl phthalisoimidium perchlorates." page 913; column 2; XP002128154 * abstract * & POL. J. CHEM., vol. 69, no. 8, 1995, pages 1114-1119, -/	1	TECHNICAL FIELDS SEARCHED (Int.CI.6)



Application Number

	DOCUMENTS CONSIDERED TO BE RELEVANT		CLASSIFICATION OF THE APPLICATION (Int.CI.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
x	CHEMICAL ABSTRACTS, vol. 122, no. 19, 8 May 1995 (1995-05-08) Columbus, Ohio, US; abstract no. 239480f, RANADIVE, V. B. ET AL.: "Nucleophilic reactions of N-hydroxy-,methoxy-, 2,3-epoxypropoxy-phthalimides" page 1039; column 1; XP002128155 * abstract * & INDIAN J. CHEM., SECT. B: ORG. CHEM. INCL. MED. CHEM., vol. 33B, no. 12, 1994, pages 1175-1177,	1	
X	CHEMICAL ABSTRACTS, vol. 108, no. 1, 4 January 1988 (1988-01-04) Columbus, Ohio, US; abstract no. 5636v, CREMLYN, R. J. ET AL.: "Phthalimidobenzenesulfonyl derivatives" page 533; column 1; XP002128156 * abstract * & QUIM. NOVA, vol. 8, no. 1, 1985, pages 61-62,	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	PATENT ABSTRACTS OF JAPAN vol. 009, no. 119 (C-282), 23 May 1985 (1985-05-23) & JP 60 008247 A (SHOWA DENKO KK), 17 January 1985 (1985-01-17) * abstract *	1	-



Application Number

	DOCUMENTS CONSIDERED TO BE RELEVANT	CLASSIFICATION OF THE APPLICATION (Int.CI.6)	
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	CHEMICAL ABSTRACTS, vol. 79, no. 23, 10 December 1973 (1973-12-10) Columbus, Ohio, US; abstract no. 136746m, IMAI, YOSHIO ET AL.: "Reaction of N-(phenylsulfonyl)phthalimde" page 347; column 2; XP002128157 * abstract * & NIPPON KAGAKU KAISI, vol. 9, 1973, pages 1729-1733,	1	
X	CHEMICAL ABSTRACTS, vol. 57, no. 7, 1 October 1962 (1962-10-01) Columbus, Ohio, US; abstract no. 8530g, WALTER RIED ET AL.: "Monoaddition of acetylene to N-substituted phthalimides" page 1962; column 2; XP002128158 * abstract * & BER., vol. 95, 1962, pages 1562-1564,		TECHNICAL FIELDS SEARCHED (Int.Cl.6)



Application Number

CLAIMS INCURRING FEES
The present European patent application comprised at the time of filing more than ten claims.
Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):
No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.
LACK OF UNITY OF INVENTION
The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:
see sheet B
All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.
Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
1-11
None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



LACK OF UNITY OF INVENTION SHEET B

Application Number

EP 98 12 2107

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims: 1 - 11 (partly)

compounds of formula (I) wherein at least one of R1, R2 is hydrogen, C1-C8 alkyl, C3-C6 alkenyl, C3-C6 alkynyl, unsubstituted or halo-substituted C3-C6 cycloalkyl and unsubstituted or halo-substituted C3-C6 cycloalkenyl.

2. Claims: 1 - 11

compounds according to formula (I) wherein at least one of R1 or R2 = cyano

3. Claims: 1 - 11

compounds according to formula (I) wherein at least one of R1 or R2 = A1-Q(1) wherein A1 is -0- or -S-.

4. Claims: 1 - 11

compounds according to formula (I) wherein at least one of R1 or R2 = A1-Q(1) wherein A1 is -S02-.

5. Claims: 1 - 11

compounds according to formula (I) wherein at least one of R1 or R2 = A1-Q(1) wherein A1 is -C(=0)-.

6. Claims: 1 - 11

compounds according to formula (I) wherein at least one of R1 or R2 = A1-Q(1) wherein A1 is -N(R4)-.

7. Claims: 1 - 11

compounds according to formula (I) wherein at least one of R1 or R2 = -(CH2)n-X; wherein n is not 0 and X = aryl, heteroaryl or a phosphore derivative.

8. Claims: 1 - 11

compounds according to formula (I) wherein R1 and R2 together form a cycle

9. Claims: 1 -11



LACK OF UNITY OF INVENTION SHEET B

Application Number

EP 98 12 2107

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely: compounds according to formula (I) wherein at least one of R1 or R2 = heterocycle



INCOMPLETE SEARCH SHEET C

Application Number EP 98 12 2107

	J .	SHEE	1 C	£1 70	12 210,
	Claim(s) searched incomple 1-11	etely:			
	Reason for the limitation	of the sear	ch:		
	Due to the 5 disclaimers substituents in the main of the application is not therefore mainly based on	claim as wel regarded as	l as the dependent clear. The search	claims, the	the scope
	•				
				ė,	
ŀ					

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 98 12 2107

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-01-2000

Patent document cited in search repor	t	Publication date	Patent family member(s)	Publication date
DE 3802175	A	03-08-1989	CN 1037143 A EP 0325983 A JP 2001441 A PT 89542 A	15-11-19 02-08-19 05-01-19 04-10-19
JP 61180753	Α	13-08-1986	NONE	
EP 0119428	A	26-09-1984	DE 3305569 A DK 76684 A ES 529686 A ES 538151 A JP 59163353 A ZA 8401183 A	23-08-19 19-08-19 01-04-19 01-11-19 14-09-19 31-10-19
JP 03198049	Α	29-08-1991	NONE	`.
WO 9520014	A	27-07-1995	AU 1323295 A CA 2180448 A EP 0740690 A JP 9507693 T US 5739299 A ZA 9500311 A	08-08-19 27-07-19 06-11-19 05-08-19 14-04-19 28-11-19
WO 9325517	A	23-12-1993	AT 150447 T AU 670949 B AU 4347093 A CA 2114114 A DE 69309030 D DE 69309030 T DK 607373 T EP 0607373 A ES 2102036 T GR 3023446 T JP 6509820 T US 5550137 A HU 9500453 A US 5340827 A	15-04-19 08-08-19 04-01-19 23-12-19 24-04-19 25-09-19 13-10-19 27-07-19 16-07-19 29-08-19 02-11-19 27-08-19 30-10-19 23-08-19
US 4732845	A	22-03-1988	JP 2107437 C JP 7107601 B JP 62103638 A JP 2019681 C JP 7045449 B JP 62103053 A	06-11-19 15-11-19 14-05-19 19-02-19 17-05-19 13-05-19
US 4694016	Α	15-09-1987	NONE sean Patent Office, No. 12/82	

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 98 12 2107

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-01-2000

Patent document cited in search report	t	Publication date	Patent family member(s)	Publication date
US 3502685	A	24-03-1970	BE 721984 A CA 933161 A DE 1793490 A FR 1589007 A GB 1224066 A LU 57114 A NL 6813843 A US 3549348 A	08-04-196 04-09-197 24-02-197 16-03-197 03-03-197 16-04-197 22-04-196 22-12-197
JP 09059236	Α	04-03-1997	NONE	
JP 60008247	Α	17-01-1985	NONE	
				•
				٠

o For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
□ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
□ FADED TEXT OR DRAWING
□ BLURRED OR ILLEGIBLE TEXT OR DRAWING
□ SKEWED/SLANTED IMAGES
□ COLOR OR BLACK AND WHITE PHOTOGRAPHS
□ GRAY SCALE DOCUMENTS
□ LINES OR MARKS ON ORIGINAL DOCUMENT
□ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

☐ OTHER: ____

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.